


Clemson University



3 1604 019 891 516



I 19.3:168

DEPARTMENT OF THE INTERIOR


BULLETIN

OF THE

UNITED STATES

GEOLOGICAL SURVEY

No. 168



ANALYSES OF ROCKS, LABORATORY OF THE U. S. GEOLOGICAL
SURVEY, 1880 TO 1899.—CLARKE

WASHINGTON
GOVERNMENT PRINTING OFFICE
1900

Joseph Hyde Pratt
DEPARTMENT OF THE INTERIOR


BULLETIN

OF THE

UNITED STATES

GEOLOGICAL SURVEY

No. 168



WASHINGTON
GOVERNMENT PRINTING OFFICE
1900

UNITED STATES GEOLOGICAL SURVEY

CHARLES D. WALCOTT, DIRECTOR

ANALYSES OF ROCKS

FROM THE


LABORATORY OF THE UNITED STATES GEOLOGICAL SURVEY

1880-1899

TABULATED BY

F. W. CLARKE

CHIEF CHEMIST



WASHINGTON

GOVERNMENT PRINTING OFFICE

1900



Digitized by the Internet Archive
in 2013

CONTENTS.

	Page.
Introduction	11
Analyses.....	19
Igneous and crystalline rocks	19
Maine	19
1. Rocks from Aroostook County.....	19
2. Litchfieldite and associated minerals	21
New Hampshire.....	23
Vermont.....	24
1. Rocks of Mount Ascutney	24
2. Miscellaneous rocks.....	27
Massachusetts	28
1. Rocks of the Connecticut Valley.....	28
2. Miscellaneous rocks	33
Connecticut	35
New York	36
1. Rocks of the Adirondack region	36
2. Peridotite from near Syracuse	38
New Jersey	39
Pennsylvania	40
Delaware	41
Maryland, District of Columbia.....	42
1. Peridotite and pyroxenite	42
2. Gabbro and diorite	44
3. Granite and gneiss	46
4. Miscellaneous rocks.....	50
Virginia	51
North Carolina.....	52
Georgia.....	55
Kentucky.....	56
1. Elliott County dike	56
2. Crittenden County dike	58
Tennessee.....	59
Missouri	59
Arkansas	60
Texas.....	60
Lake Superior region	64
1. Marquette region, Michigan.....	64
2. Crystal Falls district, Michigan	66
3. Menominee River.....	70
First. Sturgeon Falls gabbro	70
Second. Lower Quinnesec Falls	71
Third. Upper Quinnesec Falls.....	72
Fourth. Schist from the Aragon iron mine	72

Analyses—Continued.

Page.

Igneous and crystalline rocks—Continued.

Lake Superior region—Continued.

4. Penokee-Gogebic region	73
5. Rocks of Pigeon Point, Minnesota	76
6. Miscellaneous rocks from Minnesota	81

South Dakota	84
--------------------	----

Wyoming	85
---------------	----

Yellowstone National Park and the Absaroka Range	87
--	----

1. Electric Peak	87
2. Sepulchre Mountain	90
3. Absaroka Range	92
First. Crandall Basin	92
Second. Sunlight intrusives	95
Third. Ishawooa intrusives	96
Fourth. Dikes in breccia	97
4. Absarokite-shoshonite-banakite series	99
5. Miscellaneous rocks	104

Montana	111
---------------	-----

1. Madison and Gallatin valleys	111
2. Butte and Boulder districts	116
3. Crazy Mountains	120
4. Little Belt Mountains	124
5. Castle Mountain district	129
6. Highwood Mountains	131
7. Little Rocky Mountains	134
8. Bearpaw Mountains	135

Idaho	137
-------------	-----

Colorado	140
----------------	-----

1. Denver Basin	140
2. Pikes Peak district	141
3. Silver Cliff and Rosita	147
4. Buffalo Peaks	153
5. Leadville region	154
6. Tenmile district	157
7. Elk Mountains	159
8. West Elk Mountains	160
9. San Juan region	161
10. Miscellaneous rocks	164

Utah	166
------------	-----

1. Rocks from the Tintic district	166
2. Rocks from the Henry Mountains	167
3. Miscellaneous rocks	168

New Mexico	169
------------------	-----

1. Basalts from the Rio Grande Canyon	169
2. Mount Taylor region	170
3. Colfax County	171
4. Miscellaneous rocks	172

Arizona	173
---------------	-----

Nevada	174
--------------	-----

1. Rocks from Washoe	174
2. Rocks from Eureka	175

California	176
------------------	-----

1. Mount Shasta	176
-----------------------	-----

Analyses—Continued.

Page.

Igneous and crystalline rocks—Continued.

California—Continued.

2. Lassen Peak region	177
First. Rhyolites	178
Second. Dacites and andesites	179
Third. Basalts	184
3. Plumas County	187
4. Butte County	190
5. Sierra County	192
6. Nevada City and Grass Valley	194
7. Placer County	197
8. Eldorado County	199
9. Amador County	200
10. Calaveras County	203
11. Tuolumne County	203
12. Mariposa County	207
13. The quicksilver region	211
14. Mount Diablo	213
15. Miscellaneous rocks	217

Oregon

1. Basalt, Mount Thielsen	220
2. Peridotite, Douglas County	221
3. Crater Lake	221
4. Rock from Wilbur, Douglas County	223

Washington

Alaska

Brazil

Stony meteorites

1. Rockwood, Tennessee	231
2. Hamblen County, Tennessee	232
3. Winnebago County, Illinois	233
4. Taney County, Missouri	234
5. Washington County, Kansas	234
6. Kiowa County, Kansas	235
7. Travis County, Texas	236
8. Bluff, Fayette County, Texas	237
9. San Bernardino County, California	238
10. Beaver Creek, British Columbia	239
11. Llano del Inca, Chile	240

Meteoric iron

Sandstones, cherts, and sinters

Ohio

California

1. Mount Diablo	245
2. Shasta County	246
3. Sulphur Bank	247

Miscellaneous sandstones

Chert

Siliceous sinters

Carbonate rocks

Vermont, Massachusetts

New York, Pennsylvania, Maryland

Virginia, West Virginia

Analyses—Continued.

Carbonate rocks—Continued.

	Page.
Georgia.....	255
Florida.....	255
1. Coral and shell rocks.....	255
2. Miscellaneous rocks.....	257
Tennessee, Alabama, Louisiana.....	258
Texas.....	259
Ohio.....	259
Indiana.....	262
Missouri.....	263
Kansas.....	263
Michigan, Wisconsin, Minnesota, Canada.....	264
The Yellowstone National Park.....	268
Montana.....	269
Colorado.....	270
1. Denver Basin.....	270
2. Leadville district.....	271
3. Fairplay, Park County.....	272
4. Pitkin County.....	272
5. Glenwood Springs.....	273
6. Tenmile district.....	274
Utah.....	275
Nevada and California.....	276
Hawaiian Islands.....	277
Slates and shales.....	278
Vermont.....	278
New York.....	281
Kentucky, Georgia, Alabama.....	285
Ohio.....	284
Michigan.....	284
Wisconsin.....	285
Colorado.....	286
California.....	287
Clays, soils, etc.....	289
Massachusetts.....	289
New York, Pennsylvania, Delaware.....	290
Maryland.....	291
Virginia, North Carolina.....	292
South Carolina, Georgia.....	293
Florida.....	293
Alabama, Mississippi.....	295
Illinois, Iowa, Minnesota.....	296
Wisconsin.....	297
Missouri, Arkansas.....	298
Colorado.....	299
Wyoming, Utah, New Mexico.....	302
Nevada.....	303
California, Washington, Hawaiian Islands.....	304
Index to rocks.....	305

LETTER OF TRANSMITTAL.

DEPARTMENT OF THE INTERIOR,
UNITED STATES GEOLOGICAL SURVEY,
Washington, D. C., February 27, 1900.

SIR: I submit herewith the manuscript for a revised and enlarged edition of Bulletin No. 148, entitled Analyses of Rocks, etc. Dr. Hillebrand's portion of the former bulletin is so much extended, and represents so distinct a demand, that it will be presented by him as a separate bulletin.

Very respectfully,

F. W. CLARKE,
Chief Chemist.

HON. CHARLES D. WALCOTT,
Director United States Geological Survey.

ANALYSES OF ROCKS, LABORATORY OF THE UNITED STATES GEOLOGICAL SURVEY, 1880-1899.

By F. W. CLARKE.

INTRODUCTION.

The present Geological Survey of the United States was organized in 1879. In 1880 a chemical laboratory was established at Denver, in connection with the Colorado work, in charge of Dr. W. F. Hillebrand, with whom were associated Mr. Antony Guyard and, later, Mr. L. G. Eakins. In 1882 Dr. W. H. Melville was placed in charge of a second laboratory at San Francisco, and in the autumn of 1883 the central laboratory was started in Washington, with myself as chief chemist. In November, 1885, Dr. Hillebrand was transferred to Washington; early in 1888 he was followed by Mr. Eakins, and the Denver laboratory was discontinued. In the spring of 1890 Dr. Melville also was transferred to Washington, and since then the chemical work of the Survey has been concentrated at headquarters.

Up to January 1, 1900, nearly 4,400 analyses have been made in the laboratory at Washington. These represent rocks, minerals, ores, waters, sediments, coals, metals, and so on through all the range of substances with which geology has to do. There were also some hundreds of analyses made in the laboratories at Denver and San Francisco. A fair amount of research work upon mineralogical and analytical problems has also been done. In all of this work the following chemists have been employed: Charles Catlett, T. M. Chatard, F. W. Clarke, L. G. Eakins, F. A. Gooch, Antony Guyard, W. F. Hillebrand, W. H. Melville, R. B. Riggs, E. A. Schneider, George Steiger, H. N. Stokes, William Valentine, and J. E. Whitfield. As many as eight of these have been at work simultaneously; at present only four are connected with the Survey. Other officers of the Survey have been occupied more or less with chemical questions; but the men named in this list were connected directly with the laboratory. Some work for the chemical division has also been done by chemists not regularly on the rolls of the Survey; but their analyses, with the exception of a single group to be noted later, do not fall within the scope of this paper.

Quite naturally, on account of the activity of the petrographers, the dominant feature of the laboratory work has been the analysis of rocks. These have been studied in great numbers and in the most thorough way. The results have appeared in widely scattered publications,

official reports, monographs, bulletins, American and foreign journals, and so on. The object of this bulletin is to bring together this valuable material, together with such bibliographic and petrographic data as seems to be necessary in order to identify the specimens and to facilitate chemical discussion. Analyses of minerals have been included only when related to petrographic studies, appearing then in connection with the rocks to which they belong. Meteorites, of which twenty-seven have been analyzed, are brought into the work on account of their petrographic relations; and the groups of clays and soils have been admitted because of the bearing of these substances upon the study of slates and shales. The actual number of analyses given in the bulletin is as follows:

Igneous and crystalline rocks.....	854
Mineral separations.....	90
Meteorites, and separations from them.....	58
Sandstones, cherts, and sinters.....	48
Carbonate rocks.....	208
Slates and shales	48
Clays, soils, etc.....	98
Total.....	1,404

It may be observed that the classification thus indicated has not been rigorously followed. In a few instances the study of a sedimentary rock has been so related to that of its igneous neighbors that the analyses are best tabulated together; but these exceptional cases are few, and all are properly noted. The heading "igneous and crystalline rocks" has been used in the broadest and most liberal way, and doubtless many of the analyses given under it might properly be otherwise classified. In such cases of uncertainty, convenience has furnished the rule to follow.

Within each division of the analyses the classification chosen has been geographic. The petrographic grouping of the rocks would doubtless be best were petrographers agreed upon it; but their differences are many, and the chemist will do well to avoid them. The geographic method, moreover, has some advantages of its own; it facilitates the study of areas, it simplifies the bibliographic references, and it brings together, in great measure, the work of each petrographer for whom analyses have been made. Thus, most of Diller's work has been in California, most of Cross's in Colorado, and most of Iddings's in the Yellowstone National Park, and in each case the analyses are massed, and their discussion is practically uniform in character. As regards nomenclature, each rock has received the designation given it by its describer, and no liberties have been taken. This plan may cause some lack of uniformity; but no other procedure seemed to be practicable.

It will be noticed by anyone who uses this bulletin that the analyses vary as regards completeness. Among the sedimentary rocks, espe-

cially, partial analyses are common; but in the igneous group thoroughness is more general. In the early days of the chemical division many analyses were made along the older lines, just as they are still made in many laboratories to-day—that is, only the main constituents, those having direct petrographic significance, were determined. In such analyses the minor ingredients, like titanium, phosphorus, barium, strontium, chlorine, etc., were ignored; and, although the results are satisfactory in some respects, they leave much to be desired. Latterly, greater completeness has been sought for, the work done has been much fuller, and the data obtained can be discussed with much higher approaches to accuracy. The old form of “complete analysis” is to be discouraged; it leads too often to erroneous conclusions; and only the best modern methods of work and of statement should be tolerated. The fuller analyses, moreover, have brought some interesting points to light; titanium now appears to be one of the more abundant elements, and barium and strontium are found to be almost universally diffused in igneous rocks in quite perceptible quantities.

On general principles the analysis of a rock and its petrographic description should be two parts of the same investigation, matching each other completely. In practice, however, this rule does not always hold, and the departures from it are in two opposite directions. For example, an analysis of the older type says nothing of titanium and phosphorus, while the microscope reveals the presence of sphene and apatite. In this case the petrographer has been more thorough than the chemist. On the other hand, a full and perfect analysis may be given, accompanied by a petrographic description of the most general kind, in which only the main mineral constituents of the rock are noted. Here the analysis has been incompletely used, and the petrographic discussion is defective. It is hoped that the publication of this material may lead to a clearer recognition of the mutuality which should exist between the chemical and the microscopic researches, and so bring, in the future, both lines of investigation more into harmony. Hitherto the chemist and the petrographer have worked too much apart, and each has too often misunderstood the purpose of the other. If the study of the thin section could always precede the analysis, the petrographic problems could be stated more clearly, and the chemical evidence might be rendered much more pertinent and satisfactory.

In a paper published some years ago,¹ on the relative abundance of the chemical elements, I computed the average composition of the primitive crust of the earth from 880 analyses of eruptive and crystalline rocks. Of these analyses only 207 were from the laboratories of the survey, while 673 were collected from various other American and foreign sources. A large proportion of them were incomplete, regarded from a modern point of view, and yet the results obtained were fairly conclusive. The material now available for similar discussion is much

¹ Bull. U. S. Geol. Survey No. 78, 1891, p. 34.

better than that which was formerly used, and an average based upon it may not be out of place here.

In the bulletin now presented there are 830 complete analyses of rocks which are suitable for my purpose. I have also taken from the partial analyses given in the laboratory records 180 additional determinations of silica, 90 of lime, and 130 of alkalies. In 490 of the analyses there is discrimination between the water lost below 110° and that which is essential to the composition of the rocks; and this amounts to 0.40 per cent. Omitting this water, the average found may fairly represent the composition of the older crust of the earth, as deduced from a mass of data which are reasonably uniform in character and entitled to a high degree of credence. The mean for the more important constituents is as follows, with the old average given in a parallel column for comparison:

	New mean.	Old mean.
SiO ₂	59.71	58.59
Al ₂ O ₃	15.41	15.04
Fe ₂ O ₃	2.63	3.94
FeO.....	3.52	3.48
CaO.....	4.90	5.29
MgO.....	4.36	4.49
K ₂ O.....	2.80	2.90
Na ₂ O.....	3.55	3.20
H ₂ O.....	1.52	^a 1.96
TiO ₂60	.55
P ₂ O ₅22	.22
	99.22	99.66

^a Including hygroscopic water; probably 0.40 per cent.

If we reduce these figures to elementary form and include the minor constituents which are frequently found in rocks, and which in this laboratory are often estimated, the two averages compare as follows:

	New mean.	Old mean.
Oxygen	47.02	47.29
Silicon	28.06	27.21
Aluminum.....	8.16	7.81
Iron	4.64	5.46
Calcium	3.50	3.77
Magnesium	2.62	2.68
Sodium.....	2.63	2.36
Potassium	2.32	2.40
Titanium41	.33
Hydrogen17	.21
Carbon12	.22
Phosphorus.....	.09	.10
Manganese07	.08
Sulphur07	.03
Barium.....	.05	.03
Strontium02
Chromium.....	.01	.01
Nickel.....	.01
Lithium01
Chlorine01	.01
Fluorine.....	.01
	100.00	100.00

As the old mean represents an attempt to measure the composition of the entire solid crust of the earth, and so includes an allowance for the carbon in the limestones, the two columns are not strictly comparable. They are, however, corroborative of each other, and show that within reasonable limits the statistical method is applicable to the problem under consideration. For the arguments upon which the discussion is based the original paper should be consulted. The distribution of the rarer elements has also been elaborately discussed by Vogt,¹ and their percentages may be regarded as small corrections to be applied to the table at some future time.

By a similar statistical process I have attempted to ascertain something with regard to the relative abundance of the more important rock-forming minerals. Nearly 500 analyses of igneous rocks were

¹Zeitsch. Prakt. Geologie, 1898, pp. 225, 314, 377, 413; and 1899, pp. 10, 274.

discussed, and the subjoined percentages, which are probably nothing more than rough approximations to the truth, were obtained:

Quartz	12.0
Feldspars	60.0
Pyroxenes and amphiboles	18.0
Micas	4.0
	<hr/>
	94.0

The less frequent minerals make up the remaining 6 per cent. The computation, although it is by no means conclusive, is perhaps a little more satisfactory than any similar estimate which has so far been made.

For computing the average composition of the sedimentary rocks the existing analyses of individual samples are inadequate. They are too few and too incomplete to yield any conclusions of value. Attempts have been made to partly use the data, as, for example, by Joly;¹ and it seems probable, therefore, that better material will not be without interest or scientific value.

Some five years ago, at the request of Mr. G. K. Gilbert, a series of composite analyses of sedimentary rocks was made in this laboratory. Many samples were mixed into one uniform sample, from which, by a single analysis, an average composition was determined. The material was selected and the samples were prepared by Mr. Gilbert, assisted by Mr. G. W. Stose, and the analyses were made by Dr. H. N. Stokes. The data obtained may be tabulated as follows:

- A. Composite analysis of 27 Mesozoic and Cenozoic shales. Each individual shale was taken in amount roughly proportional to the mass of the formation which it represented.
- B. Composite analysis of 51 Paleozoic shales, weighted as in the former case.
- C. General average of A and B, giving them, respectively, weights as 3 to 5. This average represents 78 rocks.
- D. Composite analysis of 253 sandstones, about one gramme of each being taken in preparing the average sample.
- E. Composite analysis of 371 sandstones used for building purposes. Equal weights taken.
- F. Composite analysis of 345 limestones, equal weights being taken.
- G. Composite analysis of 498 limestones used for building purposes, equal weights taken.

¹ An estimate of the geological age of the earth: Sci. Trans. Royal Dublin Soc., vol. 7, p. 23, 1899.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	55.43	60.15	58.38	78.66	84.86	5.19	14.09
TiO ₂46	.76	.65	.25	.41	.06	.08
Al ₂ O ₃	13.84	16.45	15.47	4.78	5.96	.81	1.75
Fe ₂ O ₃	4.00	4.04	4.03	1.08	1.39	.54	.77
FeO.....	1.74	2.90	2.46	.30	.84	}undet.	}undet.
MnO	trace	trace	trace	trace	trace		
CaO.....	5.96	1.41	3.12	5.52	1.05	42.61	40.60
SrO	none	none	none	trace	none	none	none
BaO.....	.06	.04	.05	.05	.01	none	none
MgO	2.67	2.32	2.45	1.17	.52	7.90	4.49
K ₂ O.....	2.67	3.60	3.25	1.32	1.16	.33	.58
Na ₂ O	1.80	1.01	1.31	.45	.76	.05	.62
Li ₂ O	trace	trace	trace	trace	trace	trace	trace
H ₂ O at 110°.....	2.11	.89	1.34	.31	.27	.21	.30
H ₂ O above 110° ..	3.45	3.82	3.68	<i>a</i> 1.33	<i>a</i> 1.47	<i>a</i> .56	<i>a</i> .88
P ₂ O ₅20	.15	.17	.08	.06	.04	.42
CO ₂	4.62	1.46	2.64	5.04	1.01	41.58	35.58
S.....						.09	.07
SO ₃78	.58	.65	.07	.09	.05	.07
Cl.....				trace	trace	.02	.01
Carbon <i>b</i>69	.88	.81				
	100.48	100.46	100.46	100.41	99.86	100.09	100.34

a Includes organic matter. *b* Of organic origin.

These analyses may be used for a variety of purposes. For example, they can help in tracing the change from an average igneous rock to an average sediment. They suggest something as to the characteristic features which distinguish a good building stone from other limestones and sandstones. They are applicable to the discussion of a variety of large theoretical problems, like that chosen by Professor Joly. These considerations alone justify their publication here.

In the former edition of this bulletin (No. 148) a chapter upon analytical methods, by Dr. W. F. Hillebrand, was included. For that chapter there has been a large separate demand, and for that reason it will be expanded into a work of greater detail and issued as a distinct bulletin. It will be noticed that to Dr. Hillebrand many of the best analyses in this compilation are due, and a full statement of methods, embodying his experience, will be of the utmost value.

During the preparation of this bulletin much assistance was rendered by the petrographers and geologists connected with the Survey, especially with reference to analyses hitherto unpublished. In each case credit has been given for the data thus added. Twenty-eight analyses

of rocks from Montana, executed by or under the direction of Prof. L. V. Pirsson, of Yale University, and having been made in connection with regular Survey work, are included in the tabulations. With this exception all of the analyses given were made in the Survey laboratories. To those executed in the laboratory at Washington "record numbers" are attached, which serve to identify them on the record books of the Division of Chemistry. Of the abbreviations used for bibliographic reference only three need explanation, and they refer to the official publications of the Survey. "Ann." for Annual Report, "Mon." for Monograph, and "Bull." for Bulletin are the three in question. The others relate to well-known journals, and are familiar to all geologists. The letters P. R. C., following the description of a rock, refer to the Petrographic Reference Collection of the Survey, and are followed by the number assigned to the rock in that series.

ANALYSES.

IGNEOUS AND CRYSTALLINE ROCKS.

MAINE.

I. ROCKS FROM AROOSTOOK COUNTY.

Described by H. E. Gregory in Bull. 165. Analyses by W. F. Hillebrand, record No. 1795.

A. Quartz-trachyte (bostonite), Quoggy Joe Mountain, Presque Isle Township. Contains quartz, orthoclase, albite, and magnetite, with siderite, kaolin, and chlorite.

B. Teschenite, Mapleton Township. In dikes cutting shales. Contains andesine, augite, biotite, apatite, analcite, and magnetite.

C. Andesite, Edmunds Hill, Chapman Township. Contains labradorite, orthoclase, pyroxene, apatite, and magnetite.

D. Calciferous sandstone, New Sweden Township. Contains calcite, alkali, feldspar, quartz, magnetite, muscovite, and siderite. Included here because studied as one of the group.

	A.	B.	C.	D.
SiO ₂	72.77	46.77	61.40	54.23
Al ₂ O ₃	12.15	14.91	16.59	7.38
Fe ₂ O ₃44	7.80	2.13	.54
FeO	3.06	4.90	3.05	1.37
MgO22	2.94	2.73	3.29
CaO07	6.30	6.17	14.56
Na ₂ O	3.38	4.97	3.83	1.65
K ₂ O	4.67	2.37	1.34	1.74
H ₂ O at 105°17	.92	.82	.25
H ₂ O above 105°55	4.28	.88	1.22
TiO ₂20	2.31	.79	.28
P ₂ O ₅	trace	.98	.20	.07
ZrO ₂04	none	none	(?)
Cr ₂ O ₃	none	none	trace	(?)
V ₂ O ₃	(?)	.02	.02	(?)
FeS ₂12	.07	none	(?)
NiO	none	trace	trace	none
MnO16	.29	.13	undet.
SrO	none	.03	trace?	none
BaO03	.04	.02	none
CO ₂	2.06	trace?	none	13.48
	100.09	99.90	100.10	100.06

Traces of lithia present in all. F and Cl not looked for.

E. Diabase (basaltic glass), Mars Hill. Not resolvable into minerals.

F. Rhyolite, Haystack Mountain. Contains quartz in a groundmass of orthoclase.

G. Volcanic tuff, southeast base of Castle Hill. Contains fragments of trachyte, andesite, devitrified glass, and lapilli.

H. Diabase, Aroostook Falls, near Maine boundary line. Dike. Contains plagioclase, pyroxene, and pyrite.

	E.	F.	G.	H.
SiO ₂	42.25	75.98	31.42	49.64
Al ₂ O ₃	16.87	12.34	11.57	15.07
Fe ₂ O ₃	5.24	.85	2.37	1.66
FeO.....	10.72	.93	7.48	8.82
MgO.....	6.91	.15	5.32	5.43
CaO.....	3.33	.13	16.71	7.23
Na ₂ O.....	3.96	4.02	2.26	4.19
K ₂ O.....	.77	4.44	.74	.89
H ₂ O at 105°.....	.43	.24	.76	.45
H ₂ O above 105°.....	5.58	.64	4.17	2.81
TiO ₂	2.93	.17	2.30	2.32
P ₂ O ₅34	.03	.46	.29
ZrO ₂	none	.03	none	none
Cr ₂ O ₃03	(?)	trace	trace
V ₂ O ₃07	(?)	.06	.04
FeS ₂	trace	none	.16	.79
NiO.....	.01	none	trace	trace
MnO.....	.40	trace?	.38	.25
SrO.....	none	trace?	none	.05
BaO.....	trace?	.07	.64	.02
CO ₂	none	none	13.13	.32
	99.84	100.02	99.93	100.27

Traces of lithia present in all. Cl and F not looked for.

2. LITCHFIELDITE AND ASSOCIATED MINERALS.

A. Elæolite-syenite, var. litchfieldite, from Litchfield. Described by Bayley in Bull. 150, p. 201; and in Bull. Geol. Soc. Amer., vol. 3, p. 231. Contains elæolite, two feldspars, and lepidomelane, with sodalite, cancrinite, and zircon as accessories. Analysis by L. G. Eakins, record No. 1298. P. R. C. 77.

B. Albite from A. Sp. gr. 2.622.

C. Potash feldspar from A. Sp. gr. 2.56. Analyses B and C by W. H. Melville, record No. 1275. P. R. C. 77.

	A.	B.	C.
SiO ₂	60.39	68.28	65.14
Al ₂ O ₃	22.57	19.62	18.19
Fe ₂ O ₃42		
FeO.....	2.26	.23	.25
MgO.....	.13	.09	.16
CaO.....	.32	.31	.33
Na ₂ O.....	8.44	10.81	1.68
K ₂ O.....	4.77	.39	14.14
H ₂ O.....	.57	.09	.17
MnO.....	.08		
	99.95	99.82	100.06

The minerals found in this rock were quite fully described by Clarke in Bull. 42, pp. 28-38. Analyses by F. W. Clarke.

D. Elæolite, dark gray.

E. Hydronephelite.

F. Lepidomelane. Contains no fluorine.

	D.	E.	F.
SiO ₂	43.74	38.99	32.09
Al ₂ O ₃	34.48	33.62	18.52
Fe ₂ O ₃			19.49
FeO.....			14.10
MgO.....	trace		1.01
CaO.....	trace	.07	
Na ₂ O.....	16.62	13.07	1.55
K ₂ O.....	4.55	1.12	8.12
H ₂ O.....	.86	12.98	4.62
MnO.....			1.42
	100.25	99.85	100.92

G. Sodalite, deep blue. P. R. C. 77.

H. Cancrinite, bright orange yellow. P. R. C. 77.

I. Cancrinite, pale yellow, cleavable.

J. Cancrinite, pale yellow, granular.

	G.	H.	I.	J.
SiO ₂	37.33	36.29	35.83	37.22
Al ₂ O ₃	31.87	30.12	29.45	28.32
Fe ₂ O ₃		trace	trace	trace
MgO.....				.07
CaO.....		4.27	5.12	4.40
Na ₂ O.....	24.56	19.56	19.33	19.43
K ₂ O.....	.10	.18	.09	.18
H ₂ O.....	1.07	2.98	3.79	3.86
MnO.....		trace	trace	trace
CO ₂		6.96	6.50	6.22
Cl.....	6.83			
	101.76	100.36	100.11	99.70
O=Cl.....	1.54			
	100.22			

NEW HAMPSHIRE.

A. Elæolite-syenite, Red Hill, Moultonboro. Described by Bayley in Bull. Geol. Soc. Amer., vol. 3, p. 231. Contains elæolite, augite, hornblende, biotite, sodalite, albite, and orthoclase, with accessory apatite, sphene, magnetite, and an occasional zircon. Fibrous decomposition products are also present. P. R. C. 203.

B. Mixed albite and orthoclase from A.

C. Nepheline (elæolite) from A. Analyses by W. F. Hillebrand, record No. 1321. The mixed nepheline and feldspar were treated with dilute hydrochloric acid, and the residue was extracted with sodium carbonate solution. C represents the soluble part, and B the insoluble.

D. Camptonite, Campton Falls. Analysis by L. G. Eakins, record No. 1298. Described by J. P. Iddings in Bull. 150, p. 239. Contains hornblende, plagioclase, orthoclase, augite, iron ore, biotite, apatite, pyrite, and a mineral which appears to be analcite. Also variable calcite, serpentine, and chlorite. P. R. C. 92.

E. Quartz-porphyry, Pemigewasset. Analysis by Eakins, No. 1298.

	A.	B.	C.	D.	E.
SiO ₂	59.01	66.85	45.31	38.45	65.02
Al ₂ O ₃	18.18	19.50	32.67	19.68	17.93
Fe ₂ O ₃	1.63	} .13	-----	4.01	4.69
FeO.....	3.65		-----	11.15	.17
MgO.....	1.05	trace	.16	6.65	1.24
CaO.....	2.40	.11	2.00	9.37	1.34
Na ₂ O.....	7.03	7.44	12.60	2.77	3.04
K ₂ O.....	5.34	5.80	5.70	1.72	5.98
H ₂ O at 100°.....	.15	-----	-----	} 1.49	} .86
H ₂ O above 100°.....	.50	.31	1.56		
TiO ₂81	-----	-----	-----	-----
P ₂ O ₅	trace	-----	-----	-----	-----
MnO.....	.03	-----	-----	trace	.11
SrO.....	trace	-----	-----	-----	-----
BaO.....	.08	.07	-----	-----	-----
Li ₂ O.....	trace	none	-----	-----	-----
CO ₂12	-----	-----	4.82	-----
	99.98	100.21	100.00	100.11	100.38

VERMONT.

1. ROCKS OF MOUNT ASCUTNEY.

Analyses by W. F. Hillebrand, record Nos. 1621, 1657. Samples received from T. A. Jaggar, jr. Petrographic data furnished by R. A. Daly.

A. Typical granite. Contains quartz, orthoclase, plagioclase (microperthite), biotite, magnetite, sphene, apatite, and zircon.

B. Basic segregation in granite. Contains biotite, hornblende, quartz, plagioclase, microperthite, much sphene and apatite, some magnetite and zircon.

C. Another sample of B.

D. Syenite. Contains hornblende, augite, orthoclase, microperthite, plagioclase, biotite, quartz, magnetite, sphene, apatite, and zircon.

E. Basic segregation in syenite. Contains hornblende, augite, microperthite, orthoclase, plagioclase, quartz, magnetite, zircon, and apatite.

F. Syenite-porphry. Contains orthoclase, quartz, hornblende, magnetite, apatite, and zircon.

	A.	B.	C.	D.	E.	F.
SiO ₂	71.90	56.01	59.27	65.43	56.51	73.03
Al ₂ O ₃	14.12	15.19	15.76	16.11	16.59	13.43
Fe ₂ O ₃	1.20	2.34	2.07	1.15	1.35	.40
FeO.....	.86	4.89	3.57	2.85	6.59	1.49
MgO.....	.33	4.67	3.04	.40	2.52	.14
CaO.....	1.13	4.85	3.69	1.49	4.96	.79
Na ₂ O.....	4.52	5.66	5.63	5.00	5.15	4.91
K ₂ O.....	4.81	2.16	3.33	5.97	3.05	4.54
H ₂ O at 110°...	.18	.36	.23	.19	.21	.18
H ₂ O above 100°	.42	.90	.74	.39	.71	.35
TiO ₂35	1.13	1.12	.50	1.20	.30
P ₂ O ₅11	.53	.42	.13	.41	.06
ZrO ₂04	-----	.04	.11	.04	.06
MnO.....	.05	.40	.37	.23	.24	.15
(NiCo)O.....	none	.03	trace	(?)	trace?	(?)
BaO.....	.04	trace?	trace?	.03	.03	trace
CO ₂21	undet.	.30	trace?	.33	trace?
F.....	.06	undet.	.42	.08	.24	.08
Cl.....	.02	undet.	.03	.05	.07	.03
FeS ₂	trace	.09	.07	.07	.06	.09
	100.35	99.21	100.10	100.18	100.26	100.03
Less O.....	.03		.19	.04	.11	.04
	100.32		99.91	100.14	100.15	99.99

a Including ZrO₂.

G. Syenite-porphry. Contains orthoclase, plagioclase, micropertite, hornblende, quartz, augite, magnetite, biotite, apatite, and zircon.

H. Granite-porphry. Composition like D, but with more quartz.

I. Basic segregation in H. Contains hornblende, green and brown augite, biotite, quartz, micropertite, magnetite, little apatite, and zircon.

J. Diorite. Contains hornblende, augite, biotite, plagioclase, titaniferous magnetite, sphene, zircon, and quartz.

K. Diorite. Composition like J.

L. Another sample of K.

	G.	H.	I.	J.	K.	L.
SiO ₂	64.88	73.69	56.53	52.12	57.97	64.62
Al ₂ O ₃	16.24	12.46	16.47	16.35	<i>a</i> 17.28	16.46
Fe ₂ O ₃	1.37	1.21	1.58	3.68	2.23	1.82
FeO	2.70	1.75	5.40	6.02	3.75	2.14
MgO89	.17	2.67	4.14	2.20	1.10
CaO	1.92	.36	4.90	7.25	4.33	2.39
Na ₂ O	5.00	4.47	5.59	3.65	4.31	4.57
K ₂ O	5.61	4.92	3.80	2.34	4.12	5.21
H ₂ O at 110°19	.14	.23	.25	.18	.13
H ₂ O above 110°	.46	.24	.60	.88	.57	.39
TiO ₂69	.28	1.40	2.10	1.54	.81
P ₂ O ₅13	.04	.27	.89	.64	.21
ZrO ₂13	.14	.03	.0203
MnO14	.15	.20	.17	.15	.12
(NiCo)O	none	none	trace	trace	trace	none
BaO06	none	trace	.04	.07?	.03
CO ₂	none	trace	.05	.07	.05	.11
F08	.05	.19	.03	.04	undet.
Cl04	.02	.07	.09	undet.	.05
FeS ₂	none	none	trace	.24	.32	.19
Less O	100.53	100.09	99.98	100.33	99.75	100.38
	.04	.02	.09	.03	.02	.01
	100.49	100.07	99.89	100.30	99.73	100.37

a Including ZrO₂.

M. Basic segregation in diorite. Contains hornblende, augite, biotite, plagioclase, magnetite, apatite, zircon, and a little quartz.

N. Diabase. Contains plagioclase, augite, and magnetite.

O. Camptonite. Contains brown hornblende, plagioclase, a little augite, olivine, magnetite, and apatite.

P. Phyllite. Contains quartz, sericite, graphite?, magnetite, pyrite, rare orthoclase, plagioclase, sphene, and rutile.

Q. Micropertthite-hornfels. Contains biotite, quartz, red garnet, corundum, magnetite, iolite, micropertthite, and rarely pleonaste.

R. Cordierite-hornfels. Contains quartz, biotite, pleonaste, corundum, iolite, magnetite, plagioclase, red garnet, and epidote?.

	M.	N.	O.	P.	Q.	R.
SiO ₂	55.28	49.63	48.22	90.91	58.35	45.30
Al ₂ O ₃	17.23	14.40	14.27	4.18	21.30	30.51
Fe ₂ O ₃	1.54	2.85	2.46	.22	.03	.24
FeO.....	6.23	8.06	9.00	1.27	6.41	8.80
MgO.....	2.69	7.25	6.24	.37	2.10	3.11
CaO.....	5.60	9.28	8.45	.22	.85	.90
Na ₂ O.....	5.42	2.47	2.90	.77	1.60	1.65
K ₂ O.....	2.12	.70	1.93	.58	5.63	4.84
H ₂ O at 110°...	.20	.27	.28	.06	.31	.26
H ₂ O above 110°	.71	1.47	1.65	.74	.86	1.05
TiO ₂	1.64	1.68	2.79	.28	.87	1.48
P ₂ O ₅73	.25	.64	.05	.18	.12
ZrO ₂	trace	trace?	.03	.02	none	none
MnO.....	.24	.17	.20	trace	.13	.20
(NiCo)O.....	(?)	.04	.03	none	.03	.02
BaO.....	.06	trace?	.04	trace	.05	.03
CO ₂04	1.36	.15	.18		
C.....				.10	.40	.17
SO ₃	none	none	none	none	none	.04
Cl.....	.07	.07	.10		.03	.04
F.....	.28	trace	.05	trace	(?)	.04
FeS ₂07	.22	.36	.11	.58	1.07
	100.15	100.17	99.80	100.06	99.71	99.87
Less O.....	.13	.02	.04			
	100.02	100.15	99.76			

In these rocks the sulphur is all reckoned as pyrite, although pyrrhotite also is probably present. The carbonic acid represents either dolomite or siderite; not calcite. Traces of lithia and strontia occur in nearly all. Samples H, L, O, and Q contain traces of copper. In N there is 0.03 V₂O₅, a supplementary determination by Hillebrand.

2. MISCELLANEOUS ROCKS.

A. Amphibolite, Guilford. Described by Emerson in Mon. XXIX. Contains hornblende needles, with albite and rutile. Analysis by L. G. Eakins, record No. 1326.

B. Granitoid gneiss, north of Lincoln's.

C. The same, west slope of Little Peco.

D. Chloritic granite, East Clarendon section.

E. Hornblende-granite, East Clarendon section.

Rocks B, C, D, and E collected by C. L. Whittle, but not described. Analyses by H. N. Stokes, record No. 1396.

	A.	B.	C.	D.	E.
SiO ₂	49.16	71.02	69.97	67.33	52.60
Al ₂ O ₃	16.43	15.00	14.90	16.20	18.45
Fe ₂ O ₃	3.92	1.12	2.16	1.40	2.47
FeO	7.19	1.81	.96	2.73	6.11
MgO	8.19	.69	.37	1.31	4.22
CaO	9.21	.31	.45	2.81	7.55
Na ₂ O	3.70	2.48	2.85	3.15	3.24
K ₂ O41	5.79	6.54	2.14	1.12
H ₂ O45	1.14	.66	1.84	2.53
TiO ₂	1.03	.35	.44	.80	1.11
P ₂ O ₅16	.13	.12	.16	.20
Cr ₂ O ₃	trace	-----	-----	-----	-----
MnO23	trace	trace	trace	.23
BaO02	trace	.09	.05	-----
	100.10	99.84	99.51	99.92	99.83

MASSACHUSETTS.

1. ROCKS OF THE CONNECTICUT VALLEY.

Described by Emerson in Mon. XXIX.

A. Serpentine derived from salite, Osburn's soapstone quarry, Blandford.

B. Dark-green, oily serpentine, center of large Middlefield bed.

C. Enstatite, slightly altered, from Granville. For comparison with D.

D. Serpentine derived from enstatite, Granville.

Analyses A, B, and C by W. F. Hillebrand, record No. 1555. Analysis D by Geo. Steiger, No. 1536.

	A.	B.	C.	D.
SiO ₂	40.77	38.62	54.04	37.82
Al ₂ O ₃	1.16	.35	.52	.61
Fe ₂ O ₃	3.56	3.44	1.51	7.92
FeO.....	1.47	3.99	3.90	1.15
MgO.....	39.37	40.61	34.40	37.94
CaO.....	none	.40	none	none
Na ₂ O.....	.14	.10	.08	trace
K ₂ O.....	.10	.08		
H ₂ O at 110°.....	.49	.36	.70	.75
H ₂ O above 110°.....	12.48	10.91	3.07	12.50
TiO ₂	none	none	none	trace
P ₂ O ₅	trace	trace	none	trace
Cr ₂ O ₃28	.39	.14	.19
MnO.....	.09	.10	.11	-----
NiO.....	.17	.21	.23	.45
CoO.....	-----	-----	-----	.05
Li ₂ O.....	trace	trace	-----	-----
CO ₂	none	.52	1.32	-----
	100.08	100.08	100.02	99.38

E. Rich, dark-green serpentine, Rowe.

F. Black serpentine containing marmolite, Atwater's quarry, Russell.

G. Blackish-green serpentine containing much chromite. From
"The Crater," North Blandford.

H. Gray, splintery serpentine, Chester.

Analyses by Steiger, record No. 1536.

	E.	F.	G.	H.
SiO ₂	40.42	36.94	39.14	33.87
Al ₂ O ₃	1.86	.50	1.18	.77
Fe ₂ O ₃	2.75	6.04	4.46	2.81
FeO.....	4.27	1.94	3.14	4.25
MgO.....	35.95	38.35	41.45	38.57
CaO.....	.66	none	none	none
Na ₂ O.....	} .16	} none	} none	} none
K ₂ O.....				
H ₂ O at 100°.....	.21	.71	.34	.38
H ₂ O above 100°.....	10.51	12.07	9.48	7.00
TiO ₂	none	trace	none	none
P ₂ O ₅	trace	trace	.02	trace
Cr ₂ O ₃28	.33	.33	.38
MnO.....	trace	trace	none	.04
NiO.....	.53	.40	.47	} .33
CoO.....	trace	none	trace	
CO ₂	1.44	1.85	none	10.82
SO ₃	trace	.20	none	.20
FeS ₂43
	99.47	99.33	100.01	99.42

I. Peridotite, Belchertown. Contains hornblende, pyroxene, biotite, olivine, and magnetite. Analysis by L. G. Eakins, record No. 1326.

J. Massive, coarse, altered diabase, Leverett. Contains saussuritic feldspar with black hornblende. Analysis by Eakins, record No. 1325.

K. Tonalite, South Leverett. Dark green, chloritic. Contains reddish feldspar, dark hornblende, and a network of dark-green epidotic quartz veins. Analysis by Eakins, record No. 1326.

L. Biotite-granite, Moore's quarry, Florence. Very feldspathic. Quartz rare, with fluid inclusions. Feldspar mostly triclinic, orthoclase and microcline present in small quantities. Little muscovite, some rutile. Analysis by Eakins, record No. 1414.

	I.	J.	K.	L.
SiO ₂	48.63	51.56	55.51	73.27
Al ₂ O ₃	5.32	14.82	16.51	15.51
Fe ₂ O ₃	2.91	4.30	1.68	.33
FeO.....	3.90	7.21	4.57	1.14
MgO.....	21.79	7.36	6.73	.15
CaO.....	13.04	7.09	6.73	2.74
Na ₂ O.....	.34	4.21	3.19	4.79
K ₂ O.....	.23	.17	2.46	1.66
H ₂ O.....	2.81	1.47	1.53	.68
TiO ₂47	1.97	.91	.10
P ₂ O ₅21	.09	.17	trace
Cr ₂ O ₃36			
MnO.....	.12	trace	.11	trace
BaO.....	trace	trace	.02	
CO ₂	trace			
	100.13	100.25	100.12	100.37

M. Amphibolite, Bernardston. A black, heavy, massive hornblende rock. Analysis by L. G. Eakins, record No. 1327.

N. Porphyritic amphibolite, Heath. Analysis by Eakins, record No. 1325.

O. Amphibolite, New Salem. Analysis by Eakins, record No. 1325.

P. Amphibolite, Whitmans Ferry, Sunderland. Thin, shaly, aphanitic. Analysis by Eakins, record No. 1325.

	M.	N.	O.	P.
SiO ₂	51.72	51.38	45.48	49.86
Al ₂ O ₃	16.51	18.01	19.43	15.50
Fe ₂ O ₃	1.72	3.30	.13	2.99
FeO.....	9.56	8.53	6.58	8.01
MgO.....	6.58	5.08	11.08	7.79
CaO.....	8.89	6.27	10.66	8.89
Na ₂ O.....	2.74	5.34	2.28	3.26
K ₂ O.....	.34	.18	.11	.72
H ₂ O.....	.51	.56	3.17	1.51
TiO ₂	1.39	1.07	.77	1.58
P ₂ O ₅23	.18	.14	.11
Cr ₂ O ₃			trace	
MnO.....	trace	.19	trace	.07
BaO.....	trace	trace	.01	trace
CO ₂20	
	100.19	100.09	100.04	100.29

Q. Amphibolite, South Leverett. Deep green, ligniform. Analysis by L. G. Eakins, record No. 1327.

R. Amphibolite, Goshen. Derived from Conway limestone. Analysis by Eakins, record No. 1414.

S. Black, fissile amphibolite, Worthington. Nearly pure, matted hornblende. Titanite and sometimes zircon present. Analysis by Eakins, record No. 1326.

T. Black, fissile, porphyritic amphibolite, Warwick. Analysis by Eakins, record No. 1414. Collected by Emerson, but not described in Monograph XXIX.

	Q.	R.	S.	T.
SiO ₂	47.56	55.64	48.53	50.65
Al ₂ O ₃	16.13	16.27	16.35	13.03
Fe ₂ O ₃	1.80	1.22	2.03	.27
FeO.....	9.39	7.20	10.52	12.67
MgO.....	9.21	5.58	9.71	16.96
CaO.....	6.67	9.23	9.83	1.73
Na ₂ O.....	2.52	.91	1.36	1.37
K ₂ O.....	1.58	.19	.32	.04
H ₂ O.....	3.51	3.11	.79	2.96
TiO ₂	1.24	.50	.51	.50
P ₂ O ₅21	.23	.07	trace
Cr ₂ O ₃	trace	trace
MnO.....	.08	.28	.17	.15
BaO.....	trace	trace
	99.90	100.36	100.19	100.33

2. MISCELLANEOUS ROCKS.

Collected by B. K. Emerson and unpublished at the date of writing.

A. Wehrlite, New Braintree. Contains diallage, enstatite, augite, anorthite, biotite, apatite, chromite, magnetite, and pyrrhotite. Analysis by L. G. Eakins, record No. 1327

B. Black, serpentinized boltonite, Stow. Analysis by W. F. Hillebrand, record No. 1555.

C. Highly metamorphosed feldspathic conglomerate, graduating into arkose-gneiss, electric railroad cut, Marlboro. Analysis by George Steiger, record No. 1536.

D. Phonolite, Southboro. No description furnished. Analysis by H. N. Stokes, record No. 1653. Contains traces of chlorine and fluorine; 63.2 per cent of the rock is decomposable by hydrochloric acid.

	A.	B.	C.	D.
SiO ₂	50.64	36.92	75.35	54.22
Al ₂ O ₃	7.93	.10	13.03	20.20
Fe ₂ O ₃	1.41	1.19	.62	2.35
FeO.....	14.82	.87	.94	1.02
MgO.....	18.58	43.99	.21	.29
CaO.....	3.41	.59	1.33	.70
Na ₂ O.....	.96	} .05	2.44	9.44
K ₂ O.....	.21		5.14	4.85
H ₂ O at 100°.....	} .87	.72	.15	.42
H ₂ O above 100°.....		14.70	.73	5.57
TiO ₂82	none	.21	.38
P ₂ O ₅27	trace	.08	.11
Cr ₂ O ₃05	none	none
MnO.....	.16	trace	none	.19
BaO.....	none	.07	trace
CO ₃90	.03	trace
SO ₃03	none
	100.13	100.03	100.36	99.74

Keratophyr, Marblehead Neck. Described by Sears, in Bull. Museum Compar. Zoölogy, vol. 16, No. 9, p. 170. Contains crystals of feldspar, with a decomposed base, irregular patches of quartz, some scales of biotite and grains of magnetite, and also some limonite and earthy matter. The feldspar is anorthoclase. Analyses by T. M. Chatard, record No. 1176.

A. The rock.

B. The separated feldspar.

	A.	B.
SiO ₂	70.23	65.66
Al ₂ O ₃	15.00	20.05
Fe ₂ O ₃	1.99	trace
FeO.....	undet.	trace
MgO38	.18
CaO.....	.33	.67
Na ₂ O	4.98	6.56
K ₂ O.....	4.99	6.98
H ₂ O at 110°91	.04
H ₂ O at redness.....	1.28	.37
TiO ₂ ?.....	.03	undet.
P ₂ O ₅06	undet.
MnO24	.13
	100.42	100.64

Feldspars from schists of Berkshire County. Described by Wolff, Mon. XXIII, pp. 60 and 187. All albite. Analyses by R. B. Riggs, record Nos. 507 and 567.

A. From feldspathic schist, central shaft of the Hoosac tunnel. P. R. C. 129.

B, C. From the porphyritic mica-schist of Greylock Mountain.

	A.	B.	C.
SiO ₂	69.69	68.08	67.83
Al ₂ O ₃ , Fe ₂ O ₃	18.60	20.11	19.92
MgO20	(?)	(?)
CaO	trace	trace	trace
Na ₂ O	10.28	11.00	11.65
K ₂ O.....	.40	.36	.25
Ignition42	.31	.12
MnO		trace	trace
	99.59	99.86	99.77

Fe₂O₃ less than 0.5 in any case.

CONNECTICUT.

A. Olivine-basalt, main flow, Pine Hill, South Britain. Contains plagioclase, pink augite, olivine, and magnetite. The rock is quite fresh.

B. Olivine-basalt, anterior flow, South Britain. Contains plagioclase, uraltized augite, olivine, and magnetite. Rock much altered and containing numerous amygdules of calcite and prehnite, stained by iron.

Rocks A and B studied by W. H. Hobbs, who furnishes the petrographic data. Hitherto unpublished. Analyses by W. F. Hillebrand, record No. 1842.

C. Basic pitchstone (tachylyte), from the so-called "ash-bed" north-east of Meriden. Described by Emerson in Bull. Geol. Soc. Amer., vol. 8, p. 77. Analysis by H. N. Stokes, record No. 1641.

	A.	B.	C.
SiO ₂	52.40	47.52	46.86
Al ₂ O ₃	13.55	13.91	13.96
Fe ₂ O ₃	2.73	7.06	5.23
FeO.....	9.79	3.76	4.67
MgO.....	5.53	6.84	7.69
CaO.....	10.01	5.71	9.42
Na ₂ O.....	2.32	3.06	1.85
K ₂ O.....	.40	.77	2.02
H ₂ O at 105°.....	.62	1.75	1.29
H ₂ O above 105°.....	1.05	4.55	3.43
TiO ₂	1.08	1.19	1.13
P ₂ O ₅12	.15	.15
NiO.....	trace	trace
MnO.....	.26	.18	trace
SrO.....	none	none	trace
BaO.....	trace?	trace	.03
Li ₂ O.....	none	trace	trace
CO ₂	3.68	2.19
F.....	trace
FeS ₂ or Fe ₇ S ₈ <i>a</i>13
	99.99	100.13	99.92

a Calculated as pyrite.

NEW YORK.

I. ROCKS OF THE ADIRONDACK REGION.

Collected by J. F. Kemp, who furnishes the petrographic data. A and G are described by Kemp in 19th Ann., Part III, p. 383. The other descriptions are hitherto unpublished. Analyses A and B by George Steiger, record No. 1715; C to G by W. F. Hillebrand, record No. 1717.

A. Wall rock of iron mine near Lincoln Pond, Elizabethtown. Varies from norite to gabbro. Chief minerals, green augite, hypersthene, brown hornblende, plagioclase, and magnetite. Microperthitic feldspar less common. Garnet varies from absence to abundance.

B. Coarse gabbro, top of Whiteface Mountain. A pyroxenic phase of the anorthosite. Contains much labradorite, abundant light-green augite, brown hornblende, shreds of more or less bleached brown biotite, and magnetite.

	A.	B.
SiO ₂	44.77	53.18
Al ₂ O ₃	12.46	23.25
Fe ₂ O ₃	4.63	1.53
FeO.....	12.99	1.82
MgO.....	5.34	2.60
CaO.....	10.20	11.18
Na ₂ O.....	2.47	3.97
K ₂ O.....	.95	.86
H ₂ O at 100°.....	.12	.15
H ₂ O above 100°.....	.48	.98
TiO ₂	5.26	.45
P ₂ O ₅28	.09
NiO, CoO.....	trace?	(?)
MnO.....	.17	.11
BaO.....	trace?	trace?
CO ₂37	.34
S.....	a. 26	trace
	100.75	100.51

a Mainly present as pyrrhotite.

C. Pyroxenic anorthosite, Elizabethtown. Contains largely labradorite, subordinate light-green augite, less brown hornblende, and a little magnetite.

D. Norite, with close affinity to gabbro. Intrusion in C. Contains labradorite, hypersthene, garnets, green augite, brown hornblende, a little brown biotite, magnetite, and apatite.

E. Diabasic norite or gabbro, Elizabethtown. Contains the same minerals as D.

F. Gneissoid derivative, by pressure, of E. The same minerals, but with hornblende more abundant.

G. Gabbro, wall rock of Split Rock mine. Contains augite, hypersthene, brown hornblende, garnet, plagioclase, magnetite, and possibly spinel.

	C.	D.	E.	F.	G.
SiO ₂	56.94	47.16	44.97	46.74	47.88
Al ₂ O ₃	20.82	14.45	15.38	16.63	18.90
Fe ₂ O ₃83	1.61	2.29	2.17	1.39
FeO.....	3.02	13.81	12.39	10.60	10.45
MgO.....	2.36	5.24	10.89	6.11	7.10
CaO.....	9.41	8.13	7.50	8.66	8.36
Na ₂ O.....	3.36	3.09	3.02	3.81	2.75
K ₂ O.....	1.58	1.20	.56	.86	.81
H ₂ O at 110°.....	.21	.12	.10	.12	.18
H ₂ O above 110°.....	.59	.48	.65	.73	.43
TiO ₂44	3.37	1.18	2.54	1.20
P ₂ O ₅07	.57	.14	.33	.20
V ₂ O ₃	(?)	(?)	.02	(?)	(?)
NiO, CoO.....	trace	.02	.02	.03	.02
MnO.....	.11	.24	.22	.26	.16
BaO.....	.05	trace	trace	trace	trace
CO ₂45	.35	.23	.07	.12
S.....	trace	.14	.06	.11	.07
	100.24	99.98	99.64	99.77	100.02

ZrO₂, Cl, and F not looked for. Cr₂O₃, Li₂O, and SrO present in traces. S is in part, at least, present as pyrrhotite.

2. PERIDOTITE FROM DEWITT, NEAR SYRACUSE.

A rock described by Darton and Kemp, Amer. Journ. Sci., 3d series, vol. 49, p. 456. Contains olivine, partly serpentized, biotite, and augite, with magnetite, apatite, and perovskite. Possibly a little chromite also. Classed by Kemp as limburgite.

Analysis by H. N. Stokes, record No. 1491. The FeO represents the total iron, because the sulphides present, possibly as pyrrhotite, prevent the separate estimation of the two iron oxides.

SiO ₂	36.80	NiO.....	.09
Al ₂ O ₃	4.16	MnO.....	.13
FeO.....	8.33	BaO.....	.12
MgO.....	25.98	SrO.....	trace
CaO.....	8.63	CO ₂	2.95
Na ₂ O.....	.17	SO ₃06
K ₂ O.....	2.48	S.....	.95
H ₂ O below 100°.....	.51		100.22
H ₂ O above 100°.....	6.93	O=S.....	.47
TiO ₂	1.26		99.75
P ₂ O ₅47		
Cr ₂ O ₃20		

NEW JERSEY.

Rocks A, B, and C, described by J. P. Iddings in Bull. *150, pp. 254, 209, 236. Analyses by Eakins, record No. 1299.

A. Basalt, Watchung Mountain, Orange. Contains pyroxene, mostly malacolite, plagioclase, magnetite, and glass, with variable amounts of serpentine or chlorite. The feldspar is partly altered into a mineral which is probably prehnite. P. R. C. 102.

B. Elæolite-syenite, Beemersville. Contains nephelite, orthoclase, ægirite, and biotite, with melanite, sphene, apatite, zircon, and magnetite in smaller amounts. Sodalite is probably present also. P. R. C. 78.

C. Minette, Franklin Furnace. Contains alkali-feldspar, biotite, monoclinic pyroxene, magnetite, epidote, calcite, chlorite, apatite, sphene, pyrite, and a little secondary quartz. P. R. C. 91.

D. Nepheline-syenite, Brookville. Collected by N. H. Darton and described by F. L. Ransome in Amer. Journ. Sci., 4th series, vol. 8, p. 417. Contains alkali-feldspars, nepheline largely altered to secondary minerals, brown amphibole, biotite, cancrinite, plagioclase, muscovite, ægirine-augite, apatite, titanite, fluorite, and a few specks of magnetite. Also secondary analcite, sericite, and natrolite (?). Analysis by Steiger, record No. 1807.

	A.	B.	C.	D.
SiO ₂	51.36	53.56	40.71	54.68
Al ₂ O ₃	16.25	24.43	19.46	21.63
Fe ₂ O ₃	2.14	2.19	7.46	2.22
FeO	8.24	1.22	6.83	2.00
MgO	7.97	.31	6.21	1.25
CaO	10.27	1.24	11.83	2.86
Na ₂ O	1.54	6.48	1.80	7.03
K ₂ O	1.06	9.50	3.26	4.58
H ₂ O at 100°	1.33	.93	1.53	.27
H ₂ O above 100°				1.88
TiO ₂79
P ₂ O ₅28
MnO09	.10	.18	trace
NiO03			
BaO05
F22
SO ₃07
CO ₂74	none
	100.28	99.96	100.01	99.81
Less O=F09
				99.72

PENNSYLVANIA.

A. Aporhyolite, Monterey, Franklin County. Described by Florence Bascom in Bull. 150, p. 343. Contains alkali feldspars, quartz, sericite, epidote, titaniferous magnetite, leucoxene, and rarely sphene. Analysis by H. N. Stokes, record No. 1479. P. R. C. 136.

B. Quartz-porphyry, same locality. Analyzed for G. H. Williams, but never described. Analysis by L. G. Eakins, record No. 1350

	A.	B.
SiO ₂	76.34	73.85
Al ₂ O ₃	11.60	13.15
Fe ₂ O ₃	2.41	3.27
FeO.....	.30	.36
MgO.....	.06	.32
CaO.....	.55	.82
Na ₂ O.....	5.50	2.29
K ₂ O.....	2.75	5.42
H ₂ O at 100°.....	.10	} .71
H ₂ O above 100°.....	.39	
TiO ₂26
P ₂ O ₅	trace	.06
MnO.....	trace	.09
BaO.....	.09
CO ₂	trace
	100.35	100.34

DELAWARE.

Feldspars from gabbros. Separated by Diller, described by Chester in Bull. 59. Analyses by R. B. Riggs, record Nos. 459, 460, 461.

A. Sp. gr. 2.592 to 2.877, from a typical gabbro, Brandywine Creek, Wilmington.

B. Sp. gr. 2.592 to 2.780, from a typical hypersthene-gabbro, Fifth and Van Buren streets, Wilmington.

C. Sp. gr. 2.592 to 2.749, from "Iron Hill hyperite changing to diorite," near Whitaker's ore pit.

	A.	B.	C.
SiO ₂	70.37	51.44	44.09
Al ₂ O ₃	18.36	30.05	35.41
Fe ₂ O ₃58	.96	.51
MgO.....	.04	trace	none
CaO.....	5.08	13.19	18.47
Na ₂ O.....	4.32	4.07	.99
K ₂ O.....	.63	.21	.19
MnO.....	trace	trace
Ign.....	.45	.35	.35
	99.83	100.27	100.01

Samples dried at 105°. FeO not determined separately.

MARYLAND, INCLUDING THE DISTRICT OF COLUMBIA.

1. PERIDOTITE AND PYROXENITE.

Rocks A to I, inclusive, described by G. H. Williams in Amer. Geologist, vol. 6, p. 35.

A. Porphyritic lherzolite, Johnny Cake Road, Baltimore County. Contains olivine, bronzite, and diallage, the olivine partly serpentinized. Analysis by T. M. Chatard, record No. 1094.

B. Pyroxenite, Johnny Cake Road. Consists entirely of hypersthene and diallage. P. R. C. 110.

C, D. Alterations of B. B, C, and D dried at 104°.

E. Smaragdite rock, altered pyroxenite, Dogwood Road, Baltimore County.

Analyses B, C, D, and E by J. E. Whitfield, record Nos. 975, 976. C, D, and E are from the laboratory record, and do not appear in the published paper.

	A.	B.	C.	D.	E.
SiO ₂	43.87	50.80	50.10	51.94	53.22
Al ₂ O ₃	1.64	3.40	2.00	2.53	3.14
Fe ₂ O ₃	8.94	1.39	2.38	2.88	-----
FeO.....	2.60	8.11	8.68	9.38	7.95
MgO.....	27.32	22.77	26.85	25.97	20.09
CaO.....	6.29	12.31	5.06	3.60	14.44
Na ₂ O.....	.50	} trace	} none	} none	} trace
K ₂ O.....					
H ₂ O at 110°.....	1.08	} .52	} 4.16	} 2.82	} .98
H ₂ O at red heat.....	7.64				
TiO ₂12	none	none	none	none
P ₂ O ₅	trace	trace	none	none	-----
Cr ₂ O ₃44	.32	.36	.60	.23
NiO.....	trace	-----	-----	-----	-----
MnO.....	.19	.17	.29	trace	.11
SO ₃	-----	trace	trace	.19	trace
Cl.....	-----	.24	.26	.16	.26
	100.63	100.03	100.14	100.07	100.42

F, G. Two samples of websterite, Hebbville, 6 miles west of Baltimore.

H. Bronzite from G.

I. Diopside from G.

The rock consists entirely of bronzite and diopside. Analyses by T. M. Chatard, record Nos. 1094, 1123.

J. Websterite, Oakwood, Cecil County. Composed of hypersthene and diallage. Analysis by W. F. Hillebrand, record No. 1755. Described by A. G. Leonard, but hitherto unpublished.

K. Cortlandtite, Ilchester, Howard County. Analysis by W. F. Hillebrand, record No. 1422. Published by Williams in Fifteenth Ann., page 674. The rock consists of olivine, pyroxene, and large hornblende crystals, the latter considerably altered to talc.

	F.	G.	H.	I.	J.	K.
SiO ₂	53.98	52.55	54.53	51.80	53.21	39.20
Al ₂ O ₃	1.32	2.71	1.93	2.21	1.94	4.60
Fe ₂ O ₃	1.41	1.27	1.70	1.29	1.44	3.45
FeO	3.90	4.90	8.92	3.50	7.92	6.15
MgO	22.59	20.39	29.51	17.76	20.78	31.65
CaO	15.47	16.52	2.25	20.99	13.12	3.23
Na ₂ O	undet.	} .27	-----	undet.	.11	.42
K ₂ O	undet.			undet.	.07	.14
H ₂ O at 100°09	} 1.09	} 1.14	} .65	.14	.50
H ₂ O above 100°	.83				.87	9.38
TiO ₂15	.14	undet.	.13	.26	.52
P ₂ O ₅	trace	trace	trace	trace	trace	trace
ZrO ₂	-----	-----	-----	-----	trace	-----
V ₂ O ₃	-----	-----	-----	-----	.03	-----
Cr ₂ O ₃53	.44	.30	.51	.20	.41
NiO	trace	-----	-----	-----	} .03	.30
CoO	-----	-----	-----	-----		-----
MnO21	.24	.28	trace	.22	.20
SrO	-----	-----	-----	-----	none	-----
BaO	-----	-----	-----	-----	none	-----
Li ₂ O	-----	-----	-----	-----	trace	trace
FeS ₂	-----	-----	-----	-----	.03	-----
CO ₂	-----	-----	-----	-----	.10	-----
	100.48	100.52	100.56	98.84	100.47	100.15

2. GABBRO AND DIORITE.

Rocks A to E, inclusive, described by G. H. Williams, but not in detail, in 15th Ann., pp. 673, 674. Some of the rocks, with other analyses, are discussed by him in Bull. 28. Analyses by W. F. Hillebrand, record No. 1422.

A. Olivine-gabbro, Orange Grove, Baltimore County. Contains plagioclase, diallage, hypersthene, fresh olivine, magnetite, and apatite, and sometimes hornblende.

B. Hypersthene-gabbro, Wetheredville, Baltimore County. Rich in hypersthene and diallage, with plagioclase, magnetite, and apatite.

C. Gabbro-diorite, Ilchester, Howard County. A coarse anorthite-hornblende rock, probably an altered gabbro.

D. Biotite-diorite, Triadelphia, Montgomery County.

E. Biotite-diorite, Georgetown, D. C.

To these may be added an apparently unpublished analysis made for Williams by L. G. Eakins, record No. 1350, as follows:

F. Hornblende-diorite, Rock Creek tunnel shaft, Washington, D. C. Relations to E not stated.

	A.	B.	C.	D.	E.	F.
SiO ₂	48.91	44.76	43.42	55.97	56.41	56.18
Al ₂ O ₃	8.81	18.82	22.37	15.60	15.19	14.76
Fe ₂ O ₃	1.04	2.19	.81	1.21	1.60	2.12
FeO	9.52	4.73	9.25	6.28	6.24	6.98
MgO	15.19	11.32	5.75	6.83	7.18	8.11
CaO	14.69	14.58	13.34	7.31	6.77	7.97
Na ₂ O64	.89	1.24	2.23	2.21	1.62
K ₂ O10	.11	1.13	1.25	1.34	.80
H ₂ O at 100°07	.17	.09	.18	.08	} 1.37
H ₂ O above 100°	.52	2.36	1.54	1.85	2.00	
TiO ₂37	.13	1.25	1.11	.69
P ₂ O ₅	trace	none	.10	.16	.05	.08
SnO ₂ , ZrO ₂ ?14
Cr ₂ O ₃15	.08	trace	.04	.05
MnO16	.15	.06	.08	.11	.17
Li ₂ O	trace	trace	trace	trace	trace
	100.17	100.29	100.35	100.10	100.06	100.16

The following rocks, from Cecil County, have been studied by A. G. Leonard. Descriptions as yet unpublished.

G. Quartz-biotite-hornblende-diorite, near the foundry on Stone Run. Granitic in appearance.

H. Quartz-biotite-hornblende-diorite, near Porter's bridge on Octoraro Creek.

I. A true diorite with very little quartz, Stone Run, one-half mile northwest of the mill near Rising Sun.

J. Norite, three-fourths of a mile northwest of McKinsey's mill.

Analyses by Hillebrand, record No. 1755.

	G.	H.	I.	J.
SiO ₂	58.57	55.16	44.04	48.02
Al ₂ O ₃	16.10	17.51	20.01	20.01
Fe ₂ O ₃	2.89	2.62	4.22	1.13
FeO.....	6.12	5.83	8.61	7.29
MgO.....	2.33	4.35	5.01	10.05
CaO.....	7.39	8.50	11.68	11.42
Na ₂ O.....	2.11	1.83	1.24	.51
K ₂ O.....	1.01	1.08	.15	.05
H ₂ O at 105°.....	.21	.18	.11	.10
H ₂ O above 105°.....	1.27	2.01	1.90	.57
TiO ₂	1.41	.64	2.24	.23
P ₂ O ₅37	.21	.52	trace
ZrO ₂09	.02	.10	none
V ₂ O ₃02	.04	.05	.02
Cr ₂ O ₃	none	trace	none	.03
NiO, CoO.....	none	.01	.01	.01
MnO.....	.18	.15	.28	.18
SrO.....	trace	trace	none	none
BaO.....	trace	trace	none	none
Li ₂ O.....	trace	trace	trace	trace
FeS ₂	trace	.03	.25	.11
CO ₂	none	none	none	.25
	100.07	100.17	100.42	99.98

3. GRANITE AND GNEISS.

For descriptions see Williams, 15th Ann., p. 657, and Keyes, *ibid.*, p. 685. The Rowlandsville granite is described by Grimsley in Journ. Cincinnati Soc. Nat. Hist., vol. 17, p. 78.

A. White granite, Brookville, Montgomery County. (Williams.)

B. Binary granite, Guilford, Howard County. Contains quartz, orthoclase, a little plagioclase, muscovite, and biotite, with occasional zircon and apatite. (Keyes.)

C. Biotite-granite, Woodstock. Light colored, fine grained. Quartz, feldspar, and biotite, with accessory allanite and epidote. (Keyes.)

D. Biotite-granite, Rowlandsville, Cecil County. Dark colored. Contains plagioclase, orthoclase, quartz, epidote, biotite, sphene, magnetite, and apatite, with a little secondary muscovite. The percentages of the several minerals are computed by Grimsley from the analysis.

Analyses by W. F. Hillebrand, record Nos. 1220, 1422, 1455. In B and C manganese was present, barium and strontium were not tested for, and the alumina contains possible titanium and phosphoric acid.

	A.	B.	C.	D.
SiO ₂	74.87	72.57	71.79	66.68
Al ₂ O ₃	14.27	15.11	15.00	14.93
Fe ₂ O ₃	trace	.59	.77	1.58
FeO.....	.51	1.02	1.12	3.32
MgO.....	.16	.30	.51	2.19
CaO.....	.48	1.65	2.50	4.89
Na ₂ O.....	3.06	3.92	3.09	2.65
K ₂ O.....	5.36	4.33	4.75	2.05
H ₂ O at 100°.....	.26	} .47	} .64	.16
H ₂ O above 100°.....	.66			1.09
TiO ₂05	undet.	undet.	.50
P ₂ O ₅21	undet.	undet.	.10
MnO.....	trace	undet.	undet.	.10
SrO.....		undet.	undet.	trace
BaO.....		undet.	undet.	.08
Li ₂ O.....	trace	trace	trace	trace
	99.89	99.96	100.17	100.32

E. Biotite-granite, Dorseys Run, Howard County. Typical, dark colored. Quartz, feldspar, and biotite, with accessory allanite and epidote. (Keyes.)

F. Same locality, light-colored dikes.

G. Inclusions in E. Derived from gneiss.

H. The typical gneiss of the Dorseys Run area. Perhaps of sedimentary origin.

Descriptions by Keyes. Analyses by Hillebrand. The remarks appertaining to B and C apply here also. Record No. 1220.

	E.	F.	G.	H.
SiO ₂	62.91	70.45	57.33	48.92
Al ₂ O ₃	19.13	15.98	15.31	16.57
Fe ₂ O ₃98	.75	3.39	4.21
FeO.....	3.20	1.84	8.19	9.18
MgO.....	1.69	.77	4.36	5.98
CaO.....	4.28	2.60	3.95	9.69
Na ₂ O.....	3.94	3.83	1.22	2.47
K ₂ O.....	3.38	3.59	4.57	1.56
H ₂ O.....	.63	.45	1.80	1.68
Li ₂ O.....	trace	trace	trace	trace
	100.14	100.26	100.12	100.26

I. Biotite-granite, Sykesville. Contains quartz, feldspar, and biotite, with accessory magnetite, zircon, and apatite.

J. Inclusion in I, derived from limestone. Yellowish central portion. Consists of epidote and quartz, with a little chlorite.

K. Same inclusion, fine-grained dark outer zone. Quartz, garnet, and epidote, with a little magnetite. Feldspar and muscovite in some portions. Descriptions by Keyes. Analyses by Hillebrand, record No. 1220, with the same limitations as in the Guilford, Woodstock, and Dorseys Run granites.

	I.	J.	K.
SiO ₂	71.45	67.02	47.35
Al ₂ O ₃	14.36	13.77	29.76
Fe ₂ O ₃	2.07	4.64	2.94
FeO.....	2.78	1.02	3.15
MgO.....	1.17	.65	1.60
CaO.....	1.58	11.09	2.20
Na ₂ O.....	1.95	.66	2.84
K ₂ O.....	3.28	.09	6.83
H ₂ O.....	1.30	1.16	3.15
Li ₂ O.....	trace	trace	trace
	99.94	100.10	99.82

L. Typical gneiss of Washington, D. C., from quarry of Potomac Stone Company, 1 mile below Chain Bridge. A basic granite.

M. Fine-grained, fissile, chloritic gneiss, from northwest of Cabin John Bridge. Also a basic granite.

N. Fine-grained, hard gneiss, from the second lock at the Great Falls of the Potomac. Probably of sedimentary origin.

Described by Williams. Analyses by Hillebrand, record No. 1459.

	L.	M.	N.
SiO ₂	67.22	63.43	78.28
Al ₂ O ₃	15.34	16.69	9.96
Fe ₂ O ₃	2.78	3.36	1.85
FeO.....	3.41	3.87	1.78
MgO.....	1.65	2.33	.95
CaO.....	1.36	.80	1.68
Na ₂ O.....	2.00	2.38	2.73
K ₂ O.....	3.26	3.22	1.35
H ₂ O at 110°.....	.29	.23	.12
H ₂ O above 110°.....	1.68	2.67	.83
TiO ₂84	.91	.70
P ₂ O ₅14	.11	.11
MnO.....	.13	.09	.08
SrO.....	trace	trace	trace
BaO.....	.04	.03	.02
Li ₂ O.....	trace	trace	trace
	100.14	100.12	100.44

To these may be added three analyses of feldspars from the pegmatite of Jones Falls, Baltimore. Described by S. L. Powell in Johns Hopkins Univ. Circular, vol. 12, p. 49. Analyses by W. F. Hillebrand, record No. 1421.

O. Flesh-colored microcline.

P. Green microcline.

Q. Albite-oligoclase, near Ab_4An .

	O.	P.	Q.
SiO ₂	65.06	68.48	63.72
Al ₂ O ₃	18.41	16.11	22.26
Fe ₂ O ₃	} trace	.20	} trace
FeO17	
MgO04	.03	.06
CaO26	.23	3.58
Na ₂ O	1.60	1.27	8.98
K ₂ O	14.30	12.99	.76
H ₂ O at 100°04	.06	.09
H ₂ O above 100°26	.26	.43
SrO	trace	trace	trace
BaO13	.05	-----
	100.10	99.85	99.88

P contained a little unseparated quartz. All three contained traces of lithia. In O and Q the traces of iron were weighed with the alumina.

Bull. 168—4

4. MISCELLANEOUS ROCKS.

A. Typical diabase, Rocky Ridge. Analyzed for J. S. Diller, but not described. Analysis by E. A. Schneider, record No. 1370.

B. Ottrelite-phyllite rock, Liberty, Frederick County. Analyzed for S. H. Williams, but not described.

C. Ottrelite separated from B. Analyses B and C by L. G. Eakins, record No. 1349.

D. Quartz-schist, Shoemaker quarry, near Stevenson Station, Green Spring Valley. Described by Bayley in Bull. 150, p. 302. Contains quartz, muscovite, occasional tourmalines, microcline, zircon, and iron stains. Analysis by Schneider, No. 1370. P. R. C. 119.

E. Mica separated from D. Analysis by Schneider, No. 1377.

F. Sericite-schist, Ladiesburg. Described by Bayley in Bull. 150, p. 317. Contains quartz, sericite, chlorite, a mineral thought to be kaolin, zircon, iron oxide, and rutile. Analysis by George Steiger, record No. 1600. P. R. C. 126.

G. Another sample of F. Analysis by Schneider, No. 1370.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	51.68	34.92	23.40	91.65	44.93	57.24	58.11
Al ₂ O ₃	15.87	32.31	39.31	1.59	29.81	23.48	21.84
Fe ₂ O ₃	1.46	10.21	5.14	3.57	6.10	3.19	2.62
FeO.....	8.43	8.46	21.94	.21		4.87	5.63
MgO.....	7.84	1.13	2.18	.17	1.16	.93	1.85
CaO.....	11.08	.36	trace	none	-----	.09	none
Na ₂ O.....	1.86	2.12	.20	.07	.50	1.18	.97
K ₂ O.....	.34	1.87	.20	1.93	10.28	3.55	3.66
H ₂ O at 100°...	.16	} 5.29	} 6.81	-----	1.38	.33	.35
H ₂ O above 100°	.15			.60	4.88	4.65	4.05
TiO ₂72	3.37	1.19	.13	1.05	.08	.81
P ₂ O ₅12	.23	trace	none	-----	.09	.21
MnO.....	.15	trace	trace	trace	trace	none	.19
Li ₂ O.....	-----	-----	-----	-----	trace	-----	-----
F.....	-----	-----	-----	-----	.22	-----	-----
	99.86	100.27	100.37	99.92	100.31	99.68	100.29

VIRGINIA.

A. Andesite, $3\frac{1}{2}$ miles east of Front Royal. Described by Keith in 14th Ann., p. 305. Intermediate between diabase and quartz-porphry. Contains plagioclase, quartz, magnetite, ilmenite, and a little epidote. Analysis by George Steiger, record No. 1450.

B. Porphyritic felsite or felsophyre, forks of Straight Creek, 3 miles ENE. of Monterey. Described by Darton and Keith in Am. Journ. Sci., 4th series, vol. 6, p. 305. Contains phenocrysts of biotite, orthoclase, and plagioclase, the mica and feldspar being about equal in amount. In the groundmass are feldspar, quartz, magnetite, or ilmenite, and a little secondary chlorite with less muscovite. No glass. Analysis by Hillebrand, record No. 1665.

	A.	B.
SiO ₂	51.08	69.56
Al ₂ O ₃	11.37	15.52
Fe ₂ O ₃	11.17	1.67
FeO.....	5.64	1.19
MgO.....	3.96	.41
CaO.....	5.20	1.20
Na ₂ O.....	5.54	4.46
K ₂ O.....	1.50	4.68
H ₂ O at 110°.....	1.31	.34
H ₂ O above 110°.....	.19	.67
TiO ₂	2.67	.31
P ₂ O ₅39	.08
MnO.....	.22	.07
SrO.....		trace
BaO.....		.10
Li ₂ O.....		trace
CO ₂		none
Cl, F.....		(?)
S.....		trace
	100.24	100.26

NORTH CAROLINA.

Rocks A to F collected by Arthur Keith, who furnishes the petrographic data. Analyses B, C, D, and E by W. F. Hillebrand, record No. 1707. Analyses A and F by H. N. Stokes, record No. 1710.

A. Porphyritic rhyolite, 2 miles west of Barmers Elk, Watauga County. Contains orthoclase and plagioclase, with less quartz, epidote chlorite, and pyrite.

B. Quartz-porphyry, $2\frac{1}{2}$ miles northwest of Blowing Rock, Watauga County. Contains quartz and orthoclase, with subordinate sericite chlorite, and biotite.

C. Diorite, east end of Hump Mountain, Mitchell County. Contains plagioclase, orthoclase, and hornblende, with less quartz, biotite magnetite, and garnet.

D. Garnetiferous diabase, $1\frac{3}{4}$ miles southeast of Cranberry. Contains plagioclase and hornblende, with less garnet, biotite, and magnetite.

	A.	B.	C.	D.
SiO ₂	62.35	79.75	46.91	52.11
Al ₂ O ₃	13.24	10.47	15.85	13.70
Fe ₂ O ₃	3.52	.64	2.86	1.22
FeO.....	6.33	.92	9.95	9.86
MgO.....	.85	.13	7.01	8.08
CaO.....	3.34	.15	9.62	12.16
Na ₂ O.....	2.79	1.36	2.65	1.31
K ₂ O.....	3.95	6.01	.69	.16
H ₂ O at 110°.....	.11	.08	.24	.06
H ₂ O above 110°.....	1.21	.60	1.62	.53
TiO ₂	1.18	.15	2.03	.32
P ₂ O ₅57	trace	.26	.05
ZrO ₂05	none	none
Cr ₂ O ₃	none		.01	
V ₂ O ₃03	
CoO, NiO.....	none	none	.03	.03
MnO.....	.08	trace	.22	.20
SrO.....	trace	trace	trace?	none
BaO.....	.16	.06	trace?	none
Li ₂ O.....	trace	trace	trace	trace
S.....		none	none	trace
	99.68	100.37	99.98	99.79

E. Epidote-chlorite-schist, $\frac{1}{4}$ mile northeast of Montezuma, Mitchell County. Contains epidote and feldspar, with less chlorite, hornblende, and magnetite.

F. Metamorphosed amygdaloid, 3 miles southeast of Boone, Watauga County. Contains orthoclase and plagioclase, with less sericite, chlorite, and magnetite.

G. Pyroxenite, var. websterite, from Webster. Described by Williams, Amer. Geologist, vol. 6, p. 35. Consists of diopside and bronzite. Analysis by E. A. Schneider, record No. 1096. Material dried at 105°. P. R. C. 388.

H. Spherulitic rhyolite, Sam Christian gold mine, Montgomery County. Described by Diller, Amer. Journ. Sci., 4th ser., vol. 7, p. 337. The rock of the supposed fossil, palæotrochis. Contains feldspar and quartz, with a little biotite and sericite. Analysis by Hillebrand, record No. 1796.

	E.	F.	G.	H.
SiO ₂	47.85	43.62	55.14	79.57
Al ₂ O ₃	16.51	17.30	.66	11.41
Fe ₂ O ₃	4.16	14.13	3.48	.20
FeO.....	7.43	6.83	4.73	.70
MgO.....	6.24	2.34	26.66	a little
CaO.....	7.00	1.63	8.39	.21
Na ₂ O.....	3.20	3.45	.30	3.46
K ₂ O.....	.82	3.03	-----	3.52
H ₂ O at 110°.....	.21	.30	} .38	.18
H ₂ O above 110°.....	4.00	2.93		.61
TiO ₂	2.28	2.75	trace	.11
P ₂ O ₅35	1.34	.23	trace
ZrO ₂03	-----	-----	-----
Cr ₂ O ₃01	none	.25	-----
V ₂ O ₃05	-----	-----	-----
CoO, NiO.....	.03	none	.11	-----
MnO.....	.24	trace	.03	none
SrO.....	trace?	trace	-----	-----
BaO.....	trace?	.09	-----	.05
Li ₂ O.....	trace	trace	-----	-----
	100.41	99.74	100.36	100.02

The following rocks from Corundum Hill were collected, analyzed, and described by T. M. Chatard, Bull. 42, p. 45.

I. Altered gneiss. Made up of micaceous scales, with grains of quartz and some earthy matter. Record No. 238.

J. Dunite. Olivine rock containing a little chromite. Record No. 256.

K, L. Yellow, clay-like alterations of dunite. Record Nos. 254, 255.

Other analyses of altered rocks are given in the paper, and also analyses of associated minerals.

	I.	J.	K.	L.
SiO ₂	64.27	40.11	40.18	40.04
Al ₂ O ₃	16.75	.88	1.35	3.17
Fe ₂ O ₃	6.08	1.20	10.97	12.15
FeO89	6.09
MgO	1.74	48.58	43.84	42.97
CaO25
Na ₂ O89
K ₂ O	3.09
H ₂ O (Ign.)	4.97	2.74	2.01	2.14
TiO ₂	1.32	none	none
P ₂ O ₅05
Cr ₂ O ₃18	1.41
MnO07
Chromite5617
	100.37	100.34	99.76	100.64

GEORGIA.

Rocks collected by A. H. Brooks, who supplies the petrographic data. Hitherto unpublished. Analyses by H. N. Stokes, record No. 1727.

A. Meta-quartz-diorite, 2 miles southwest of Sweden, Gordon County. Contains plagioclase, near labradorite, green hornblende, sometimes diallage, much vitreous quartz, and accessory magnetite, ilmenite, and orthoclase. Also secondary epidote, zoisite, urallite, chlorite, garnet, calcite, and leucoxene.

B. Augite-microcline-granite, 1 mile east of Rowland, Bartow County. Contains microcline, some plagioclase, abundant pyroxene partly altered into chiefly urallite and chlorite, some biotite with frequent inclusions of rutile, much blue vitreous quartz, apatite, zircon, and magnetite.

C. Quartz-gabbro, 2 miles southeast of Walleska, Cherokee County. Closely related to B. Contains essentially plagioclase, near labradorites, and augite. Accessory magnetite, ilmenite, apatite, and zoisite. Orthoclase is sparingly present. Quartz occurs in vitreous masses.

	A.	B.	C.
SiO ₂	69.87	67.98	56.20
Al ₂ O ₃	12.02	14.84	15.46
Fe ₂ O ₃	1.42	1.00	1.54
FeO.....	3.49	3.15	9.76
MgO.....	2.30	.91	1.83
CaO.....	7.86	2.17	5.39
Na ₂ O.....	.66	2.66	2.78
K ₂ O.....	.11	4.76	2.56
H ₂ O at 110°.....	.18	.14	.16
H ₂ O above 110°.....	.89	.49	.59
TiO ₂69	.84	2.25
P ₂ O ₅17	.34	1.13
MnO.....	.16	trace	.13
BaO.....	none	.20	.17
SrO.....	none	trace	trace
Li ₂ O.....	trace	trace	none
CO ₂43	none	none
S.....	none	.08	.07
SO ₃	none	trace	trace
Cl.....	none	trace	trace
F.....	(?)	trace	trace
C (graphite).....		.21	
	100.25	99.77	100.02

KENTUCKY.

1. THE ELLIOTT COUNTY DIKE.

Described by Diller in Bull. 38. Also in Amer. Journ. Sci., 3d series, vol. 32, p. 125.

A peridotite, var. kimberlite, consisting largely of olivine, sometimes altered to serpentine. Pyrope, ilmenite, a few scales of biotite, a little enstatite, and a trace of apatite are present as primary minerals. Secondary minerals are serpentine, dolomite, magnetite, and octahedrite. P. R. C. 799.

A granitic rock occurs with the peridotite. It consists chiefly of feldspar, orthoclase, and plagioclase, with a considerable amount of quartz and ilmenite, and traces of hornblende, sphene, and apatite.

Analyses by T. M. Chatard, record Nos. 272, 273, 282, 305, 351, 352, 353, 354, and 358.

A. Granite.

B. Peridotite (kimberlite).

C. Olivine from peridotite.

D. Garnet from peridotite.

E. Ilmenite from peridotite.

	A.	B.	C.	D.	E.
SiO ₂	60.56	29.81	40.05	41.32	.76
Al ₂ O ₃	16.19	2.01	.39	21.21	2.84
Fe ₂ O ₃	5.19	5.16	2.36	4.21	9.13
FeO.....	2.41	4.35	7.14	7.93	27.81
MgO.....	1.30	32.41	46.68	19.32	8.68
CaO.....	2.09	7.69	1.16	4.94	.23
Na ₂ O.....	4.78	.11	.08	.07	} .19
K ₂ O.....	4.82	.20	.21	
H ₂ O, 110°.....	} 8.92	.14	} .17	} .20
H ₂ O, ign.....	.51		.66		
TiO ₂	1.19	2.20	.07	.16	49.32
P ₂ O ₅30	.35	.04	none	trace
Cr ₂ O ₃43	.24	.91	.74
MnO.....	.36	.23	.20	.34	.20
NiO.....05
CoO.....	trace
CO ₂	6.66	?
SO ₃28
	99.70	100.86	99.42	100.58	100.10

The following analyses are of sedimentary rocks adjoining the dike:
F. Calcareous standstone.
G. Fine-grained fissile sandstone.
H. Indurated shale.
I. Fragment of shale included in the peridotite.

	F.	G.	H.	I.
SiO ₂	60.78	60.25	41.32	35.53
Al ₂ O ₃	10.54	20.18	20.71	18.23
Fe ₂ O ₃	3.27	1.53	2.59	2.46
FeO.....		3.42	5.46	4.81
MgO.....	1.59	3.52	1.91	2.01
CaO.....	10.15	.51	9.91	21.17
Na ₂ O.....	1.41	.39	7.19	2.53
K ₂ O.....	2.36	3.17	.88	1.08
H ₂ O, 110°.....	.85	1.94		1.40
H ₂ O, ign.....	2.32	5.17	8.78	9.00
TiO ₂03	.23	.48	.95
P ₂ O ₅09	.10	.08	.08
Cr ₂ O ₃			trace	
MnO.....	.10	.10	.17	.13
CO ₂	6.29		.55	.88
	99.78	100.51	100.03	100.26

H was dried five hours at 110°, previous to analysis.

2. THE CRITTENDEN COUNTY DIKE.

A mica-peridotite, described by Diller in Amer. Journ. Sci., 3d series, vol. 44, p. 286. Contains biotite, serpentine, and perovskite, with less apatite, muscovite, magnetite, calcite, chlorite, and some other secondary products. P. R. C. 800.

A. The rock described by Diller. Analysis by W. F. Hillebrand, record No 1241.

B. An unpublished analysis of probably the same rock, from a shaft 40 feet deep at Marion. Collected by J. R. Procter, analyzed by L. G. Eakins, record number 965.

	A.	B.
SiO ₂	33.84	34.50
Al ₂ O ₃	5.88	14.37
Fe ₂ O ₃	7.04	2.85
FeO.....	5.16	4.46
MgO.....	22.96	21.81
CaO.....	9.46	11.43
Na ₂ O.....	.33	.51
K ₂ O.....	2.04	1.50
H ₂ O.....	7.50	7.14
TiO ₂	3.78	-----
F ₂ O ₅89	.77
Cr ₂ O ₃18	-----
MnO.....	.16	-----
NiO.....	.10	-----
CoO.....	trace	-----
BaO.....	.06	-----
Cl.....	.05	-----
CO ₂43	.21
SO ₃	-----	.60
	99.86	100.15

TENNESSEE.

Gabbro, 2 miles south of Limestone Cove, Unicoi County. Collected by Arthur Keith, who furnishes the petrographic data. Analysis by W. F. Hillebrand, record No. 1707. Contains hypersthene, plagioclase, and magnetite.

SiO ₂	48.11	ZrO ₂	(?)
Al ₂ O ₃	14.74	Cr ₂ O ₃01
Fe ₂ O ₃	2.54	V ₂ O ₃03
FeO.....	11.85	NiO, CoO03
MgO	5.10	MnO19
CaO.....	6.72	SrO02
Na ₂ O	2.92	BaO.....	.04
K ₂ O.....	1.92	Li ₂ O	trace
H ₂ O at 110°.....	.27	FeS ₂	a. 13
H ₂ O above 110°	1.73		99.96
TiO ₂	3.17		
P ₂ O ₅44		

a Equivalent to 0.07 S. Assumed to be pyrite, no pyrrhotite being present.

MISSOURI.

Granite and porphyry, 6 miles east of Ironton. Described by Haworth in Mo. Geol. Survey, vol. 8, Annual Report, pp. 140, 180, and 213. Analyses by W. H. Melville, record No. 1206.

A, B. Granite. A, P. R. C. 1027; B, P. R. C. 1028.

C, D. Porphyry. C, P. R. C. 1029; D, P. R. C. 1030.

Rocks composed principally of orthoclase and quartz, with some microcline, plagioclase, and biotite, and minor accessory minerals.

	A.	B.	C.	D.
SiO ₂	69.94	72.35	71.33	71.88
Al ₂ O ₃	15.19	13.78	12.55	12.88
Fe ₂ O ₃	1.88	1.87	3.75	3.05
FeO60	.36	.85	1.05
MgO92	.42	.58	.33
CaO	1.15	.87	.94	1.13
Na ₂ O	3.95	4.44	4.52	4.21
K ₂ O	4.29	4.49	4.20	4.46
H ₂ O at 100°14	.22	.12	.17
H ₂ O above 100°85	.54	.30	.26
TiO ₂25	.44	.55	.22
P ₂ O ₅13	.13	.16	.15
NiO	trace	.20	.15	.02
MnO03	.06	.04	trace
	99.32	99.87	100.04	99.81

ARKANSAS.

Ouachitite, dike near Maple Spring, 4 miles southwest of Hot Springs. Described by Kemp, in Ann. Rep. Geol. Survey Arkansas, 1890, vol. 2, p. 399. A dike rock of the monchiquite group. Contains abundant and conspicuous augite and biotite, magnetite, and minor accessory minerals in a groundmass considered by Kemp as glass. Calcite and other secondary products are also present. According to Pirsson, Journ. Geol., vol. 4, p. 679, the so-called "glass" in the monchiquites is really analcite. Analysis by L. G. Eakins, record No. 1023.

SiO ₂	36.40	K ₂ O	3.01
Al ₂ O ₃	12.94	H ₂ O	2.36
Fe ₂ O ₃	8.27	TiO ₂42
FeO	4.59	P ₂ O ₅	1.04
MgO	11.44	CO ₂	3.94
CaO	14.46		99.84
Na ₂ O97		

TEXAS.

A. Quartz-pantellarite, Vieja Mountains, San Carlos, Presidio County. Description furnished by E. C. E. Lord. Contains anorthoclase, augite, and grains of quartz in a groundmass of ægirine augite, a brown hornblende, which is probably barkevikite, orthoclase, and quartz. Magnetite and apatite are present as accessory minerals. Analysis by George Steiger, record No. 1581.

The following rocks from Uvalde County were collected by T. Way-
and Vaughan. Petrographic data furnished by Whitman Cross,
Analyses by W. F. Hillebrand, record No. 1681.

B. Plagioclase-basalt, Pinto Mountain, Brackett quadrangle. Con-
tains olivine, augite, plagioclase (labradorite), biotite, a very little
alkali feldspar (?), magnetite, and apatite. Sp. gr. 3.118, 20°. P. R.
1. 1067.

C. Basanite, Mount Inge, Uvalde quadrangle. Contains sanidine,
nephelinite, hornblende, augite, ægirine augite, olivine, magnetite, apa-
tite, and a trace of pyrite. Sp. gr. 2.770, 20°. P. R. C. 1069.

D. Rock of basaltic habit, allied to C, 1 mile northeast of Big
Mountain, Uvalde quadrangle. Contains alkali feldspar, augite, mag-
netite, and variable amounts of olivine, nephelinite, ægirite, biotite, and
eolitic minerals. Sp. gr. 2.742, 23°. P. R. C. 1068.

	A.	B.	C.	D.
SiO ₂	68.71	45.11	48.13	48.23
Al ₂ O ₃	13.45	a 12.44	18.44	17.43
Fe ₂ O ₃	5.31	2.67	3.41	2.77
FeO.....	.75	9.36	4.30	5.92
MgO.....	.19	11.56	3.06	2.99
CaO.....	.96	10.61	5.89	6.38
Na ₂ O.....	4.63	3.05	8.00	6.87
K ₂ O.....	5.51	1.01	3.80	2.78
H ₂ O at 110°.....	.13	.16	.18	.54
H ₂ O above 110°.....	.36	.78	1.59	2.84
TiO ₂21	2.34	1.74	2.00
P ₂ O ₅04	.51	.49	.69
ZrO ₂		(?)	.05	.04
Cr ₂ O ₃			none	none
V ₂ O ₃04		.04
NiO.....		.04	.02	trace
MnO.....	.14	.22	.19	.18
SrO.....	none	trace	.10	.08
BaO.....	none	trace	.10	.08
Li ₂ O.....	none	none	trace	trace
S.....		.01	.09	.08
SO ₃05			
Cl.....		.11	.29	.03
F.....		undet.	.06	undet.
	100.44	100.02	99.93	99.97
Less O.....		.02	.09	
		100.00	99.84	

a Including Cr₂O₃.

E. Phonolite, hill between Black and Big mountains, Uvalde quadrangle. Contains sanidine, nephelite, and ægirite, and very little brown hornblende, augite, and magnetite. Sp. gr. 2.559, 19.5°. P. R. C. 1070.

F. The portion of E soluble in 1:40 dilute nitric acid.

G. Nepheline-basalt, Tom Munns Hill, Uvalde quadrangle. Contains olivine, augite, nephelite, magnetite, and apatite. Sp. gr. 3.148 19°. P. R. C. 1065.

H. The portion of G soluble in 1:40 dilute nitric acid.

	E.	F.	G.	H.
SiO ₂	54.42	26.90	40.32	12.27
Al ₂ O ₃	20.76	14.34	19.46	6.09
Fe ₂ O ₃	2.64	4.75
FeO.....	1.33	.60	7.48	2.83
MgO.....	.22	(?)	18.12	6.48
CaO.....	1.34	.30	10.55	1.45
Na ₂ O.....	10.41	8.32	2.62	2.39
K ₂ O.....	4.89	1.22	1.10	.93
H ₂ O at 110°.....	.22	(?)	.57	(?)
H ₂ O above 110°.....	2.50	(?)	1.25	(?)
TiO ₂40	2.66
P ₂ O ₅11	.11	.68	.68
ZrO ₂15	none
Cr ₂ O ₃	none
NiO.....	none06
MnO.....	.1525
SrO.....	trace03
BaO.....	.0406
Li ₂ O.....	trace	trace
S.....	.0101
SO ₃03
Cl.....	.2305
F.....	none04
	99.82	51.79	100.09	33.12
Less O.....	.0503
	99.77	100.06

a Including Cr₂O₃

I. Nepheline-basalt, Black Mountain, Uvalde quadrangle. Contains olivine, augite, nephelinite, magnetite, and apatite. Sp. gr. 3.200, 21.5°. P. R. A. 1066.

J. The portion of I soluble in 1:40 dilute nitric acid.

K. Augite from I. Violet in color, very pure.

L. Nepheline-melilite-basalt, from about 3* miles southwest of Uvalde. Contains nephelinite, melilite, olivine, augite, magnetite, and apatite. Sp. gr. 3.150, 20.5°. P. R. C. 1064.

M. The portion of L soluble in 1:40 dilute nitric acid.

	I.	J.	K.	L.	M.
SiO ₂	39.92	12.00	45.23	37.96	19.32
Al ₂ O ₃	8.60	5.15	7.73	10.14	7.12
Fe ₂ O ₃	4.40	-----	2.95	3.69	-----
FeO.....	8.00	3.38	4.07	7.59	3.16
MgO.....	20.17	7.16	12.25	14.69	6.52
CaO.....	10.68	1.33	23.37	16.28	7.75
Na ₂ O.....	1.91	1.77	.47	2.18	2.11
K ₂ O.....	1.03	.77	.12	.69	.67
H ₂ O at 110°.....	.43	(?)	} .37	.39	(?)
H ₂ O at above 110°....	1.45	(?)		1.82	(?)
Ti ₂ O.....	2.70	-----	4.28	2.93	trace
P ₂ O ₅51	.51	none	1.13	1.13
ZrO ₂	none	-----	-----	none	-----
Cr ₂ O ₃14	-----	-----	.08	-----
V ₂ O ₃04	-----	-----	.05	-----
NiO.....	.06	-----	.05	.04	-----
MnO.....	.24	-----	.07	.22	-----
SrO.....	.04	-----	none	.05	-----
BaO.....	.06	-----	none	.06	-----
Li ₂ O.....	trace	-----	trace	trace	-----
SO ₃	-----	-----	-----	.03	-----
S.....	trace	-----	-----	.04	-----
Cl.....	trace	-----	-----	trace	-----
F.....	.07	-----	-----	.07	-----
	100.45	32.07	100.96	100.13	47.78
Less O.....	.03	-----	-----	.03	-----
	100.42	-----	-----	100.10	-----

LAKE SUPERIOR REGION.

1. MARQUETTE REGION, MICHIGAN.

Rocks mostly described by Van Hise and Bayley. When not otherwise specified the descriptions have been published in Mon. XXVIII, and partly in 15th Ann., p. 485.

A. Peridotite, near Opin Lake, E. $\frac{1}{4}$ sec. 27, T. 48 N., R. 27 W. Contains diallage, olivine, magnetite, and plagioclase. The diallage is partly chloritized, and the olivine is partly serpentinized. Analysis by W. F. Hillebrand, record No. 1452. P. R. C. 996.

B. Altered greenstone, Marquette district. Analysis by George Steiger, record No. 1586, hitherto unpublished. P. R. C. 988.

C. Grünerite-magnetite-schist, sec. 11, T. 47 N., R. 27 W. Mainly impure grünerite, with magnetite and quartz. Analysis by W. H. Melville, record No. 1403. P. R. C. 984.

D. Like C, from sec. 18, T. 47 N., R. 28 W. Analysis by Melville, record No. 1403. P. R. C. 980.

E. Like C and D, from sec. 20, T. 46 N., R. 30 W. Analysis by H. N. Stokes, record No. 1546. C, D, and E are similar schists, and alike in mineralogical composition.

	A.	B.	C.	D.	E.
SiO ₂	39.37	48.85	46.94	49.70	46.25
Al ₂ O ₃	4.47	15.83	.66	1.35	.92
Fe ₂ O ₃	4.96	2.50	4.51	3.10	30.62
FeO.....	9.13	10.79	33.72	37.19	16.92
MgO.....	26.53	5.82	6.64	5.72	2.13
CaO.....	3.70	6.20	3.22	.68	1.69
Na ₂ O.....	.50	2.79	.16	trace	none
K ₂ O.....	.26	1.31	none
H ₂ O at 110°.....	.87	.27	} .67	} 1.40	.42
H ₂ O above 110°.....	7.08	3.77			
TiO ₂66	1.28	none	none
P ₂ O ₅17	.22	.07	.12	.07
Cr ₂ O ₃68
MnO.....	.12	.11	.31	.93	1.01
NiO.....	.21	none
CuO.....	trace
SrO.....	trace	.09	none
BaO.....	trace	none	none
CO ₂	1.23	none	2.79	none
SO ₃06
	99.94	99.89	99.69	100.19	100.03

F. Green schist, sec. 34, T. 48 N., R. 27 W., near center of section. Contains plagioclase, chlorite, and quartz. Analysis by George Steiger, record No. 1447. P. R. C. 986.

G. Sericite-schist associated with the preceding. Mainly sericite and quartz. Analysis by Steiger, No. 1447. P. R. C. 985.

H. Dark-gray, foliated schist, SE. corner of sec. 30, T. 47 N., R. 30 W. Contains quartz, plagioclase, orthoclase, and biotite. Analysis by Steiger, No. 1525. P. R. C. 991.

I. Schistose gneiss, NW. $\frac{1}{4}$ of sec. 35, T. 47 N., R. 26 W. Contains quartz, kaolin, sericite, plagioclase, chlorite, magnetite, and apatite. Analysis by Steiger, No. 1525.

J. Novaculite from Marquette. Described by Williams in Bull. 62. Contains quartz and sericite. Analysis by W. F. Hillebrand, record No. 759.

	F.	G.	H.	I.	J.
SiO ₂	61.35	70.76	63.50	82.38	76.99
Al ₂ O ₃	16.45	14.83	17.89	11.32	13.92
Fe ₂ O ₃94	1.46	1.12	.97	.45
FeO.....	4.20	3.09	5.38	.26	.77
MgO.....	3.12	1.99	1.22	.17	1.12
CaO.....	3.46	.36	2.34	.22	.32
Na ₂ O.....	5.24	.47	2.55	.59	.56
K ₂ O.....	1.05	3.50	2.43	1.04	3.65
H ₂ O at 100°.....	.10	.09	.22	.18	} 2.35
H ₂ O above 100°.....	2.51	2.70	2.04	2.33	
TiO ₂26	.33	.62	.14	-----
P ₂ O ₅18	.26	.19	.09	trace
MnO.....	-----	-----	trace	none	trace
Li ₂ O.....	-----	-----	-----	-----	trace
CO ₂	1.98	-----	-----	-----	-----
	100.84	99.84	99.50	99.69	100.13

2. CRYSTAL FALLS DISTRICT, MICHIGAN.

Rocks described by J. Morgan Clements and H. L. Smyth, in Mon. XXXVI. A, B, C, and D by Smyth, p. 274. Analyses by H. N. Stokes, record No. 1721.

A. Granite, sec. 2, T. 41 N., R. 30 W. Contains quartz, orthoclase microcline, plagioclase, biotite, muscovite, magnetite, hematite, titanite, and occasional apatite.

B. Gneiss, sec. 35, T. 42 N., R. 29 W. Same minerals as in A.

C. Mica-schist, sec. 35, T. 42 N., R. 29 W. Contains biotite, quartz, some microcline, and magnetite.

D. Amphibolite, sec. 32, T. 42 N., R. 28 W. Contains hornblende, plagioclase, biotite, and quartz, with a little rutile and magnetite.

	A.	B.	C.	D.
SiO ₂	76.10	74.37	64.71	50.36
Al ₂ O ₃	12.95	13.34	16.43	13.26
Fe ₂ O ₃65	.92	1.83	6.30
FeO09	.21	3.84	9.34
MgO14	.27	2.97	5.55
CaO12	.50	.08	7.85
Na ₂ O	2.36	2.50	.11	2.11
K ₂ O	6.50	6.70	5.63	1.14
H ₂ O at 110°17	.12	.31	.16
H ₂ O above 110°48	.44	2.79	1.55
TiO ₂07	.07	.72	1.77
P ₂ O ₅02	.01	.02	.20
MnO	trace	trace	trace	trace
	99.65	99.45	99.44	99.59

Ba, Sr, Cl, F, S, SO₃ not looked for.

The following rocks, with one exception, are described by Clements in his portion of Mon. XXXVI:

E. Mica-diorite, sec. 28, T. 42 N., R. 32 W., southeast of Crystal Falls. Contains plagioclase (andesine), orthoclase, quartz, biotite, hornblende, and titaniferous magnetite. Analysis by Stokes, record No. 1721.

F. Hornblende-gabbro, sec. 29, T. 42 N., R. 31 W., west bank Michigamme River, east of Crystal Falls. Contains labradorite, hornblende, and iron oxide, with subordinate pyroxene, biotite, and orthoclase. Analysis by George Steiger, record No. 1712.

G. Bronzite-norite, same locality as F. Contains bronzite, hornblende, and labradorite. Analysis by Steiger, record No. 1712.

H. Wehrlite, sec. 29, T. 42 N., R. 31 W., on Michigamme River, east of Crystal Falls. Contains hornblende, pyroxene, olivine, biotite, and iron oxide. Analysis by Stokes, record No. 1721.

I. Picrite-porphry, sec. 27, T. 44 N., R. 32 W., northwest of Mansfield. Principal minerals serpentine, amphibole, chlorite, ilmenite, all but the last being secondary. Analysis by Stokes, record No. 1721.

	E.	F.	G.	H.	I.
SiO ₂	58.51	49.80	48.23	44.99	37.36
Al ₂ O ₃	16.32	19.96	18.26	5.91	4.76
Fe ₂ O ₃	2.11	6.32	1.26	3.42	6.61
FeO.....	4.43	.49	6.10	8.30	6.12
MgO.....	3.73	7.05	10.84	21.02	31.11
CaO.....	3.92	11.33	9.39	9.79	1.19
Na ₂ O.....	3.11	2.22	1.34	.91	trace
K ₂ O.....	4.08	.61	.73	.74	trace
H ₂ O at 110°.....	.23	.13	.26	.63	.65
H ₂ O above 110°.....	2.00	1.71	2.00	3.19	10.37
TiO ₂72	.79	1.00	.97	.79
P ₂ O ₅30	.07	.07	.05	.06
Cr ₂ O ₃25	.62
MnO.....	trace			trace	trace
NiO.....					.04
CO ₂	none	.15	.43	trace?	none
"	99.46	100.63	99.91	99.17	99.68

Ba, Sr, Cl, F, S, SO₃ not looked for.

J. Pre-Cambrian nonporphyritic metabasalt, from the Hemlock formation. Contains plagioclase, light-green hornblende, epidote-zoisite, chlorite, calcite, muscovite, apatite, sphene, quartz, pyrite, and magnetite.

K. Porphyritic metabasalt, Hemlock formation. Same minerals as in J, with feldspar phenocrysts.

L. Porphyritic metabasalt, like K.

M. Metadolerite, large dike in T. 43 N., R. 31 W., east of Mansfield. Hitherto unpublished. Petrographic data furnished by C. R. Van Hise. Contains uralite, labradorite, and ilmenite.

Analyses J, K, and L by H. N. Stokes, record No. 1617.

Analysis M by George Steiger, record No. 1814.

	J.	K.	L.	M.
SiO ₂	46.47	47.20	52.59	44.29
Al ₂ O ₃	16.28	15.36	15.89	17.46
Fe ₂ O ₃	3.15	3.06	6.12	3.82
FeO.....	8.96	8.87	3.96	10.35
MgO.....	6.56	4.20	5.04	7.03
CaO.....	7.90	5.05	5.55	8.68
Na ₂ O.....	3.64	4.72	5.79	2.19
K ₂ O.....	.21	1.40	.67	.71
H ₂ O at 110°.....	.28	.16	.16	.21
H ₂ O above 110°.....	3.89	3.04	2.16	4.11
TiO ₂	1.28	3.30	1.36	1.40
P ₂ O ₅13	.36	.15	.20
V ₂ O ₃04	
Cr ₂ O ₃01	none	trace	
MnO.....	.09	.20	.25	trace
CuO.....	trace	trace	trace	
SrO.....	none	trace	none	
BaO.....	none	trace	trace	none
Li ₂ O.....	trace	trace	none	
CO ₂	1.26	3.34	none	
S.....	none	trace	none	
F.....	trace	trace	trace	
	100.11	100.26	99.73	100.45

N. Adinole, sec. 8, T. 43 N., R. 31 W., near Mansfield. Contains actinolite, albite, quartz, some chlorite, and epidote. Analysis by George Steiger, record No. 1709.

O. Spilosite. Contains quartz, feldspar, chlorite, epidote, and a little biotite.

P. Spilosite. Contains quartz, feldspar, actinolite, and epidote. Analyses O and P by H. N. Stokes, record No. 1617. These three rocks are contact derivatives of the Mansfield clay slate, q. v.

	N.	O.	P.
SiO ₂	74.16	52.51	57.77
Al ₂ O ₃	11.85	19.00	19.33
Fe ₂ O ₃82	3.31	1.29
FeO	1.66	7.19	3.37
MgO	2.10	3.29	4.35
CaO.....	2.10	1.55	1.71
Na ₂ O.....	6.57	6.72	8.22
K ₂ O.....	.15	.70	.22
H ₂ O at 110°05	.34	.18
H ₂ O above 110°.....	.52	3.26	2.34
TiO ₂37	1.70	.92
P ₂ O ₅08	.15	.04
MnO06	trace	trace
SrO	none	trace	trace
BaO.....	none	trace	none
V ₂ O ₃02
Li ₂ O		trace	none
CO ₂09	none	none
C18		
F		trace	none
	100.76	99.72	99.76

3. MENOMINEE RIVER.

1ST. STURGEON FALLS GABBRO.

From Sturgeon Falls, Menominee River, sec. 27, T. 39 N., R. 29 W., Michigan. Described by Williams, Bull. 62, p. 67.

A. Saussurite-gabbro. Contains plagioclase, almost wholly altered to saussurite, diallage, hornblende, and ilmenite, with quartz, calcite, and chlorite as alteration products.

B. The same, altered and somewhat schistose. Feldspar much altered into calcite, with secondary quartz and sericite; pyroxene and hornblende changed to chlorite. Leucoxene common.

C. Light-gray, silvery schist, derived from gabbro. Contains chlorite, calcite, and a little quartz, with remnants of feldspar and some leucoxene. Analyses by R. B. Riggs, record Nos. 389, 390, 391. Material dried at 105°. TiO_2 undetermined.

	A.	B.	C.
SiO_2	51.46	38.05	45.70
Al_2O_3	14.35	24.73	16.53
Fe_2O_3	3.90	5.65	4.63
FeO	5.28	6.08	3.89
MgO	9.54	11.58	9.57
CaO	9.08	1.25	4.28
Na_2O	2.92	2.54	.55
K_2O24	1.94	3.82
H_2O	3.30	7.53	4.70
CO_220	.93	5.95
	100.27	100.28	99.62

2D. LOWER QUINNESEC FALLS, MENOMINEE RIVER.

Described by Williams, Bull. 62, pp. 89, 91.

A. Gabbro-diorite, shore below falls, Wisconsin side. Contains saussurite, hornblende, and ilmenite. Some calcite in the saussurite. Hornblende partly altered to chlorite and ilmenite to leucoxene.

B. The same, schistose form. Contains less saussurite. Hornblende and ilmenite completely altered into chlorite and leucoxene. Calcite present, and also porphyritic feldspar.

C. Silvery schist, adjoining B, and derived from gabbro. Resembles B in general, without the feldspar crystals. Calcite and sericite are present, and rutile in place of leucoxene.

D. Dark, massive greenstone. Contains hornblende, chlorite, epidote, quartz, leucoxene, with some ilmenite and traces of original feldspar.

E. Dark, schistose greenstone, forming a band in D. Chlorite entirely replaces hornblende, and rutile replaces leucoxene. Some feldspar, quartz, and calcite.

Analyses by R. B. Riggs, record Nos. 384, 385, 386, 387, 388. Material dried at 105°.

	A.	B.	C.	D.	E.
SiO ₂	47.96	49.19	46.21	43.80	44.49
Al ₂ O ₃	16.85	18.71	18.38	16.08	16.37
Fe ₂ O ₃	4.33	5.03	3.30	9.47	5.07
FeO	4.17	4.04	3.90	10.50	5.50
MgO	9.15	7.98	7.03	6.54	7.50
CaO.....	13.25	5.92	6.28	7.81	7.94
Na ₂ O	1.25	1.44	2.14	1.96	2.59
K ₂ O.....	.30	.77	.35	.34	.56
H ₂ O.....	2.89	5.05	3.82	3.99	4.99
CO ₂08	1.82	8.32	.08	5.38
	100.23	99.95	99.73	100.57	100.39

3D. UPPER QUINNESEC FALLS, MENOMINEE RIVER.

Described by Williams, Bull. 62, pp. 104, 113, 114, 121.

A. Light greenstone. Contains hornblende, feldspar much altered to saussurite, ilmenite with leucoxene border, and some secondary quartz.

B. Mica-diorite-porphyry. Mainly andesitic feldspar and biotite, with apatite, zircon, sphene, and some calcite.

C. Biotite-gneiss, Michigan side. Contains biotite, soda-orthoclase, and quartz. Sphene common, zircon and apatite present.

D. Schistose quartz-porphyry. Contains quartz, feldspar, sericite, some chlorite, anatase, and tourmaline, with zircon and apatite sparingly.

A, B, and D are from the Wisconsin side of the river.

Analyses by R. B. Riggs, record Nos. 392, 393, 394, 395. Material dried at 105°.

	A.	B.	C.	D.
SiO ₂	48.35	54.83	67.77	66.69
Al ₂ O ₃	15.40	25.49	16.61	16.69
Fe ₂ O ₃	4.04	1.61	2.06	2.06
FeO.....	4.63	1.65	1.96	.93
MgO.....	11.61	1.96	1.26	1.15
CaO.....	10.38	6.08	1.87	1.40
Na ₂ O.....	1.87	5.69	4.35	2.46
K ₂ O.....	.35	1.87	2.35	5.23
H ₂ O.....	3.60	1.18	1.69	1.70
CO ₂08	.18	.19	1.42
	100.31	100.54	100.11	99.73

4TH. SCHIST FROM THE ARAGON IRON MINE.

Talco-schist. Description, hitherto unpublished, furnished by W. S. Bayley. Analysis by George Steiger, record No. 1835. Consists mainly of talc and quartz, with a little limonite and other iron oxides.

SiO ₂	49.56	H ₂ O at 110°.....	4.50
Al ₂ O ₃	10.12	H ₂ O above 110°.....	7.66
Fe ₂ O ₃	5.87	TiO ₂60
FeO.....	.13	P ₂ O ₅04
MgO.....	20.53	MnO.....	trace
CaO.....	.72	CO ₂	none
Alkalies.....	none		99.73

4. PENOKEE-GOGEBIC REGION.

Rocks mostly described by Van Hise in Mon. XIX. Analyses A to G by T. M. Chatard, record Nos. 991, 992, 993, 994, 995.

A. Diabase, near SE. corner of sec. 13, T. 47 N., R. 46 W., Michigan. Contains plagioclase, augite, magnetite, apatite, and olivine, with some ilmenite and leucoxene. P. R. C. 1001.

B. Same dike as A, partly altered. The pyroxene is altered to amphibole, and the latter partly to biotite. Biotite has also been derived from feldspar. P. R. C. 1000.

C. Feldspar separated from A.

D. Altered diabase, Aurora mine, NE. $\frac{1}{4}$ of SW. $\frac{1}{4}$ of sec. 23, T. 47 N., R. 47 W., Michigan. P. R. C. 1002.

	A.	B.	C.	D.
SiO ₂	47.90	46.85	51.18	41.60
Al ₂ O ₃	15.60	22.62	27.00	37.20
Fe ₂ O ₃	3.69	5.12	3.19	3.21
FeO.....	8.41	1.58	undet.	.30
MgO.....	8.11	2.01	1.92	.02
CaO.....	9.99	1.25	11.70	.23
Na ₂ O.....	2.05	.80	3.48	.07
K ₂ O.....	.23	2.66	.41
H ₂ O at 105°.....	.15	3.12	} 1.19	.29
H ₂ O above 105°.....	2.34	8.25		13.54
TiO ₂82	1.12	3.79
P ₂ O ₅13	.1614
Cr ₂ O ₃	trace
MnO.....	.17	2.54	.17	.08
NiO.....	.10	.08
BaO.....	.05	.10	trace
CO ₂38	1.8938
	100.12	100.15	100.24	100.85

E. Feldspar separated from gabbro, southern half of sec. 14, T. 44 N., R. 4 W., Wisconsin.

F, G. Feldspars separated from olivine-diabase, NE. $\frac{1}{4}$ of sec. 13, T. 45 N., R. 1 W., Wisconsin.

	E.	F.	G.
SiO ₂	51.99	61.65	56.15
Al ₂ O ₃	29.32	19.91	26.05
Fe ₂ O ₃	1.23	2.28	1.98
FeO		undet.	undet.
MgO63	.61	.54
CaO	12.60	4.12	8.70
Na ₂ O	2.91	4.74	4.79
K ₂ O28	5.72	1.56
H ₂ O at 105°03	.95	.13
H ₂ O above 105°54		.64
MnO	trace	trace	.13
	99.53	99.98	100.67

The following analyses, by W. F. Hillebrand, of separations from the gabbro of Ashland County, Wisconsin, were made for the late R. D. Irving, but were never published by him:

- H. Magnetite, record No. 687. Partial analysis.
- I. Feldspar, record No. 688.
- J. Diallage, record No. 686.

For convenience I also include here, although it is decidedly out of classification—

K. Graywacke, Hurley, Wisconsin. Described by Bayley in Bull. 150, p. 84. Analysis by H. N. Stokes, record No. 1314. Contains quartz, feldspars, iron oxides, and probably kaolin. In the cement are chlorite, quartz, magnetite, pyrite, rutile, occasionally biotite, and either muscovite or kaolin. P. R. C. 20.

	H.	I.	J.	K.
SiO ₂	-----	53.30	49.80	76.84
Al ₂ O ₃	-----	29.03	2.86	11.76
Fe ₂ O ₃	50.29	.55	2.48	.55
FeO.....	30.70	.23	10.82	2.85
MgO.....	-----	.13	15.33	1.39
CaO.....	-----	11.40	16.50	.70
Na ₂ O.....	-----	4.87	.51	2.57
K ₂ O.....	-----	.40	.12	1.62
H ₂ O.....	-----	.23	.33	1.87
TiO ₂	8.77	trace	1.29	-----
P ₂ O ₅	-----	trace	trace	-----
MnO.....	-----	none	.37	trace
SrO.....	-----	trace	none	-----
BaO.....	-----	trace	none	-----
Li ₂ O.....	-----	none	trace?	-----
	89.46	100.14	100.41	100.18

5. ROCKS OF PIGEON POINT, MINNESOTA.

Described by Bayley in Bull. 109. Also partially in Amer. Journ. Sci., 3d series, vol. 37, p. 54. A and B are also described in Bull. 150, p. 274.

A. Olivine diabase. Contains labradorite, olivine, pyroxene, apatite, and magnetite. Analysis by W. F. Hillebrand, record No. 496.

B. Diallage separated from diabase. Partial analysis by R. B. Riggs, record No. 607.

C. Rock lying between the diabase and the adjacent red porphyry. Contains red feldspar, quartz, hornblende, chlorite, magnetite, apatite, and rutile. Analysis by W. F. Hillebrand, record No. 500. P. R. C. 1018.

	A.	B.	C.
SiO ₂	49.88	48.34	57.98
Al ₂ O ₃	18.55	2.90	13.58
Fe ₂ O ₃	2.06	4.68	3.11
FeO.....	8.37	14.15	8.68
MgO.....	5.77	11.34	2.87
CaO.....	9.72	15.10	2.01
Na ₂ O.....	2.59	3.56
K ₂ O.....	.68	3.44
H ₂ O.....	1.04	2.47
TiO ₂	1.19	1.98	1.75
P ₂ O ₅1629
ZrO ₂	none
MnO.....	.0913
SrO.....	trace	trace
BaO.....	.0204
Li ₂ O.....	none	trace
Cl.....	trace	trace
	100.12	98.49	99.91

D. Red soda-granite. Contains feldspar, quartz, chlorite, some muscovite, rutile, leucoxene, hematite and apatite, with sometimes secondary calcite. Granular variety. Analysis by W. F. Hillebrand, record No. 495. Dried at 110°.

E. Same as D, porphyritic variety. Called by Bayley a quartz-keratophyre. Analysis by Hillebrand, record No. 497.

F. Similar red porphyry, Little Brick Island, Lake Superior, near Pigeon Point. Consists chiefly of feldspar and quartz, with augite occasionally present. Analysis by L. G. Eakins, record No. 1059.

G. Red feldspar, anorthoclase, separated from D. Contains hematite inclusions. Analysis by J. E. Whitfield, record No. 521. Dried at 104°.

	D.	E.	F.	G.
SiO ₂	72.42	74.00	73.70	65.00
Al ₂ O ₃	13.04	12.04	12.87	18.22
Fe ₂ O ₃68	.78	3.76	2.64
FeO.....	2.49	2.61	.31	-----
MgO.....	.58	.42	.11	.06
CaO.....	.66	.85	.14	1.06
Na ₂ O.....	3.44	3.47	3.63	8.40
K ₂ O.....	4.97	4.33	4.56	4.18
H ₂ O.....	1.21	.86	.57	.46
TiO ₂40	.34	.12	-----
P ₂ O ₅20	.06	trace	-----
MnO.....	.09	.05	.07	-----
SrO.....	trace?	trace	-----	-----
BaO.....	.15	.12	-----	-----
Li ₂ O.....	trace?	trace?	-----	-----
Cl.....	trace	trace	-----	-----
	100.33	99.93	99.84	100.02

H. Dark vitreous quartzite. Contains quartz, a little red feldspar, some chlorite, some green mica, a few scales of biotite, and grains of magnetite. P. R. C. 1007.

I. Lighter interbedded quartzite. Feldspar in it much altered. P. R. C. 1004.

J. Epidotic quartzite. P. R. C. 1023.

K. Unaltered quartzite. P. R. C. 1006.

Analyses by R. B. Riggs, record Nos. 569, 571, 572, 575. Material dried at 105°.

	H.	I.	J.	K.
SiO ₂	74.22	73.65	73.14	73.64
Al ₂ O ₃	10.61	11.08	12.60	11.25
Fe ₂ O ₃	7.45	7.24	7.57	6.24
FeO.....	.85	.77	1.31	1.04
MgO.....	1.48	1.52	1.67	1.57
CaO.....	.56	.40	.43	.36
Na ₂ O.....	2.12	1.67	1.78	3.04
K ₂ O.....	1.08	1.65	1.00	1.42
Ign.....	1.79	1.88	.83	1.98
TiO ₂16	trace?	.04	trace
MnO.....	none	trace	trace	none
	100.32	99.86	100.37	100.54

Ba and Sr not looked for.

L. Altered quartzite. Contains more fibrous chlorite, some sericite, a little kalin and biotite, grains of magnetite, earthy matter, and sometimes a little calcite. Analysis by R. B. Riggs, record No. 570. P. R. C. 1009.

M. Mottled quartzite. Analysis by Riggs, record No. 573. P. R. C. 1021.

N. Purplish slate, containing many small crystals of red feldspar and scales of mica. Analysis by J. E. Whitfield, record No. 520. P. R. C. 1024.

O. Very slightly altered slate. Analysis by Riggs, record No. 576. P. R. C. 1011.

Material for analysis dried at 104° – 105° .

	L.	M.	N.	O.
SiO ₂	71.00	72.25	63.82	59.71
Al ₂ O ₃	12.88	10.73	14.65	18.32
Fe ₂ O ₃	6.69	8.01	3.16	8.11
FeO.....	.65	.38	5.12	.85
MgO.....	1.68	1.85	2.08	3.54
CaO.....	.21	.42	.70	1.05
Na ₂ O.....	1.43	2.03	1.95	1.93
K ₂ O.....	2.95	2.56	2.81	3.43
H ₂ O.....	2.03	2.05	2.62	3.24
TiO ₂44	trace	2.66	trace
P ₂ O ₅19
MnO.....	trace	trace	none	none
Li ₂ O.....			none
SO ₃33
	99.96	100.28	100.09	100.18

Ba and Sr not looked for.

P. Red granitic rock resembling D, but with more dark spots, found in contact with the purplish slate, N. Consists mainly of red feldspar, quartz, and chlorite. Analysis by J. E. Whitfield, record No. 519. Dried at 104° . P. R. C. 1014.

Q. Groundmass of red mottled quartzite. Analysis by W. F. Hillebrand, record No. 499. P. R. C. 1022.

R. Green mottlings from Q. Mostly quartz and sericite. Analysis by Hillebrand, record No. 498.

S. Another sample of the green mottlings, selected with especial care. Analysis by L. G. Eakins, record No. 1058.

T. Brilliantly red vitreous quartzite. Analysis by R. B. Riggs, record No. 574. Dried at 105° . P. R. C. 1008.

	P.	Q.	R.	S.	T.
SiO ₂	68.36	76.57	77.70	83.27	83.69
Al ₂ O ₃	13.76	9.21	7.67	7.81	7.50
Fe ₂ O ₃	2.65	1.67	3.55	1.99	1.81
FeO.....	2.75	3.94	3.29	1.81	.38
MgO.....	.68	1.51	1.83	1.59	.35
CaO.....	.70	.73	.26	.20	.39
Na ₂ O.....	3.56	3.07	1.96	.19	2.46
K ₂ O.....	4.48	1.02	1.04	1.11	2.61
H ₂ O.....	.98	1.89	^a 2.36	2.32	.72
TiO ₂	1.57	.42	.30	trace	trace?
P ₂ O ₅33	trace	none	trace
ZrO ₂	none
MnO.....	trace	.05	.04	trace
SrO.....	undet.	trace?	(?)	undet.
BaO.....	undet.	none	none	undet.
Li ₂ O.....	none	trace	trace
SO ₃66
	100.48	100.08	100.00	100.29	99.91

^a Determined by difference.

6. MISCELLANEOUS ROCKS FROM MINNESOTA.

A. Granulitic hypersthene-gabbro, from SE. $\frac{1}{4}$ sec. 20, T. 65 N., R. 4 W. Described by Bayley, *Journal of Geology*, vol. 3, p. 1. Contains hypersthene, biotite, diallage, magnetite, and plagioclase. Poor in feldspar. Analysis by H. N. Stokes, record No. 1267.

B. Hypersthene from A. Analysis by E. A. Schneider, record No. 1358.

C. Granulitic diallage-gabbro, SE. $\frac{1}{4}$ of NW. $\frac{1}{4}$ of sec. 26, T. 64 N., R. 8 W. Described by Bayley, *Journal of Geology*, vol. 3, p. 1. Contains diallage, hypersthene, magnetite, and plagioclase. Analysis by W. H. Melville, record No. 1403.

D. Olivine-gabbro, west side of Birch Lake, SE. $\frac{1}{4}$ of sec. 19, T. 63 N., R. 4 W. Described by Bayley, *Journal of Geology*, vol. 1, p. 688. Contains a large proportion of diallage and olivine. Analysis by Stokes, No. 1267.

E. Olivine from D. Analysis by W. F. Hillebrand, record No. 1308.

	A.	B.	C.	D.	E.
SiO ₂	46.96	48.44	49.56	45.66	35.58
Al ₂ O ₃	14.13	7.91	17.81	16.44	.92
Fe ₂ O ₃76	.33	2.76	.66	} 33.91
FeO.....	14.95	20.88	9.48	13.90	
MgO.....	15.97	19.35	5.93	11.57	26.86
CaO.....	2.32	1.44	9.70	7.23	.90
Na ₂ O.....	.35	2.87	2.13
K ₂ O.....	1.6841
H ₂ O at 105°.....	.07	.08	} .50	.07	.11
H ₂ O above 105°.....	1.26	none		.83	.20
TiO ₂62	undet.	.48	.92	1.22
P ₂ O ₅0367	.05
Cr ₂ O ₃	trace	trace	trace
MnO.....	.93	.92	.06	trace	.35
CoO.....	} .20
NiO.....	.0616	
CO ₂	trace
	100.09	99.35	99.82	100.03	100.25

F. Average gabbro, south quarter post, sec. 35, T. 61 N., R. 12 W. Described by Bayley in Journal of Geology, vol. 1, p. 688. Contains plagioclase, olivine, pyroxene, and magnetite. Analysis by H. N. Stokes, record No. 1267.

G. Feldspar from preceding gabbro. Analysis by W. F. Hillebrand, record No. 1308.

H. Feldspar from gabbro, east side of North Fowl Lake. Analysis by Hillebrand, No 1308.

I. Feldspar from gabbro, NW. $\frac{1}{4}$ of SE. $\frac{1}{4}$ of sec. 23, T. 62 N., R. 10.

J. Feldspar from gabbro, center of sec. 25, T. 64 N., R. 8.

K. Feldspar from gabbro, Duluth and Iron Range Railroad.

Analyses I, J, and K by Hillebrand, record No. 1115.

H, I, J, and K analyzed for W. S. Bayley, but analyses hitherto unpublished.

	F.	G.	H.	I.	J.	K.
SiO ₂	46.45	51.89	62.71	52.50	52.61	53.45
Al ₂ O ₃	21.30	29.68	19.20	30.15	29.80	29.77
Fe ₂ O ₃81	.32	1.08	.47	.57	.33
FeO.....	9.57	.37	.93	.15	.23	.15
MgO.....	7.90	.38	.81	.10	.20	.11
CaO.....	9.83	12.62	.44	12.82	12.25	11.33
Na ₂ O.....	2.14	3.87	2.96	3.72	3.80	4.33
K ₂ O.....	.34	.50	10.41	.53	.53	.68
H ₂ O at 100°...	.14	.07	.23	} .25 }	} .29 }	} .23 }
H ₂ O above 100°	1.02	.39	.92			
TiO ₂	1.19	trace	trace	trace	trace
P ₂ O ₅02
NiO.....	.04
MnO.....	trace	trace	trace
SrO.....	trace
	100.75	100.09	99.69	100.69	100.28	100.38

L. Garnetiferous gabbro, Granite Falls. Described by W. S. Bayley in Bull. 150, p. 282. Contains plagioclase, augite, garnet, magnetite, a little hornblende, some quartz grains, and apatite. Analysis by H. N. Stokes, record No. 1296. Sp. gr. 3.105. P. R. C. 109.

M. Gabbro-diorite, Minnesota Falls. Described by Bayley in Bull. 150, p. 369. Essentially plagioclase and hornblende, with some kaolin, augite, and biotite. Analysis by Stokes, No. 1296. Sp. gr. 2.935. P. R. C. 144.

N. Quartz-norite-gneiss, Odessa. Described by Bayley in Bull. 150, p. 358. Contains quartz, plagioclase, pyroxene, biotite, garnet, magnetite, and sulphide of iron. Analysis by Stokes, No. 1296. Sp. gr. 2.770.

O. Mica-schist, bed of Cross River, near Gunflint Lake. Description furnished by C. R. Van Hise. Contains biotite, quartz, feldspar (?), and pyrites. Analysis by T. M. Chatard, record No. 896.

P. Actinolite-magnetite-schist, SE. $\frac{1}{4}$ of SE. $\frac{1}{4}$ of sec. 34, T. 61 N., R. 12 W. Described by Bayley in Amer. Journ. Sci., 3d series, vol. 46, p. 178. Mainly actinolite and magnetite. Analysis by W. H. Melville, record No. 1403.

Q. Magnetite rock near the preceding, also described with P by Bayley. Analysis by Melville, No. 1403.

	L.	M.	N.	O.	P.	Q.
SiO ₂	52.31	48.29	61.04	64.77	12.35	1.16
Al ₂ O ₃	18.35	20.87	16.97	14.45	.10	1.81
Fe ₂ O ₃	5.90	1.13	-----	1.84	58.68	69.08
FeO.....	11.06	4.93	5.58	4.54	21.34	27.10
MgO.....	1.00	7.54	3.62	2.34	4.08	.25
CaO.....	7.33	14.32	5.99	2.33	1.91	.53
Na ₂ O.....	2.90	1.77	1.96	1.37	trace	-----
K ₂ O.....	.49	.38	.55	5.03	-----	-----
H ₂ O at 105°.....	} .35	} .89	} .43	.07	} .19	-----
H ₂ O above 105°.....				1.92		
TiO ₂	-----	-----	-----	.60	.12	none
P ₂ O ₅	-----	-----	-----	.20	.25	.06
MnO.....	-----	-----	-----	.11	1.22	.33
SO ₃	-----	-----	-----	.60	-----	-----
CO ₂	-----	-----	-----	.41	-----	-----
FeS.....	-----	-----	a 3.73	-----	-----	-----
	99.69	100.12	99.87	100.58	100.24	100.32

a Probably pyrrhotite.

SOUTH DAKOTA.

A. Phonolite, Black Hills. Described by Whitman Cross in Bull. 150, p. 191. Contains sanidine, nephelite, ægirite, nosean, and sodalite, with accessory sphene, apatite, and zircon, and possibly some rare zirconates or titanates. Also, sparingly, secondary zeolites and calcite. No magnetite, but minute ferritic flakes are disseminated through the rock. Analysis by W. F. Hillebrand, record No. 1618. P. R. C. 73.

B. An earlier analysis of A, by H. N. Stokes, record No. 1314. This was made along the ordinary lines as regards "completeness," as requested by the collector, and without regard to minor accessory constituents. It serves well to illustrate the difference between the two modes of treatment, in comparison with the fuller analysis.

C. Tourmaline-biotite-schist, north of Harney Peak, Black Hills. Described by Bayley in Bull. 150, p. 327. Contains quartz, biotite, tourmaline, a little garnet and rarely muscovite, iron ore, and apatite. Analysis by Stokes, record No. 1296. P. R. C. 130.

	A.	B.	C.
SiO ₂	57.86	58.60	66.77
Al ₂ O ₃	20.26	20.98	17.65
Fe ₂ O ₃	2.35	2.22	1.55
FeO.....	.39	.44	3.29
MgO.....	.04	.33	2.13
CaO.....	.89	1.13	.56
Na ₂ O.....	9.47	8.38	.99
K ₂ O.....	5.19	5.49	4.49
H ₂ O at 110°.....	.21	} 1.92	} 1.89
H ₂ O above 110°.....	2.40		
TiO ₂22
P ₂ O ₅03
ZrO ₂15
MnO.....	.21	.20
SrO.....	.04
BaO.....	.09
Li ₂ O.....	trace
SO ₃06
S.....	.03
Cl.....	.08
F.....	(?)
CO ₂	none
	99.97	99.69	99.32

WYOMING.

Rocks studied by Whitman Cross. The Leucite Hills series is described in Amer. Jour. Sci., 4th ser., vol. 4, p. 115. Analysis J is by L. G. Eakins, record No. 1078. The others are by W. F. Hillebrand, one made in the Denver laboratory, and record Nos. 343, 355, 362, 368, 403, 463, 1668, and 1682.

A. Wyomingite, Boars Tusk, Leucite Hills. Contains phlogopite, leucite, diopside, and apatite. Sp. gr. 2.779, 13.5°. P. R. C. 567.

B. Wyomingite, Fifteenmile Spring, Leucite Hills. Composition like A. Sp. gr. 2.627, 30°. P. R. C. 566.

C. Portion of B soluble in 2.5 per cent nitric acid.

D. Phlogopite from wyomingite.

E. Madupite, Pilot Butte, Leucite Hills. Contains predominating diopside and phlogopite, with perofskite and magnetite, in a glassy base which has approximately the composition of leucite. Sp. gr. 2.857, 22°. P. R. C. 576.

	A.	B.	C.	D.	E.
SiO ₂	50.23	53.70	6.08	42.56	42.65
Al ₂ O ₃	11.22	11.16	.91	12.18	9.14
Fe ₂ O ₃	3.34	3.10	} .50	2.73	5.13
FeO.....	1.84	1.21		.90	1.07
MgO.....	7.09	6.44	1.51	22.40	10.89
CaO.....	5.99	3.46	2.13	.20	12.36
Na ₂ O.....	1.37	1.67	.28	.44	.90
K ₂ O.....	9.81	11.16	1.21	10.70	7.99
H ₂ O at 110°.....	.93	.80	(?)	-----	2.04
H ₂ O above 110°.....	1.72	2.61	(?)	2.35	2.18
TiO ₂	2.27	1.92	.21	2.09	1.64
P ₂ O ₅	1.89	1.75	1.54	.06	1.52
Cr ₂ O ₃10	.04	-----	.73	.07
Di ₂ O ₃ , etc. <i>a</i>03	none	-----	-----	.11
MnO.....	.05	.04	-----	-----	.12
SrO.....	.24	.19	.10	trace	.33
BaO.....	1.23	.62	.14	1.00	.89
Li ₂ O.....	trace	trace	-----	trace	trace
SO ₃74	.06	.06	-----	.58
Cl.....	.03	.03	.03	-----	.03
F.....	.50	.44	-----	2.46	.47
	100.62	100.40	14.70	100.80	100.11
Less O.....	.22	.19	-----	1.03	.20
	100.40	100.21	-----	99.77	99.91

a Probably contains other rare earths.

F. Orendite, Fifteenmile Spring, Leucite Hills. Contains predominating leucite and sanidine, with phlogopite, a little biotite, diopside, and amphibole, and accessory apatite and rutile. Sp. gr. 2.686, 23.5°.

G. Orendite, North Table Butte, Leucite Hills. Composition like F. Sp. gr. 2.699, 19°.

H. Groundmass of F. Sp. gr. 2.615, 19°. Not cited in published paper.

I. Diopside separated from mixed wyomingite and madupite. Sp. gr. 3.290, 20°. Analyzed in the Denver laboratory. P_2O_5 included with TiO_2 .

J. Dacite?, Garfield Peak. Phenocrysts of plagioclase and hornblende, with microscopic sphene and apatite in a groundmass containing plagioclase, orthoclase, quartz, and magnetite. Sp. gr. 2.576, 26.2°. P. R. C. 159.

	F.	G.	H.	I.	J.
SiO ₂	54.08	54.17	58.13	50.86	67.78
Al ₂ O ₃	9.47	10.16	11.72	16.67
Fe ₂ O ₃	3.19	3.34	3.01	1.19	1.99
FeO.....	1.03	.65	1.01	1.82	.51
MgO.....	6.74	6.62	5.79	17.42	.71
CaO.....	3.55	4.19	2.24	23.32	2.67
Na ₂ O.....	1.39	1.21	1.36	.76	4.91
K ₂ O.....	11.76	11.91	12.58	.42	3.43
H ₂ O at 110°.....	.79	.52	.99	} 1.44
H ₂ O above 110°.....	2.71	1.01	1.16	.31	
TiO ₂	2.08	2.67	1.48	3.03
P ₂ O ₅	1.36	1.59	.3219
ZrO ₂	undet.	.22	undet.
V ₂ O ₃02
Cr ₂ O ₃07	.05
NiO.....	trace
MnO.....	.05	.06	trace	.03	trace
SrO.....	.20	.18	.10
BaO.....	.67	.59	.29
Li ₂ O.....	trace	trace	trace
SO ₃29	.16	.13
CO ₂49
Cl.....	.04	.06	trace
F.....	.49	.36	trace
	99.98	100.21	100.31	99.16	100.30
Less O.....	.21	.17
	99.77	100.04

YELLOWSTONE NATIONAL PARK AND THE ABSAROKA RANGE.

1. ELECTRIC PEAK.

Rocks described by Iddings in 12th Ann., p. 577. Also in Bull. Phil. Soc. Washington, vol. 11, p. 206. For analyses O, P, see Mon. XXXII, Pt. II.

A. Pyroxene-mica-diorite. Contains augite, hypersthene, plagioclase, quartz, biotite, hornblende, magnetite, and apatite. Analysis by J. E. Whitfield, record No. 830. P. R. C. 94.

B. Pyroxene-mica-diorite. Same minerals as under A. Analysis by Whitfield, No. 921.

C. Pyroxene-mica-diorite. Like A and B. Analysis by W. H. Melville, record No. 1231.

D. Pyroxene-porphry. Contains augite, hypersthene, biotite, magnetite, plagioclase, and quartz. Analysis by Whitfield, No. 830.

E. Quartz-pyroxene-mica-diorite. Contains biotite, hornblende, augite, hypersthene, magnetite, plagioclase, orthoclase, and quartz. Analysis by Melville, No. 1231.

	A.	B.	C.	D.	E.
SiO ₂	56.28	58.05	61.22	57.38	64.07
Al ₂ O ₃	14.23	18.00	16.14	16.86	15.82
Fe ₂ O ₃	4.69	2.49	3.01	2.49	3.40
FeO.....	4.05	4.56	2.58	5.17	1.44
MgO.....	6.37	3.55	4.21	5.51	3.39
CaO.....	7.94	6.17	5.46	7.32	4.43
Na ₂ O.....	2.98	3.64	4.48	3.33	4.06
K ₂ O.....	1.23	2.18	1.87	1.45	2.27
H ₂ O at 100°.....	.93	.86	.04	.42	.10
H ₂ O above 100°.....			.40		.42
TiO ₂84	1.05	.61	trace	.45
P ₂ O ₅40	.17	.25	trace	.18
MnO.....	.16	none	trace	trace	trace
NiO.....	-----	-----	.09	-----	.05
Li ₂ O.....	.01	none	-----	.39	-----
SO ₃	trace	.07	-----	.21	-----
Cl.....	.17	trace	-----	.17	-----
	100.28	100.79	100.36	100.70	100.08

F. Quartz-mica-diorite. Contains biotite, hornblende, augite, hypersthene, plagioclase, orthoclase, and quartz. Analysis by Whitfield, No. 921.

G. Quartz-mica-diorite. Contains biotite, hornblende, plagioclase, pyroxene, quartz, and orthoclase. Analysis by Whitfield, No. 921.

H, I, J. Quartz-mica-diorites. Contain hornblende, biotite, plagioclase, orthoclase, and quartz. Analyses by Whitfield, Nos. 921 and 830.

	F.	G.	H.	I.	J.
SiO ₂	65.11	65.60	64.85	66.05	67.54
Al ₂ O ₃	16.21	17.61	16.57	16.96	17.02
Fe ₂ O ₃	1.06	.95	2.10	2.59	2.97
FeO	3.19	2.76	2.15	1.38	.34
MgO	2.57	1.49	2.14	2.08	.13
CaO	3.97	3.72	4.01	3.37	3.36
Na ₂ O	4.00	4.36	3.71	4.20	4.62
K ₂ O	2.51	2.36	3.10	2.53	2.28
H ₂ O94	.59	.35	.69	.55
TiO ₂71	.75	.91	.34	.80
P ₂ O ₅02	.16	.14	trace	trace
MnO	none	none	none	none	trace
Li ₂ O04	.03	none	none	.03
SO ₃	trace	trace	trace	.03	.26
Cl	none	none	none	trace	.15
	100.33	100.38	100.03	100.22	100.05

K. Quartz-mica-diorite-porphry. Contains biotite, hornblende, plagioclase, orthoclase, and quartz. Analysis by Whitfield, No. 830.

L. Quartz-mica-diorite-porphry. Contains quartz, biotite, plagioclase, alkali-feldspar, and hornblende. Analysis by Whitfield, No. 830.

M. Hornblende-porphry; intrusive sheet. Contains hornblende, plagioclase, magnetite, and quartz. Analysis by Whitfield, No. 429.

N. Hornblende-mica-porphry; intrusive sheet. Contains hornblende, plagioclase, biotite, magnetite, and quartz. Analysis by Whitfield, No. 420. The Indian Creek laccolith.

O. Augite-andesite-porphry; intrusive sheet, upper portion. Contains malacolite, plagioclase, probably orthoclase, magnetite, and little biotite, with secondary chlorite or serpentine and actinolite. Analysis by Whitfield, No. 830.

P. Same sheet as O, lower portion. Analysis by W. F. Hillebrand, record No. 1571. Contains 0.04 V_2O_5 .

	K.	L.	M.	N.	O.	P.
SiO ₂	65.97	69.24	58.49	61.50	52.10	50.59
Al ₂ O ₃	16.53	15.30	16.70	17.42	16.34	11.53
Fe ₂ O ₃	2.59	1.72	3.85	4.66	3.84	1.83
FeO.....	1.72	.69	2.37	1.09	6.82	7.64
MgO.....	2.11	.95	3.12	1.26	4.33	11.27
CaO.....	3.37	2.98	5.90	5.33	4.73	8.79
Na ₂ O.....	3.41	4.46	3.47	3.99	4.02	2.27
K ₂ O.....	2.67	2.52	1.59	1.29	4.20	2.33
H ₂ O at 110°....	1.23	1.30	2.44	2.44	1.74	.21
H ₂ O above 110°						1.76
TiO ₂42	.65	1.71	none	.79	.80
P ₂ O ₅	trace	trace	trace	.60	.68	.48
MnO.....	none	trace	.24	trace	trace	.17
NiO.....						.06
SrO.....						.03
BaO.....						.10
Li ₂ O.....	.09	none	.01	.03	.13	trace
SO ₃13	.27	.63	.35	.22	none
Cl.....	.09	trace			.24	trace
O=Cl.....	100.33	100.08	100.52	99.96	100.18	99.86
	.02				.05	
	100.31				100.13	

2. SEPULCHRE MOUNTAIN.

Rocks described by Iddings, 12th Ann., p. 633. Also in Bull. Phil. Soc. Washington, vol. 11, p. 210, and in Mon. XXXII, Pt. II.

A. Pyroxene-andesite. Essentially composed of augite, hypersthene, and plagioclase. Analysis by J. E. Whitfield, record No. 923.

B. Pyroxene-andesite. Composition like A. Analysis by Whitfield, No. 923.

C. Hornblende-andesite. Essentially plagioclase and hornblende. Analysis by Whitfield, No. 922.

D. Hornblende-pyroxene-andesite. Contains augite, hypersthene, plagioclase, and hornblende. Analysis by T. M. Chatard, record No. 712.

E. Hornblende-pyroxene-andesite. Minerals as in D, with magnetite also. Analysis by Chatard, No. 712.

	A.	B.	C.	D.	E.
SiO ₂	55.83	57.17	55.92	56.61	60.30
Al ₂ O ₃	17.11	17.25	17.70	13.62	16.31
Fe ₂ O ₃	4.07	2.48	3.16	5.89	4.35
FeO.....	3.75	4.31	4.48	2.60	1.41
MgO.....	5.05	4.83	4.34	5.48	2.39
CaO.....	7.40	6.61	5.90	6.61	5.62
Na ₂ O.....	2.94	3.44	4.08	3.13	3.99
K ₂ O.....	1.71	2.03	2.24	2.71	2.36
H ₂ O at 105°.....	1.28	1.20	1.42	1.20	.64
H ₂ O above 105°.....				1.07	1.86
TiO ₂	1.05	1.03	.94	.79	.76
P ₂ O ₅21	.05	.18	.06	.20
Cr ₂ O ₃05	trace?
MnO.....	none	none	trace	.35	.13
BaO.....14	.15
SrO.....	Trace	Trace
Li ₂ O.....	none	trace	.09
SO ₃	trace	trace	trace	(?)	.10
Cl.....	none	trace	none
	100.40	100.40	100.45	100.31	100.57

F. Hornblende-mica-andesite. Contains plagioclase, hornblende, biotite, and magnetite. Analysis by Whitfield, No. 924.

G. Hornblende-mica-andesite. Contains plagioclase, hornblende, and biotite. Analysis by Chatard, No. 712.

H. Dacite. Contains plagioclase, hornblende, biotite, and quartz. Analysis by Whitfield, No. 830.

I. Dacite. Minerals as in H. Analysis by L. G. Eakins, record No. 1135.

J. Andesite-breccia. Not in paper cited. See Mon. XXXII, Pt. II. Analysis by Chatard, No. 712. Reported by Iddings as hornblende-mica-andesite, containing hornblende, plagioclase, quartz, biotite, and a little magnetite.

	F.	G.	H.	I.	J.
SiO ₂	64.27	65.50	65.66	67.49	67.95
Al ₂ O ₃	17.84	14.94	15.61	16.18	14.98
Fe ₂ O ₃	3.36	1.72	2.10	1.30	2.33
FeO.....	1.29	2.27	2.07	1.22	.95
MgO.....	2.00	2.97	2.46	1.34	1.42
CaO.....	3.42	2.33	3.64	2.68	3.98
Na ₂ O.....	3.84	5.46	3.65	4.37	4.39
K ₂ O.....	2.48	2.76	2.03	2.40	2.86
H ₂ O at 105°.....	} 1.32	.24	} 1.07	} 2.69	.37
H ₂ O above 105°.....		1.13			.61
TiO ₂32	.45	1.37	.13	.45
P ₂ O ₅16	.09	trace	.13	.07
MnO.....	none	.20	none	.08	.09
BaO.....		.13			.23
SrO.....		trace?			trace?
Li ₂ O.....	.03		.36		
SO ₃	trace	.06	.13		.11
Cl.....	none		.12		
	100.33	100.25	100.27	100.01	100.79

3. ABSAROKA RANGE.

1ST. CRANDALL BASIN.

Rocks described by Iddings in Mon. XXXII, Pt. II. The analyses also appeared in Bull. Phil. Soc. Washington, vol. 12, p. 204.

A. Gabbro-porphyry, Hurricane Ridge. Contains augite, plagioclase, hypersthene, biotite, magnetite, and a little olivine. Analysis by L. G. Eakins, record No. 1089.

B. Basalt flow, north side of Timber Creek. Rich in olivine, augite, and magnetite. Analysis by Eakins, No. 1087.

C. Basalt dike, ridge south of Hurricane Ridge. Like B. Contains also labradorite and a little orthoclase. Analysis by Eakins, No. 1087.

D. Mica-gabbro-porphyry, Hurricane Ridge. Contains plagioclase, augite, hypersthene, biotite, and magnetite. Analysis by Eakins, No. 1089.

E. Mica-gabbro, Hurricane Ridge. Like D, with some orthoclase and a little quartz and olivine. Analysis by Eakins, No. 1089.

F. Basalt-glass-breccia, ridge south of Indian Peak. A glass showing crystals of olivine, augite, plagioclase, and magnetite. Analysis by Eakins, No. 1087.

	A.	B.	C.	D.	E.	F.
SiO ₂	51.81	52.09	52.11	53.56	53.71	53.89
Al ₂ O ₃	15.24	17.84	16.58	16.07	18.00	18.81
Fe ₂ O ₃	3.66	4.27	3.66	3.21	3.99	4.92
FeO	4.86	4.56	4.99	5.29	4.05	2.81
MgO	8.89	5.33	6.87	7.23	5.19	3.29
CaO	9.06	8.03	6.43	8.77	6.88	5.42
Na ₂ O	2.83	3.39	3.25	3.06	3.50	3.65
K ₂ O	2.08	1.98	3.20	1.94	3.10	2.98
H ₂ O67	1.77	1.99	.19	.55	2.99
TiO ₂77	.39	.53	.68	.74	.49
P ₂ O ₅18	.27	.63	.18	.38	.52
MnO08	.14	.23	.11	.24	.17
	100.13	100.06	100.47	100.29	100.33	99.94

G. Orthoclase-gabbro-diorite, rich in mica, Hurricane Ridge. Contains orthoclase, plagioclase, quartz, biotite, augite, hypersthene, magnetite, and hornblende. Analysis by Eakins, No. 1089.

H. Another sample, like G. Same analyst and record number.

I. Dioritic facies of gabbro, Hurricane Ridge. Very feldspathic. Same analyst and number.

J. Monzonite, Hurricane Ridge. Contains orthoclase, plagioclase, biotite, augite, hypersthene, and magnetite. Analysis by W. H. Melville, record number 1233.

K. Augite-andesite-porphyry, intrusive sheet, Hurricane Ridge. Analysis by Eakins, No. 1088. Contains plagioclase, augite, hypersthene, magnetite, biotite, quartz, and microscopic orthoclase.

	G.	H.	I.	J.	K.
SiO ₂	55.93	56.21	57.26	57.32	57.64
Al ₂ O ₃	18.32	18.24	19.40	17.29	18.43
Fe ₂ O ₃	2.39	3.26	2.49	3.89	3.63
FeO	4.91	3.69	3.29	3.03	2.84
MgO	3.97	3.38	2.57	3.56	3.32
CaO	6.17	5.91	5.68	5.81	5.49
Na ₂ O	4.29	4.15	4.21	3.89	4.03
K ₂ O	2.62	3.02	2.95	3.04	3.33
H ₂ O at 100°22	.78	.86	.33	.51
H ₂ O above 100°30	
TiO ₂81	.88	.76	.62	.77
P ₂ O ₅56	.64	.51	.50	.34
MnO14	.17	.16	.06	.10
NiO10	
Cl					trace
	100.33	100.33	100.14	99.74	100.43

L. Hornblende-mica-andesite-porphry dike, ridge south of Hurricane Ridge. Contains plagioclase, hornblende, biotite, augite, hypersthene, and magnetite, with a little chlorite or serpentine. Analysis by Eakins, No. 1087.

M. Quartz-diorite-porphry, Hurricane Ridge. Contains plagioclase, orthoclase, quartz, augite, hypersthene, magnetite, and a little biotite. Analysis by Melville, No. 1234.

N. Quartz-mica-diorite, Hurricane Ridge. Contains andesine, orthoclase, quartz, biotite, hornblende, magnetite, and a little pyroxene. Analysis by Melville, No. 1234.

O. Quartz-mica-diorite-porphry, Hurricane Ridge. Contains andesine, orthoclase, quartz, and biotite. Analysis by Melville, No. 1234.

P. Aplite dike, Hurricane Ridge. Contains quartz, orthoclase, oligoclase, biotite, magnetite, some chlorite, and a little hornblende. Analysis by Eakins, No. 1088.

	L.	M.	N.	O.	P.
SiO ₂	61.16	63.42	63.97	64.40	71.62
Al ₂ O ₃	16.17	17.16	15.78	15.77	14.99
Fe ₂ O ₃	2.89	3.09	2.35	2.47	1.27
FeO.....	2.18	1.50	1.87	1.15	1.01
MgO.....	3.89	1.64	2.84	2.12	.74
CaO.....	4.26	4.65	3.71	3.54	1.33
Na ₂ O.....	3.87	4.51	4.36	4.10	3.62
K ₂ O.....	3.20	3.04	4.01	3.81	4.81
H ₂ O at 100°.....	2.09	.16	.09	.31	.41
H ₂ O above 100°.....		.28	.49	1.93	
TiO ₂23	.35	.48	.40	.08
P ₂ O ₅13	.26	.40	.16	trace
MnO.....	trace	.04	.05	.04	.17
NiO.....		.19	trace	.17	
Cl.....					trace
	100.07	100.29	100.40	100.27	100.05

2D. SUNLIGHT INTRUSIVES.

Descriptions, hitherto unpublished, supplied by Arnold Hague and T. A. Jaggar, jr. Analyses A, B, and C by W. F. Hillebrand, record No. 1801; D by H. N. Stokes, No. 1804.

A. Quartz-syenite, Copper Creek Basin. Contains biotite, hornblende, epidote, orthoclase, augite, titanite, magnetite, and apatite. Hornblende very pale in color.

B. Syenite-porphyry, Sulphur Creek Basin. Contains oligoclase and biotite, in a groundmass of quartz and feldspar.

C. Augite-syenite-porphyry, Copper Creek Basin. Contains augite, biotite, orthoclase, a little hornblende, and quartz.

D. Gabbro, southwest of Beams Hill, Sunlight Valley. Contains plagioclase, pyroxene, magnetite, apatite, and a little biotite.

	A.	B.	C.	D.
SiO ₂	63.07	66.64	64.40	53.57
Al ₂ O ₃	17.47	16.22	16.90	17.78
Fe ₂ O ₃	2.09	1.84	1.86	3.19
FeO.....	1.38	1.06	1.37	4.93
MgO.....	1.44	1.25	1.13	4.36
CaO.....	2.27	2.41	2.60	6.22
Na ₂ O.....	5.77	5.11	5.79	4.04
K ₂ O.....	4.59	3.86	4.56	3.04
H ₂ O at 110°.....	.25	.52	.16	.27
H ₂ O above 110°.....	.43	.55	.39	.80
TiO ₂38	.29	.23	.89
P ₂ O ₅18	.16	.21	.44
ZrO ₂	trace	.01	.02	-----
Cr ₂ O ₃	trace	trace	?	none
V ₂ O ₃	trace	.01	?	-----
NiO.....	none	none	none	none
MnO.....	.03	trace	.07	.07
SrO.....	.15	.14	.14	.13
BaO.....	.32	.27	.27	.21
Li ₂ O.....	trace?	none	trace	trace
CO ₂	none	none	none	none
FeS ₂02	trace?	trace?	-----
	99.84	100.34	100.10	99.94

3D. ISHAWOOA INTRUSIVES.

Descriptions, hitherto unpublished, supplied by Arnold Hague and T. A. Jaggar, jr. Analyses A and B by H. N. Stokes, record No. 1804; C and D by W. F. Hillebrand, No. 1765.

A. Granite-porphyry, base of Crater Mountain. Contains oligoclase, orthoclase, biotite, and quartz.

B. Diorite-porphyry, Cabin Creek. Contains plagioclase, orthoclase, quartz, and abundant hornblende. This rock is intermediate between granite-porphyry and diorite-porphyry, but nearer to the latter.

C. Diorite, base of Needle Mountain. Contains plagioclase, quartz, biotite, with subordinate hornblende and orthoclase. Structure granitic.

D. Diabase, entrance to Shoshone Canyon. Contains plagioclase, augite, and chlorite.

	A.	B.	C.	D.
SiO ₂	64.23	60.00	63.76	52.18
Al ₂ O ₃	16.34	16.37	16.01	18.19
Fe ₂ O ₃	1.07	2.28	2.22	3.31
FeO.....	1.58	2.46	1.96	4.36
MgO.....	2.47	3.81	2.43	4.69
CaO.....	3.07	4.96	4.55	6.51
Na ₂ O.....	3.49	3.73	3.98	4.58
K ₂ O.....	2.59	2.70	2.84	1.88
H ₂ O at 110°.....	.47	.61	.28	.75
H ₂ O above 110°.....	1.76	1.42	.57	2.00
TiO ₂50	.59	.52	.99
P ₂ O ₅18	.35	.25	.29
NiO.....	none	none	none	trace
MnO.....	trace	.05	.09	.14
SrO.....	.06	.11	.09	.06
BaO.....	.19	.26	.17	.11
Li ₂ O.....	trace	trace	trace	trace
CO ₂30	.17	.23	none
Cl.....	trace	trace
S.....	(?)	none
FeS ₂	1.61
	99.91	99.87	99.95	100.04

4. DIKES IN BRECCIA.

Descriptions, hitherto unpublished, supplied by Arnold Hague and T. A. Jaggar, jr. Analyses A, B, D, by H. N. Stokes, record No. 1804; C, E, F, G, H, by W. F. Hillebrand, Nos. 1765 and 1801.

A. Hornblende-pyroxene-andesite, Eagle Creek. Contains plagioclase, hornblende, pyroxene, quartz, and apatite.

B. Leucite-absarokite, Sunlight Valley. Contains phenocrysts of olivine and augite, with secondary alteration of the olivine to serpentine. The groundmass contains magnetite, augite, plagioclase, and orthoclase. Leucite abundant.

C. Gabbro-porphry, Deer Creek. Contains plagioclase, pyroxene, olivine, ilmenite, and apatite.

D. Hornblende-augite-andesite, Wind River Plateau. Contains phenocrysts of augite, hypersthene, and hornblende, plagioclase, and fine magnetite grains.

	A.	B.	C.	D.
SiO ₂	50.72	47.32	50.29	60.15
Al ₂ O ₃	16.01	11.22	15.85	17.85
Fe ₂ O ₃	4.35	2.91	8.22	2.00
FeO.....	4.20	5.81	1.43	2.02
MgO.....	7.06	15.96	4.65	3.26
CaO.....	9.02	7.11	7.71	5.48
Na ₂ O.....	2.92	1.88	2.98	3.95
K ₂ O.....	1.13	3.79	3.53	2.36
H ₂ O at 110°.....	.40	.31	1.77	.25
H ₂ O above 110°.....	2.14	1.71	1.98	1.24
TiO ₂	1.08	.75	.96	.47
P ₂ O ₅29	.61	.51	.22
ZrO ₂				
Cr ₂ O ₃	none	trace	none
V ₂ O ₃				
NiO.....	none	trace	trace	none
MnO.....	.07	.11	.15	.07
SrO.....	.09	.05	.09	.10
BaO.....	.11	.22	.15	.20
Li ₂ O.....	trace	trace	trace	none
CO ₂85	.13	none	none
Cl.....			trace
FeS ₂				
	100.44	99.89	100.27	99.62

E. Augite-andesite, Dike Mountain. Contains augite, plagioclase serpentinized olivine, magnetite, and apatite.

F. Trachyte-andesite, Dike Mountain. Contains plagioclase, orthoclase, chlorite, apatite, and magnetite. Very little augite.

G. Biotite-trachyte, Dike Mountain. Contains plagioclase, orthoclase, biotite, magnetite, and chlorite.

H. Biotite-trachyte, Dike Mountain. Contains orthoclase, plagioclase, biotite, and magnetite.

	E.	F.	G.	H.
SiO ₂	51.17	52.47	63.24	57.73
Al ₂ O ₃	16.14	18.23	17.98	18.93
Fe ₂ O ₃	4.11	3.31	2.67	1.97
FeO	4.48	3.85	.85	1.92
MgO	4.82	2.85	.63	.91
CaO	7.72	4.56	.93	2.78
Na ₂ O	2.99	4.83	6.27	5.52
K ₂ O	3.54	3.81	5.47	6.11
H ₂ O at 110°63	.68	.37	.22
H ₂ O above 110°	2.24	2.03	.80	2.93
TiO ₂	1.01	.97	.38	.33
P ₂ O ₅48	.64	.22	.25
ZrO ₂	none	.02	trace	trace
Cr ₂ O ₃	trace	trace	none	trace
V ₂ O ₃04	.03	.01	.01
NiO01	trace	none	trace ?
MnO21	.15	.04	.06
SrO10	.11	.03	.09
BaO20	.23	.25	.16
Li ₂ O	trace	trace	trace	trace
CO ₂	none	1.01	none	.26
FeS ₂05	.04	trace	.02
	99.94	99.82	100.14	100.20

4. ABSAROKITE—SHOSHONITE—BANAKITE SERIES.

Rocks from the Yellowstone Park and the Absaroka Range, described in Mon. XXXII, Pt. II, and also by Iddings in Journal of Geology, vol. 3, pp. 938, 943, 947. The rock A, from Ishawooa Canyon, is also described by Hague in Amer. Journ. Sci., 3d series, vol. 38, p. 46.

A. Leucite-absarokite, Ishawooa Canyon, Wyoming. Boulder. Contains olivine and augite in a groundmass of orthoclase and leucite. Accessory minerals, magnetite, apatite, and a few flakes of brown mica. Analysis by J. E. Whitfield, record No. 1057. Material dried at 104°.

B. Absarokite dike, head of Lamar River. Contains olivine and augite in a groundmass of orthoclase and plagioclase, with accessory magnetite, biotite, and ilmenite.

C. Absarokite dike, south of Clark Fork. Contains augite, quartz, biotite, magnetite, orthoclase, plagioclase, and sometimes analcite.

D. Absarokite lava flow, head of Raven Creek. Contains olivine, augite, orthoclase, labradorite, magnetite, apatite, and a little serpentine.

E. Absarokite dike, divide east of Cache Creek. Contains orthoclase, plagioclase, augite, biotite, magnetite, serpentine, and occasional quartz.

Analyses B, C, D, and E by L. G. Eakins, record Nos. 1086, 1365.

	A.	B.	C.	D.	E.
SiO ₂	47.28	48.95	48.36	51.76	49.71
Al ₂ O ₃	11.56	12.98	12.42	12.36	13.30
Fe ₂ O ₃	3.52	3.63	5.25	4.88	4.41
FeO.....	5.71	4.68	2.48	4.60	3.37
MgO.....	13.17	11.73	9.36	9.57	7.96
CaO.....	9.20	7.66	8.65	7.14	8.03
Na ₂ O.....	2.73	2.31	1.46	1.99	1.49
K ₂ O.....	2.17	3.96	3.97	3.83	4.81
H ₂ O.....	2.96	3.16	5.54	3.05	4.07
TiO ₂88	.49	1.18	.47	1.57
P ₂ O ₅59	.67	.84	.56	.66
Cr ₂ O ₃			trace		trace
MnO.....	.13	.13	.13	.11	.17
BaO.....			.29		.46
Cl.....	.18				
	100.08	100.35	99.93	100.32	100.01
O=Cl.....	.04				
	100.04				

F. Shoshonite lava sheet, Lamar River, south of Bison Peak. Contains plagioclase, orthoclase, augite, olivine, magnetite, and serpentine, with amygdules of zeolite and calcite. Analysis by Eakins, No. 1086.

G. Shoshonite lava sheet, southeast fork of Beaverdam Creek. Contains plagioclase, orthoclase, augite, and serpentinized olivine.

H. Leucite (?) shoshonite lava sheet, mountain east of Pyramid Peak. Contains feldspars, olivine, serpentine, augite, magnetite, brown mica, and impure leucite (?).

I. Olivine-free shoshonite dike, northeast of Indian Peak. Contains augite, plagioclase, biotite, and magnetite.

J. Shoshonite lava sheet, Two Ocean Pass. Contains orthoclase, serpentinized olivine, magnetite, augite, chlorite, biotite, apatite, labradorite. Dried at 104°.

Analyses G, H, I by Eakins, Nos. 1371, 1375, 1379. Analysis J by Whitfield, No. 906.

	F.	G.	H.	I.	J.
SiO ₂	50.06	53.49	52.49	54.86	56.05
Al ₂ O ₃	17.00	17.19	17.89	17.28	19.70
Fe ₂ O ₃	2.96	4.73	5.76	4.08	3.74
FeO.....	5.42	3.25	2.08	2.28	2.32
MgO.....	3.61	4.42	3.49	4.19	2.51
CaO.....	8.14	6.34	7.01	5.42	4.34
Na ₂ O.....	3.53	3.23	3.18	3.94	3.29
K ₂ O.....	3.40	3.86	3.73	3.96	4.44
H ₂ O.....	4.85	2.17	2.63	2.16	1.86
TiO ₂51	.71	.81	.69	.98
P ₂ O ₅66	.43	.55	.48	.66
MnO.....	.14	.14	.09	.19	trace
BaO.....		.06	.30	.37	
Li ₂ O.....					.06
SO ₃19
	100.28	100.02	100.01	99.90	100.14

K. Banakite dike, head of Lamar River. Contains augite, serpentinized olivine, orthoclase, plagioclase, biotite, magnetite, ilmenite, apatite, and analcite. Adjoins rock B. Analyses by Eakins, No. 1375.

L. Banakite dike, Hoodoo Mountain. Like K, but with amygdaloidal zeolites. Analysis by Eakins, No. 1371.

M. Banakite dike, Ishawooa Canyon, Wyoming. Like K and L, but more feldspathic. Contains a little serpentine, probably from olivine; also possibly analcite or sodalite. Analysis by Eakins, No. 1086.

N. Banakite dike, near head of Stinkingwater River. Like M, but with more serpentine. Analysis by W. H. Melville, record No. 1232.

	K.	L.	M.	N.
SiO ₂	51.82	52.63	51.46	52.33
Al ₂ O ₃	16.75	16.87	18.32	18.70
Fe ₂ O ₃	4.56	4.52	4.61	4.95
FeO.....	3.36	3.11	2.71	1.83
MgO.....	4.03	3.69	2.91	2.69
CaO.....	4.94	4.77	6.03	4.71
Na ₂ O.....	3.91	3.86	4.11	4.51
K ₂ O.....	5.02	5.17	4.48	5.45
H ₂ O at 100°.....	3.97	3.65	3.89	.74
H ₂ O above 100°.....				2.71
TiO ₂71	.81	.83	.71
P ₂ O ₅52	.63	.86	.81
MnO.....	.23	.10	.17	.03
NiO.....				.14
BaO.....	.26	.29		
	100.08	100.10	100.38	100.31

O. Leucite-banakite lava sheet, southeast fork of Beaverdam Creek. Overlies rock G. Contains olivine, augite, leucite, feldspars, magnetite, apatite, and a little brown mica. Analysis by Eakins, No. 1378.

P. Earlier unpublished analysis of O, another sample, by J. E. Whitfield, record No. 907.

Q. Quartz-banakite dike, near head of Stinkingwater River. Contains plagioclase, orthoclase, quartz, biotite, magnetite, augite, and a little calcite. Analysis by Melville, No. 1232.

R. Quartz-banakite dike, near Q. Contains plagioclase, orthoclase, quartz, biotite, magnetite, augite, chlorite, and serpentine. Analysis by Melville, No. 1232.

	O.	P.	Q.	R.
SiO ₂	52.93	51.56	57.29	60.89
Al ₂ O ₃	19.67	21.00	18.45	17.14
Fe ₂ O ₃	3.07	5.17	4.38	3.32
FeO.....	3.50	2.76	1.20	.95
MgO.....	2.88	2.52	2.08	1.16
CaO.....	4.69	4.83	3.57	3.58
Na ₂ O.....	4.20	4.37	4.43	4.54
K ₂ O.....	4.75	4.13	5.43	5.71
H ₂ O at 100°.....	2.73	2.27	.17	.39
H ₂ O above 100°.....			2.01	1.22
TiO ₂72	.65	.72	.49
P ₂ O ₅59	.69	.46	.27
MnO.....	.15	trace	trace	.09
NiO.....			.12	.19
BaO.....	.21			
Li ₂ O.....		.13		
SO ₃21		
Cl.....		trace		
	100.09	100.29	100.31	99.94

S. Absarokite, Two Ocean Pass. Contains augite, olivine, orthoclase, magnetite, serpentine, little biotite, apatite, and an isotropic substance, probably glass. Analysis by J. E. Whitfield, record No. 906.

T. Shoshonite, Beaverdam Creek. Analysis by Whitfield, No. 907. Probably the rock already represented by analysis G.

U. Shoshonite, northeast spur of Sepulchre Mountain. Contains augite and serpentinized olivine, in a groundmass of plagioclase, orthoclase, augite, magnetite, and apatite. Analysis by Whitfield, No. 908.

V. Shoshonite, Baldy Mountain, Bear Gulch, Montana. Contains hypersthene, augite, olivine, plagioclase, orthoclase, and magnetite. Analysis by Whitfield, No. 909.

	S.	T.	U.	V.
SiO ₂	51.68	52.86	51.75	54.97
Al ₂ O ₃	14.07	17.51	17.48	18.38
Fe ₂ O ₃	4.71	5.18	6.42	3.06
FeO.....	4.57	3.31	1.46	4.22
MgO.....	7.72	4.18	4.05	2.38
CaO.....	6.65	6.51	8.20	5.43
Na ₂ O.....	2.45	3.22	3.33	3.45
K ₂ O.....	4.16	3.41	3.72	3.37
H ₂ O.....	2.09	1.76	2.26	.82
TiO ₂	1.08	1.04	.86	.97
P ₂ O ₅72	.53	.67	.42
MnO.....	trace	trace	trace	trace
Li ₂ O.....	trace	.04	trace	.03
SO ₃13	.22	.17	.03
Cl.....		.16	trace	trace
CO ₂				2.92
	100.03	99.93	100.37	100.45

5. MISCELLANEOUS ROCKS. FIRST GROUP.

Collected and investigated by Arnold Hague and J. P. Iddings.

A. Black obsidian, Obsidian Cliff. Described by Iddings, 7th Ann. Contains microlites of augite and magnetite, with traces of quartz and feldspar. Analysis by J. E. Whitfield, record No. 224.

B. Red obsidian, Obsidian Cliff. Described by Iddings, 7th Ann. Like A, with ferric oxide replacing magnetite. Analysis by Whitfield, No. 223.

C. Obsidian, east of Willow Park. Black and opaque. Described by Iddings, Bull. Phil. Soc. Washington, vol. 12, p. 204. Analysis by Whitfield, No. 222.

D. Lithoidite, Obsidian Cliff. Described by Iddings, Bull. 150, p. 153. Contains quartz and sanidine, with a little magnetite and augite. Analysis by Whitfield, No. 425.

E. Rhyolite, Upper Geyser Basin. See Iddings, Bull. Phil. Soc. Washington, vol. 12, p. 204. Analysis by F. A. Gooch, record No. 114.

F. Rhyolite, Tower Creek. See Iddings, Bull. Phil. Soc. Washington, vol. 12, p. 204. Analysis by Gooch, No. 115.

	A.	B.	C.	D.	E.	F.
SiO ₂	74.70	75.52	72.59	75.50	70.92	71.85
Al ₂ O ₃	13.72	14.11	13.47	13.25	13.24	13.17
Fe ₂ O ₃	1.01	1.74	1.58	1.02	3.54	2.17
FeO.....	.62	.08	1.32	.91	.66	1.34
MgO.....	.14	.10	1.05	.07	.23	.63
CaO.....	.78	.78	2.12	.90	1.42	2.25
Na ₂ O.....	3.90	3.92	4.63	4.76	4.28	4.06
K ₂ O.....	4.02	3.63	2.52	2.85	4.25	3.89
H ₂ O.....	.62	.39	.18	.41	.57	.43
TiO ₂	none	none	.52	none	.16	.43
P ₂ O ₅	none	-----	-----	none	.18	.14
MnO.....	trace	none	none	none	.14	.12
Li ₂ O.....	-----	-----	-----	.06	none	-----
SO ₃	-----	-----	-----	.32	-----	-----
FeS ₂40	.11	.26	-----	-----	-----
	99.91	100.38	100.24	100.05	100.59	100.48

G. Rhyolite, "Great Paint Pots." Analysis by Gooch, No. 113, hitherto unpublished.

H. Rhyolite, "Elephants Back." Porphyritic obsidian. Analysis by Whitfield, No. 423, hitherto unpublished. Reported by Iddings as containing quartz, sanidine, and a little augite and magnetite, in a glassy, microlitic groundmass.

I. Rhyolite, Mount Sheridan. Composition reported by Iddings as quartz and sanidine, with a little magnetite and augite. Analysis by Whitfield, No. 426, hitherto unpublished.

J. Rhyolite, Madison Plateau. Like I. Analysis by Whitfield, No. 427, not yet published.

K. Trachytic rhyolite, Elk Creek. Analysis by Whitfield, No. 428, hitherto unpublished. Reported by Iddings as containing sanidine, labradorite, and little biotite, in a groundmass of quartz and alkali feldspar

	G.	H.	I.	J.	K.
SiO ₂	75.71	75.34	75.89	75.19	64.65
Al ₂ O ₃	11.11	12.51	12.27	13.77	17.80
Fe ₂ O ₃	1.56	.42	1.12	.61	2.33
FeO.....	.37	1.55	1.37	1.37	2.10
MgO.....	.08	.32	.29	.09	.81
CaO.....	.88	1.07	.86	.68	1.73
Na ₂ O.....	4.64	3.31	3.23	3.83	4.18
K ₂ O.....	4.18	4.17	3.42	3.33	2.83
H ₂ O.....	.35	.86	.82	.65	3.06
TiO ₂	1.25	none	.50	none	trace
P ₂ O ₅		none	none	none	trace
MnO.....	.07	.07	none	trace	trace
Li ₂ O.....		trace	.01	.02	.17
SO ₃42	.28	.29	.43
	100.20	100.04	100.06	99.83	100.09

L. Trachytic rhyolite-tuff, Two Ocean Pass. Analysis by Whitfield, No. 906, hitherto unpublished. Reported by Iddings as containing sanidine, labradorite, biotite, magnetite, and augite, in a somewhat altered, glassy groundmass. Also contains fragments of andesite.

M. Glassy trachyte, approaching rhyolite, Sunset Peak, Bear Gulch. Contains phenocrysts of sanidine, plagioclase, and biotite. See Iddings, Bull. Phil. Soc. Washington, vol. 12, p. 205; also Mon. XXXII, Pt. II. Analysis by L. G. Eakins, record No. 1378.

N. Altered rhyolite, "Iron Pot," Lower Basin. Collected by W. H. Weed. Analysis by Whitfield, No. 546, hitherto unpublished.

O. Dacite-porphry, Echo Peak.

P. Same as O, Mount Holmes. Analyses O and P by Whitfield, Nos. 421, 422, hitherto unpublished. Reported by Iddings as containing plagioclase, probably oligoclase, quartz, little biotite, and magnetite.

	L.	M.	N.	O.	P.
SiO ₂	61.15	69.45	89.20	74.51	69.54
Al ₂ O ₃	15.70	14.92	2.39	14.83	17.95
Fe ₂ O ₃	4.31	3.16	1.21	1.09	2.50
FeO.....	1.12	.23	none	trace	.22
MgO.....	3.04	.05	trace	.47	.50
CaO.....	2.84	1.19	trace	.81	1.80
Na ₂ O.....	1.54	3.19	1.11	4.38	4.30
K ₂ O.....	2.22	5.95	.79	2.72	1.21
H ₂ O.....	7.05	1.69	5.09	.92	1.96
TiO ₂69	.19	none	none	none
P ₂ O ₅75	.06	none	trace	none
MnO.....	trace	.07	-----	none	none
BaO.....	-----	.03	-----	-----	-----
Li ₂ O.....	none	-----	-----	.02	trace
SO ₃18	-----	.44	.24	.37
	100.59	100.18	100.23	99.99	100.35

Q. Mica-dacite-porphyry, Bunsen Peak. Analysis by Whitfield, No. 419, hitherto unpublished. Reported by Iddings as containing oligoclase-andesine, quartz, biotite, little magnetite, apatite, and zircon.

R. Mica-dacite-porphyry, Birch Hills. Like Q in composition. Analysis by Whitfield, No. 718, hitherto unpublished.

S. Hornblende-mica-andesite-porphyry, Fan Creek. Analysis by Whitfield, No. 713, hitherto unpublished. Reported by Iddings as containing andesine-oligoclase, hornblende, and biotite, in a ground-mass of feldspar (probably oligoclase and orthoclase), with a little magnetite, biotite, and hornblende.

T. Hornblende-mica-andesite-porphyry, Gray Peak. Analysis by Whitfield, No. 715, hitherto unpublished. Composition similar to that of S.

	Q.	R.	T.	S.
SiO ₂	70.52	70.24	65.63	65.64
Al ₂ O ₃	15.85	17.36	17.00	17.29
Fe ₂ O ₃	2.28	1.38	2.55	3.07
FeO36	.79	1.19	1.29
MgO09	.53	2.03	1.78
CaO	2.59	2.74	3.48	1.98
Na ₂ O	3.93	3.69	4.42	5.77
K ₂ O	3.43	2.65	1.64	2.44
H ₂ O35	.71	2.00	1.03
TiO ₂	trace	trace	trace	none
P ₂ O ₅17	trace	.07	.23
MnO09	none	none	trace
Li ₂ O	trace	none	.04	.04
SO ₃29	trace	trace	trace
Cl	none	trace	trace
CO ₂	none	.27	.17
	99.95	100.09	100.32	100.73

U. Hornblende-mica-andesite, Crescent Hill. Analysis by Whitfield, No. 432, hitherto unpublished. Reported by Iddings as containing oligoclase-andesine, biotite, and subordinate decomposed hornblende, in a groundmass of feldspar and quartz, with a little magnetite and biotite.

V. Hornblende-andesite, Tower Creek. Analysis by Gooch, No. 117, hitherto unpublished. Reported by Iddings as containing plagioclase, hornblende, and a little augite.

W. Pyroxene-andesite, Agate Creek. Analysis by Whitfield, No. 432, hitherto unpublished. Reported by Iddings as containing augite, hypersthene, labradorite, and magnetite, in a glassy, microlitic groundmass.

X. Rhyolitic perlite, described by Iddings, Bull. 150, p. 153. Analysis by H. N. Stokes, No. 1314, hitherto unpublished. From bluff opposite the Midway Geyser Basin. Reported by Iddings as containing quartz, sanidine, plagioclase, and rarely augite and magnetite, with microscopic zircon and apatite, in a glassy groundmass. P. R. C. 61.

	U.	V.	W.	X.
SiO ₂	64.61	61.56	61.45	73.84
Al ₂ O ₃	18.62	14.73	15.07	12.47
Fe ₂ O ₃	2.78	4.47	4.46	.32
FeO.....	.95	1.23	1.18	.90
MgO.....	.85	3.57	3.02	.25
CaO.....	4.20	4.87	5.37	1.08
Na ₂ O.....	4.37	5.10	4.00	2.88
K ₂ O.....	2.36	2.24	1.22	5.38
H ₂ O.....	.93	1.42	1.23	2.76
TiO ₂	none	.87	2.80
P ₂ O ₅30	.04	trace
MnO.....	trace	.34	none	trace
Li ₂ O.....	.0105
SO ₃	trace29
Cl.....	trace
CO ₂25
	100.23	100.44	100.14	99.88

MISCELLANEOUS ROCKS. SECOND GROUP.

Collected and investigated by Arnold Hague and J. P. Iddings.

A. Pyroxene-andesite, west of Dunraven Peak. Analysis by F. A. Gooch, record No. 116, hitherto unpublished. Reported by Iddings as containing labradorite, augite, hypersthene, and magnetite, in a micro-litic groundmass.

B. Basalt, southwest of Dunraven Peak. Analysis by Gooch, No. 118, hitherto unpublished. Reported by Iddings as containing augite, olivine, labradorite-bytownite, and magnetite, in a globulitic glassy groundmass.

C. Basalt, north spur of Prospect Peak. Analysis by J. E. Whitfield, record No. 431, hitherto unpublished. Reported by Iddings as containing labradorite, augite, olivine, and magnetite. Little glass in groundmass.

D. Basalt, Yellowstone Canyon. Analysis by Whitfield, No. 430, hitherto unpublished. Reported by Iddings as containing labradorite-bytownite, augite, olivine, magnetite, and a little brown glass.

E. Basalt, Stinkingwater Canyon. See Iddings, Bull. Phil. Soc. Washington, vol. 12, p. 205. Analysis by W. H. Melville, record No. 1232.

	A.	B.	C.	D.	E.
SiO ₂	56.47	51.70	47.17	51.70	52.37
Al ₂ O ₃	15.33	15.18	17.85	17.90	16.57
Fe ₂ O ₃	2.54	2.09	7.42	7.24	6.34
FeO.....	4.53	8.54	1.18	1.00	2.35
MgO.....	5.08	8.18	6.54	2.77	5.27
CaO.....	6.93	8.73	10.12	6.94	8.54
Na ₂ O.....	3.81	2.31	2.94	4.17	2.99
K ₂ O.....	1.66	1.81	.56	1.62	2.45
H ₂ O at 100°.....	} 1.65	} .16	} .65	} 1.15	1.18
H ₂ O above 100°.....					1.04
TiO ₂99	1.24	2.13	3.17	.73
P ₂ O ₅54	.21	.20	.41	.31
Fe, metal.....			3.26	1.81	
MnO.....	.18	trace	none	trace	.07
NiO.....					.12
Li ₂ O.....			.02	.03	
SO ₃51	.32	
S.....		.09			
Cl.....		trace			
	99.71	100.24	100.55	100.23	100.33

F. Camptonite (?), dike in Stinkingwater Canyon. See Iddings. Bull. Phil. Soc. Washington, vol. 12, p. 205. Analysis by Melville. No. 1232.

G. Kersantite, Bighorn Pass. Described by Iddings in Mon. XXXII. Pt. II. Contains hornblende, plagioclase, orthoclase, quartz, augite, biotite, magnetite, chlorite, calcite, and apatite. Augite and hornblende partly decomposed. Analysis by Whitfield, No. 714.

H. Basalt, dike, north spur of Mount Washburn. Analysis by Whitfield, No. 717, hitherto unpublished. Reported by Iddings as approaching pyroxene-andesite in composition. Contains labradorite, augite, serpentinized olivine, and magnetite, in a groundmass of globulitic and microlitic brown glass.

	F.	G.	H.
SiO ₂	50.99	48.73	53.75
Al ₂ O ₃	15.62	11.92	20.75
Fe ₂ O ₃	8.47	4.79	4.50
FeO.....	1.43	4.56	3.53
MgO.....	5.23	5.93	3.76
CaO.....	6.53	9.24	7.18
Na ₂ O.....	3.39	2.62	4.16
K ₂ O.....	3.05	2.47	1.37
H ₂ O at 100°.....	1.39	} 1.52	} 1.55
H ₂ O above 100°.....	2.48		
TiO ₂67	1.34	none
P ₂ O ₅53	.32	.15
MnO.....	trace	.36	trace
NiO.....	.07		
BaO.....		trace	
Li ₂ O.....		trace	trace
SO ₃34	trace
Cl.....		.11	none
CO ₂		5.80	none
	99.85	100.05	100.70

MONTANA.

1. MADISON AND GALLATIN VALLEYS.

Rocks collected by A. C. Peale and G. P. Merrill. Described by Merrill in Proc. U. S. National Museum, vol. 17, p. 637. See also Bull. 110, p. 47.

A. Basalt?, east side of Bozeman Creek, 2½ miles southeast of Bozeman. Contains olivine and augite, chloritized and serpentinized. The colorless groundmass contains apparently two feldspars and a pyroxene, with grains of iron oxide. Analysis by T. M. Chatard, record No. 517. Sp. gr. 2.86. P. R. C. 971.

B. Pyroxene from A. Analysis by L. G. Eakins, record No. 1046.

C. Portion of A soluble in hydrochloric acid. Contains olivine, iron oxides, and decomposition products. Analysis by Eakins, record No. 817.

	A.	B.	C.
SiO ₂	46.90	52.50	20.88
Al ₂ O ₃	10.17	2.26	3.89
Fe ₂ O ₃	1.22	2.05	2.21
FeO	5.17	2.47	4.28
MgO	20.98	17.11	16.44
CaO	6.20	21.70	1.01
Na ₂ O	1.16	.35	trace
K ₂ O	2.04	.07	trace
H ₂ O at 120°	1.04	} .64	-----
H ₂ O above 120°	4.38		
TiO ₂41	-----	-----
P ₂ O ₅44	-----	-----
Cr ₂ O ₃33	1.07	-----
MnO10	trace	trace
	100.54	100.32	48.71

D. Highly altered porphyrite?, hills 1 mile north of East Gallatin River, near camp No. 6. Rock contains hornblende paramorphs after augite in a devitrified base, with amygdules of calcite, chloritic and ferruginous matter derived from porphyritic augite and olivine. The base is also filled with needles, which may be mica. Analysis by Eakins, record No. 820. P. R. C. 968.

E. Lamprophyre, Cottonwood Creek. Contains porphyritic augite and olivine in an indeterminate groundmass carrying augite, iron oxides, and mica. Analysis by Chatard, record No. 516. P. R. C. 979.

F. Augite-porphyry, Cottonwood Creek. Contains feldspars, augite, and brown mica, with iron oxides, apatite, glass, and secondary calcite and chlorite. Carries porphyritic plagioclase and augite. No unaltered olivine visible. Sp. gr. 2.785. Analysis by Eakins, record No. 819. P. R. C. 965.

	D.	E.	F.
SiO ₂	49.47	51.65	52.33
Al ₂ O ₃	12.15	13.89	15.09
Fe ₂ O ₃	1.93	2.70	4.31
FeO.....	4.07	4.80	4.03
MgO.....	10.86	11.56	6.73
CaO.....	9.30	4.07	7.06
Na ₂ O.....	2.08	2.99	3.14
K ₂ O.....	2.42	4.15	3.76
H ₂ O at 120°.....	4.14	1.30	2.68
H ₂ O above 120°.....		1.89	
TiO ₂21	.55	.14
P ₂ O ₅37	.21	1.02
Cr ₂ O ₃	trace	.08
MnO.....	.10	.15	.09
BaO.....	.03	.19	.07
SO ₃19
CO ₂	3.31
	100.44	100.37	100.45

G. Basalt?, Bear Creek. Resembles A, but with a more crystalline groundmass. Contains plagioclase, possibly sanidine, augite, olivine, and iron oxides. Analysis by Chatard, record No. 1154. P. R. C. 967.

H. Lamprophyre, between South Boulder and Antelope creeks. Contains porphyritic augite and olivine in a feldspathic groundmass, with apatite, augite, grains of iron oxide, and shreds of brown mica. Sp. gr. 2.96. Analysis by Eakins, record No. 1266. P. R. C. 966.

I. Lamprophyre, hills east of South Boulder Creek. Nodules from a decomposed mass. Shows sanidine, plagioclase, brown mica, and altered olivine.

	G.	H.	I.
SiO ₂	49.13	50.82	50.03
Al ₂ O ₃	9.05	11.44	14.08
Fe ₂ O ₃	3.57	.25	2.92
FeO.....	5.05	8.94	6.11
MgO.....	17.21	14.01	10.73
CaO.....	5.68	8.14	7.46
Na ₂ O.....	2.01	1.79	1.46
K ₂ O.....	2.24	3.45	2.64
H ₂ O at 110°.....	.84	} .58	} 3.70
H ₂ O above 110°.....	3.50		
TiO ₂42	.59	.61
P ₂ O ₅38	.20	.42
Cr ₂ O ₃39	.03	trace
NiO.....	trace	trace	trace
MnO.....	.15	.19	.08
BaO.....	.05	.06	.04
	99.67	100.49	100.28

J. Hornblende-picrite, North Meadow Creek. Contains hornblende, abundant fresh olivine, grains of pleonaste and iron oxides, and occasionally hypersthene. Sp. gr. 3.35. P. R. C. 973.

K. Pyroxenite, divide between Meadow and Granite creeks. Contains hornblende and hypersthene, with grains of iron oxide. P. R. C. 972.

L. Hypersthene-andesite, northwest of Red Bluff. Contains plagioclase and pyroxene, with an amorphous glassy base, and sometimes olivine altered to chloritic matter.

M. Peridotite, var. wehrlite, hills 3 miles northwest of Red Bluff. Contains olivine, diallage, brown mica, rarely plagioclase, and secondary iron oxides. Sp. gr. 3.37. Analyses J, K, L, and M by Eakins, record No. 1266. P. R. C. 976.

	J.	K.	L.	M.
SiO ₂	46.13	51.83	59.48	48.95
Al ₂ O ₃	4.69	7.98	16.37	5.69
Fe ₂ O ₃73	1.48	3.21	1.20
FeO.....	16.87	8.28	3.17	12.11
MgO.....	25.17	24.10	3.29	23.49
CaO.....	4.41	5.26	4.88	5.33
Na ₂ O.....	.08	.35	3.30	1.58
K ₂ O.....	trace	.06	2.81	.79
H ₂ O.....	1.38	.29	2.01	.18
TiO ₂73	.29	.93	.81
P ₂ O ₅07	.09	.41	.12
Cr ₂ O ₃04	.31	.03	.05
MnO.....	trace	trace	.19	.08
NiO.....	.09	.11	trace	.16
BaO.....	trace13	trace
S.....	.24
	100.63	100.43	100.21	100.54
Lees O.....	.12			
	100.51			

The following rocks, at first supposed to be Pliocene sandstones, were also described by Merrill, Amer. Journ. Sci., 3d series, vol. 32, p. 119. All consist of puniceous volcanic glass. C, from Idaho, is included here for convenience. Analyses by J. E. Whitfield, record No. 382.

A. Little Sage Creek.

B. Devils Pathway.

C. Marsh Creek Valley, Idaho.

Iron and alumina weighed together. The iron is mostly in the ferrous form.

	A.	B.	C.
SiO ₂	65.56	65.76	68.92
Al ₂ O ₃ , Fe ₂ O ₃	18.24	17.18	16.22
MgO72	trace	trace
CaO	2.58	2.30	1.62
Na ₂ O	2.08	2.22	1.56
K ₂ O	3.94	3.14	4.00
H ₂ O at 105°	1.12	3.46	1.60
H ₂ O at red heat	6.50	5.60	6.00
	100.74	99.66	99.92

The following examples of volcanic dust or sand, from the Gallatin Valley, were collected by A. C. Peale. Analyses A, B, C, and D by F. W. Clarke, record No. 379. Analysis E by H. N. Stokes, record No. 1314.

A. Dry Creek Valley, above the mouth of Pass Creek.

B, C. Near Bozeman.

D. Near Fort Ellis.

E. Essentially rhyolitic glass, described by Iddings in Bull. 150, p. 146. Gallatin Valley.

	A.	B.	C.	D.	E.
SiO ₂	46.09	61.82	71.01	60.98	68.68
Al ₂ O ₃	14.35	19.86	15.17	21.69	12.69
Fe ₂ O ₃					1.14
FeO					1.17
MgO	1.29	.51	.34	1.33	1.14
CaO	1.61	1.78	1.19	1.83	1.11
Na ₂ O	1.47	2.38	2.77	.80	1.23
K ₂ O		1.31	2.97	1.23	5.58
Ignition	6.45	11.47	6.34	11.96	7.99
MnO					trace
CaCO ₃	28.72				
	99.98	99.13	99.79	99.82	100.73

2. BUTTE AND BOULDER DISTRICTS.

Rocks received from W. H. Weed and G. W. Tower. The granite rocks are described by Weed in *Journ. Geol.*, vol. 7, p. 737. All the rocks will be described in the 21st Annual Report.

A. Butte granite. A quartz-monzonite, variety banatite, from Walkerville Station. Contains quartz, andesine, orthoclase, hornblende, and biotite, with a little titanite, apatite, and magnetite.

B. Biotite from A.

C. Amphibole from A.

D. Butte granite, like A, from the Frohner mine, 10 miles south of Helena.

Analyses by H. N. Stokes, record Nos. 1686, 1787, 1808.

	A.	B.	C.	D.
SiO ₂	63.88	35.79	45.73	64.17
Al ₂ O ₃	15.84	13.70	6.77	15.25
Fe ₂ O ₃	2.11	5.22	4.94	2.16
FeO	2.59	13.72	10.39	2.98
MgO	2.13	12.13	12.32	2.60
CaO	3.97	.05	11.25	4.24
Na ₂ O	2.81	.15	.77	2.62
K ₂ O	4.23	9.09	1.22	4.34
H ₂ O at 110°22	1.21	.49	.16
H ₂ O above 110°66	3.64	2.29	.65
TiO ₂65	3.51	1.43	.67
P ₂ O ₅21	.10	.35	.16
MnO07	.19	.54	.04
SrO02	none	none	trace
BaO09	.13	none	.07
Li ₂ O	trace	trace	trace	trace
SO ₃34			.07
Cl	trace	.20		trace
F76	.28	
	99.82	99.59	98.77	100.18
Less O37	.12	
		99.22	98.65	

E. Butte granite, Gagnon mine, Butte.

F. Butte granite, Atlantic mine, Butte.

G. Butte granite, Alice mine, Butte. Analyses E, F, and G by W. H. Hillebrand, record No. 1692. Mineralogically the rocks are like A and D.

H. Weathered Butte granite.

I. Altered Butte granite. Decomposed near quartz-pyrite veins. Shows opaline silica, with sericite derived from feldspar. Hornblende gone; mica recognizable only as sericite masses having the form of biotite.

Analyses H and I by Stokes, record Nos. 1792, 1787.

	E.	F.	G.	H.	I.
SiO ₂	64.05	64.34	63.87	65.14	64.81
Al ₂ O ₃	15.38	15.72	15.39	15.63	19.44
Fe ₂ O ₃	2.20	1.62	1.93	2.37	1.82
FeO	2.74	2.94	3.08	2.13	.16
MgO	2.08	2.17	2.23	1.85	.19
CaO	4.30	4.24	4.30	3.62	.18
Na ₂ O	2.74	2.76	2.76	2.63	.21
K ₂ O	4.00	4.04	4.18	4.29	5.30
H ₂ O at 110°27	.25	.19	.37	1.41
H ₂ O above 110°83	.76	.69	.75	5.25
TiO ₂60	.53	.65	.59	.73
P ₂ O ₅21	.14	.17	.16	.10
ZrO ₂	(?)	.02	.03		
MnO11	.12	.11	trace	trace
NiO	trace ?	trace ?	trace		
SrO04	.03	.04	trace	trace
BaO08	.06	.07	.10	.10
CO ₂35	.03	.15	none	none
SO ₂05	.31
FeS ₂07	.03	.07		
Cu005	.005	.005		
	100.055	99.805	99.915	99.68	100.01

Traces of lithia in all five rocks of this table.

J. Granite, Boulder type, near Boulder. Near adamellite. Contains quartz, feldspar, black mica, and dark-green hornblende. Has an unusual amount of plagioclase.

K. Bluebird granite, Nettie mine, Butte district. An aplite. Contains orthoclase and quartz, with a little plagioclase and a sprinkling of biotite.

L. Same rock and locality as K. Analysis by W. F. Hillebrand, record No. 1692.

M. Diorite, Red Mountain, east of Butte. Contains pale-green to colorless hornblende, plagioclase varying from labradorite to albite, some orthoclase, and a little quartz. Biotite, apatite, and magnetite also occur.

N. Diorite, Red Mountain. Contact of granite with limestones. Consists mainly of green hornblende and "basic" plagioclase. Little quartz present.

O. Quartz-monzonite, Red Rock Creek, north of Butte. Contains hornblende, augite, biotite, plagioclase, and orthoclase, with a little magnetite and apatite.

Analyses J, K, M, N, and O by H. N. Stokes, record Nos. 1686, 1787.

	J.	K.	L.	M.	N.	O.
SiO ₂	67.12	76.87	77.05	56.41	49.22	61.64
Al ₂ O ₃	15.00	12.52	12.84	17.62	12.02	15.63
Fe ₂ O ₃	1.62	.67	.56	1.24	2.77	3.39
FeO	2.23	none	.14	3.55	8.80	2.69
MgO	1.74	.09	trace	3.97	9.29	2.82
CaO	3.43	.49	.57	8.66	10.56	4.90
Na ₂ O	2.76	2.47	2.81	3.25	1.90	2.64
K ₂ O	4.52	5.78	5.52	2.61	1.70	3.72
H ₂ O at 110°09	.25	.22	.14	.27	.28
H ₂ O above 110°	.58	.52	.48	.76	1.63	.91
TiO ₂48	.11	.12	.68	.95	.71
P ₂ O ₅15	.05	none	.49	.43	.21
Cr ₂ O ₃	none					trace
MnO06	trace	none	.08	trace	.04
SrO03	none	none	.08	.03	.04
BaO07	none	none	.09	.03	.08
Li ₂ O	trace	trace	trace	none	trace	trace
S		trace		none	.05	
SO ₃	trace			none	.04	none
Cl		none		.07	.08	
	99.88	99.82	100.31	99.70	99.77	99.70

P. Pitchstone, or rhyolite-dacite obsidian, Butte. Phenocrysts of andesine, quartz, and orthoclase in a glassy base.

Q. Rhyolite, Hyde Park dike, Butte district. Contains sanidine, quartz, plagioclase, and biotite in a groundmass of quartz and feldspar.

R. Rhyolite, top of Red Mountain, Rimini. Quartz and feldspar phenocrysts in a groundmass of the same minerals.

S. Quartz-porphyry, Modoc mine, Butte district. Contains quartz, orthoclase, and plagioclase in a groundmass of quartz and feldspar. A few small biotites are present. Feldspar partly altered to sericite.

T. Andesite-porphyry, Hiawatha Creek, above basin. Contains plagioclase, orthoclase, and augite.

U. Andesite-porphyry, Zosel mining district, near Daylight. Somewhat altered. Contains augite, andesine, and olivine in a groundmass of plagioclase, augite and iron oxide. Ferromagnesian minerals altered to serpentine.

Analysis S by Hillebrand, record No. 1692. P, Q, R, T, and U by Stokes, record Nos. 1686, 1787.

	P.	Q.	R.	S.	T.	U.
SiO ₂	67.55	74.34	75.30	69.95	62.53	54.61
Al ₂ O ₃	15.68	12.97	11.95	15.14	19.01	15.23
Fe ₂ O ₃98	.75	} 2.17	.38	1.96	3.51
FeO	1.02	.54		.83	1.44	4.80
MgO	1.11	.86	.05	.56	1.29	4.69
CaO	2.51	.85	.62	1.45	5.17	7.41
Na ₂ O	4.15	2.49	3.09	2.70	3.45	1.46
K ₂ O	2.86	4.72	4.96	6.36	3.30	2.70
H ₂ O at 110°38	1.03	.36	.40	.21	.32
H ₂ O above 110°	2.76	1.11	.61	.91	.45	2.47
TiO ₂34	.18	.17	.24	.65	.86
P ₂ O ₅12	.07	trace	.10	.17	.35
ZrO ₂	none	.05	-----	.02	-----	none
MnO	trace	trace	trace	.08	.03	.09
SrO03	trace	none	.02	.04	.04
BaO11	.07	trace	.13	.13	.11
Li ₂ O	none	trace	trace	trace	trace	trace
CO ₂	none	none	none	.37	none	1.46
SO ₃	none	.03	.44	-----	none	none
FeS ₂	-----	-----	-----	a .39	-----	-----
Cl05	none	trace	undet.	-----	none
Cu	-----	-----	-----	.03	-----	-----
Organic matter	-----	-----	.45	-----	-----	-----
	99.65	100.06	100.17	100.06	99.83	100.11

a Other sulphides than pyrite present.

3. CRAZY MOUNTAINS.

The greater number of the rocks in this group were received from J. E. Wolff, who has supplied the petrographic data. The few exceptions are properly indicated. The three tinguaïtes and the elaeolite-syenite were described by Wolff and Tarr in Bull. Mus. Compar. Zoölogy, vol. 16, No. 12, 1893, in a paper upon "Acmite-Trachyte from the Crazy Mountains." The renaming of these rocks is due to Wolff. The analyses, except when otherwise stated, are by W. F. Hillebrand, record No. 1438.

A. Granitite, Big Timber Creek. Small dikes in main diorite stock. Contains quartz, orthoclase, plagioclase (oligoclase), and biotite.

B. Hornblende-granitite, main stock, Big Timber Creek. Contains biotite, hornblende, quartz, orthoclase, and plagioclase (oligoclase).

C. Granite-porphýry, intrusive sheet, north part of the mountains. Contains orthoclase, albite, quartz, biotite, and green hornblende in a groundmass of quartz and untwinned feldspar.

D. Porphyrite, intrusive sheet, Sweet Grass Creek. Contains hornblende and andesine in a groundmass of plagioclase, orthoclase, and quartz, with accessory magnetite.

E. Porphyrite, dike in contact zone, Sweet Grass Creek. Contains brown hornblende, biotite, and labradorite in a groundmass of plagioclase, biotite, and hornblende, with a little quartz and orthoclase.

F. Porphyrite, intrusive sheet, middle peak of Three Peaks. Contains andesine, hornblende, augite, and biotite in a groundmass of plagioclase, orthoclase, and quartz.

	A.	B.	C.	D.	E.	F.
SiO ₂	74.37	64.47	69.93	66.28	64.49	61.08
Al ₂ O ₃	13.12	15.45	14.95	16.21	17.25	16.62
Fe ₂ O ₃73	2.25	1.78	.80	.86	2.87
FeO.....	.87	2.25	.55	2.06	2.42	2.56
MgO.....	.35	2.68	.60	1.57	1.24	1.65
CaO.....	1.26	3.63	1.46	3.53	3.79	3.66
Na ₂ O.....	2.57	4.54	5.30	4.36	4.19	4.75
K ₂ O.....	6.09	3.19	3.99	3.20	4.15	3.90
H ₂ O at 110°...	.05	.05	.12	.12	.06	.44
H ₂ O above 110°	.25	.63	.32	.78	.54	.97
TiO ₂29	.75	.33	.50	.51	.73
P ₂ O ₅06	.22	.33	.20	.23	.63
MnO.....	trace	.06	trace	trace	trace	trace
SrO.....	trace	.04	.06	.05	.08	.08
BaO.....	.10	.23	.29	.34	.30	.32
Li ₂ O.....	trace	trace?	trace	trace?	trace	trace
	100.11	100.44	100.01	100.00	100.11	100.26

G. Porphyrite, intrusive sheet, north of Shields River Basin. Contains plagioclase, hornblende, and biotite, in a groundmass of plagioclase, a little orthoclase, augite, hornblende, and magnetite.

H. Porphyrite, intrusive sheet, northern part of the mountains. Contains brown hornblende, green augite, and plagioclase, in a groundmass of plagioclase, augite, and magnetite, with accessory apatite.

I. Syenite, ridge north of Shields River Basin. Contains hornblende, green augite, and anorthoclase, with accessory sphene, apatite, and magnetite.

J. Diabase-porphyrite (?), dike south of Shields River Basin. Contains labradorite and decomposed augite in a groundmass of plagioclase, epidote, and chlorite.

K. Diorite-porphyrite, Big Timber Creek. Contains labradorite, augite, hornblende, biotite, orthoclase, quartz, magnetite, and apatite.

L. Quartz-diorite, main stock on Sweet Grass Creek. Contains hornblende, biotite, augite, labradorite, orthoclase, and quartz, with accessory apatite, magnetite, olivine, and hypersthene.

	G.	H.	I.	J.	K.	L.
SiO ₂	56.75	54.69	58.28	58.28	54.56	53.48
Al ₂ O ₃	16.40	16.53	17.89	19.37	17.58	19.35
Fe ₂ O ₃	4.78	4.54	3.20	1.35	4.30	2.37
FeO.....	3.10	2.83	1.73	2.98	4.98	4.90
MgO.....	3.22	2.99	1.51	1.30	2.86	3.67
CaO.....	5.34	5.34	3.69	4.78	6.00	7.55
Na ₂ O.....	4.19	5.19	5.89	4.40	4.43	4.07
K ₂ O.....	3.36	3.93	5.34	3.75	2.70	1.41
H ₂ O at 110° ..	.40	.32	.17	.44	.02	.16
H ₂ O above 110°	.82	1.05	.98	1.78	.38	.80
TiO ₂86	.91	.64	.96	1.34	1.07
P ₂ O ₅52	.73	.26	.35	.60	.62
MnO.....	.17	.07	.06	.07	.06	.06
SrO10	.06	.05	.09	.08	.11
BaO.....	.33	.37	.36	.25	.27	.19
Li ₂ O.....	trace?	trace?	trace	trace?	trace	trace?
Cl.....	trace
CO ₂833308
	100.34	100.38	100.05	100.48	100.16	99.89

M. Diorite, head of Rock Creek. Contains biotite, labradorite, and augite, in a groundmass of plagioclase, orthoclase, and quartz, with accessory magnetite, apatite, and hornblende.

N. Diorite, main stock, Big Timber Creek. Contains biotite, augite, labradorite, quartz, orthoclase, apatite, and magnetite.

O. Olivine-gabbro, Big Timber Creek. Contains labradorite (?), brown hornblende, augite, olivine, and magnetite.

P. Rock from Musselshell River, north of Crazy Mountains. Received from J. S. Diller, but undescribed. Regarded by W. H. Weed as monchiquite. Analysis by L. G. Eakins, record No. 1021.

Q. Hornstone, metamorphosed shale, contact zone, Sweet Grass Creek. An aggregate of augite, quartz, triclinic feldspar, and biotite.

	M.	N.	O.	P.	Q.
SiO ₂	57.97	50.73	40.42	44.66	57.31
Al ₂ O ₃	15.65	19.99	9.98	12.12	14.24
Fe ₂ O ₃73	3.20	9.83	5.81	1.00
FeO.....	2.80	4.66	10.67	3.20	3.24
MgO.....	4.96	3.48	11.56	8.77	4.60
CaO.....	10.93	8.55	10.78	8.14	11.31
Na ₂ O.....	3.03	4.03	1.26	4.47	2.64
K ₂ O.....	3.16	1.89	.60	2.75	4.55
H ₂ O at 110°.....	.22	.11	.45	} 4.33	.25
H ₂ O above 110°.....	.38	.66	1.17		.24
TiO ₂60	1.59	2.51	1.02	.52
P ₂ O ₅15	.81	.63	2.02	.18
(CoNi)O.....			.02		
MnO.....	trace	.05	.25	.21	.08
SrO.....	.02	.11	.02		trace?
BaO.....	.09	.27	.05		.19
Li ₂ O.....	trace	trace	trace		trace?
CO ₂				2.19	.17
Cl.....	trace				
F.....	trace				
	100.69	100.13	100.20	99.69	100.52

R. Elæolite-syenite, Peaked Butte, northeast side of the mountains. Described by Wolff and Tarr, *l. c.* Contains anorthoclase, augite, occasionally sodalite, ægirine, apatite, magnetite, and some interstitial nephelite. Analysis by W. H. Melville, record No. 1291.

Ra. Anorthoclase from R. Analysis by W. F. Hillebrand, record No. 1297.

S. Tinguaitite, var. sölvbergite, intrusive sheet north of Shields River.

T. Tinguaitite, var. sölvbergite, dike north part of mountains.

U. Tinguaitite, var. sölvbergite, dike at head of Sixteenmile Creek.

S, T, and U are the rocks described by Wolff and Tarr in their paper upon "Acmite-Trachyte." Mineral composition the same as under R. Analyses by W. H. Melville, record No. 1291.

	R.	Ra.	S.	T.	U.
SiO ₂	59.66	62.31	58.70	62.17	64.33
Al ₂ O ₃	16.97	22.63	19.26	18.58	17.52
Fe ₂ O ₃	3.18	3.37	2.15	3.06
FeO	1.1558	1.05	.94
MgO8076	.73	.34
CaO	2.32	.63	1.41	1.57	.56
Na ₂ O	8.38	7.68	8.55	7.56	7.30
K ₂ O	4.17	4.79	4.53	3.88	4.28
H ₂ O at 105°07	.16	.07	.07	.04
H ₂ O above 105°	2.53	.72	2.57	1.63	.95
TiO ₂	trace	trace	trace	trace
P ₂ O ₅1410	.11	trace
MnO1910	trace	.35
SrO57
BaO77
	99.56	100.26	100.00	99.50	99.67

V. Theralite, Gordons Butte. Contains green augite, ægirine, biotite, olivine, nephelite, sodalite, and a feldspar, partly sanidine, containing K, Na, Ba, Sr, and Ca; also accessory apatite, magnetite, and sphene.

W. Theralite, Gordons Butte; another sample. Analysis by E. A. Schneider, record No. 1281.

X. Theralite, north of Alabaugh Creek. Described by Wolff for the Educational Series. Contains augite, ægirine, biotite, olivine, magnetite, apatite, nephelite partly zeolitized, a mineral of the sodalite group, sanidine, and analcite. Analysis by Schneider, No. 1281.

Rocks V, W, and X are described by Wolff in Bull. 150, pp. 197, 199.

Y. Altered theralite, head of Shields River, west of Loco Mountain. Received from W. H. Weed. Analysis by H. N. Stokes, record No. 1547.

Z. Hornblende-picrite, Conical Peak. Data supplied by J. P. Iddings. Contains hornblende, plagioclase, hypersthene, augite, olivine, very little primary quartz, and probably some magnetite and apatite. Analysis by L. G. Eakins, record No. 1379.

	V.	W.	X.	Y.	Z.
SiO ₂	44.65	44.31	47.67	48.90	45.71
Al ₂ O ₃	13.87	17.20	18.22	14.70	10.80
Fe ₂ O ₃	6.06	4.64	3.65	4.14	4.43
FeO	2.94	3.73	3.85	3.68	9.35
MgO	5.15	6.57	6.35	3.95	13.75
CaO	9.57	10.40	8.03	8.26	10.48
Na ₂ O	5.67	4.45	4.93	5.22	1.58
K ₂ O	4.49	3.64	3.82	.56	.85
H ₂ O at 110°96	.77	.38	.52	} .97
H ₂ O above 110°	2.10	3.30	2.97	2.44	
TiO ₂95	undet.	undet.	.95	1.83
P ₂ O ₅	1.5079	.11
Cr ₂ O ₃10
MnO17	.10	.28	.03	.17
SrO3713
BaO7631	trace
Li ₂ O	trace	trace
CO ₂11	5.42
SO ₃6104
Cl	trace
	99.92	99.11	100.15	100.04	100.13

4. LITTLE BELT MOUNTAINS.

Rocks collected by W. H. Weed and L. V. Pirsson, who furnish the petrographic data. To be described in a paper on the Neihart and Barker mining districts in Pt. III of the 20th Ann. Analyses by W. F. Hillebrand, record No. 1476, and H. N. Stokes, record No. 1547.

A. Quartz-porphry, Yogo Peak, sheet at head of Belt and Running Wolf creeks. Phenocrysts of orthoclase and quartz in a groundmass of quartz and alkali feldspar, with a little white mica and some kaolin. Chlorite, limonite, and calcite are also present, pseudomorphous after biotite and perhaps hornblende. Total amount of secondary minerals very small. Analysis by Hillebrand.

B. Granite-porphry, Wolf Butte. Phenocrysts of quartz, orthoclase, plagioclase, and biotite in a groundmass of quartz and alkali feldspar. A little apatite and iron ore, with secondary calcite, limonite, chlorite, and white mica. Analysis by Stokes.

C. Granite-porphry, top of Barker Mountain. Phenocrysts of orthoclase, oligoclase, biotite, green hornblende, sphene, and iron ore, in a groundmass of quartz and alkali feldspar. Also a little apatite and some secondary chlorite and limonite. Analysis by Stokes.

D. Syenite, Wright and Edwards mine, Hughesville, near Barker. Very fresh rock. Contains magnetite, ilmenite, hornblende, anorthoclase, albite, subordinate quartz, a little chlorite, calcite, and limonite, and white mica in traces. Analysis by Stokes.

E. Syenite, Yogo Peak. Described in Amer. Journ. Sci., 3d series, vol. 50, p. 471. Contains apatite, sphene, iron ore, pyroxene, hornblende, biotite, orthoclase, oligoclase, and quartz, with traces of chlorite and limonite and a little kaolin. Analysis by Hillebrand.

F. Syenite-porphry, dike at head of Sheep Creek. Phenocrysts of orthoclase, some plagioclase, and green hornblende, in a groundmass of alkali feldspar, with some quartz. Also contains a little apatite and iron ore, with some secondary calcite and kaolin. Analysis by Hillebrand.

	A.	B.	C.	D.	E.	F.
SiO ₂	73.12	69.68	68.60	64.64	61.65	66.29
Al ₂ O ₃	14.27	14.97	16.13	16.27	15.07	15.09
Fe ₂ O ₃51	.79	2.22	2.42	2.03	1.37
FeO.....	.26	.34	.44	1.58	2.25	1.17
MgO.....	.24	.66	.72	1.27	3.67	2.39
CaO.....	1.10	2.10	1.36	2.65	4.61	2.38
Na ₂ O.....	3.43	3.38	4.37	4.39	4.35	3.96
K ₂ O.....	4.90	4.40	4.89	4.98	4.50	4.91
H ₂ O at 110°...	.68	1.09	.20	.09	.26	.39
H ₂ O above 110°	.73	.92	.58	.27	.41	.60
TiO ₂08	.28	.32	.51	.56	.27
P ₂ O ₅03	.17	.18	.37	.33	.15
Cr ₂ O ₃	none	-----	-----	-----	trace	none
MnO.....	.06	trace	trace	trace	.09	.06
SrO.....	trace	.06	.09	.08	.10	.07
BaO.....	trace	.14	.27	.18	.27	.30
Li ₂ O.....	trace	trace	trace	-----	trace	trace
CO ₂77	.88	-----	.37	-----	.45
SO ₃	-----	trace	trace	trace	-----	-----
Cl.....	-----	trace	trace	.05	-----	-----
	100.18	99.86	100.37	100.12	100.15	99.85

G. Syenite-porphyry, intrusive sheet, between Yogo Peak and Big Baldy Mountain. Abundant phenocrysts of hornblende and orthoclase, with less biotite and plagioclase, in a groundmass of alkali feldspar with accessory quartz. Also contains iron ore and apatite, with secondary calcite, chlorite, sericite, and kaolin. Analysis by Hillebrand.

H. Granite-syenite-porphyry, north end of Thunder Mountain. Phenocrysts of orthoclase, oligoclase, hornblende, and biotite, in a groundmass of alkali feldspar and very abundant quartz. Also contains a little sphene, iron ore, and apatite, very little secondary chlorite and limonite, and a trace of kaolin. Analysis by Stokes.

I. Quartz-syenite-porphyry, top of Big Baldy Mountain. Phenocrysts of orthoclase, plagioclase, biotite, iron ore, brown hornblende, and sphene, in a groundmass of quartz and alkali feldspar. Also contains a little apatite, with traces of chlorite, limonite, and kaolin. Analysis by Hillebrand.

J. Syenite-diorite-porphyry, talus slope on west side of Bear Park. Phenocrysts of biotite, hornblende, plagioclase, and orthoclase, in a groundmass of quartz and alkali feldspar. Also contains a little magnetite, chlorite, white mica, and apatite. Analysis by Stokes.

K. Diorite-porphyry, Steamboat Mountain. Phenocrysts of orthoclase, plagioclase, hornblende, and biotite, with a little iron ore and apatite, in a groundmass of orthoclase, plagioclase, and quartz. Analysis by Stokes.

L. Diorite, Carpenter Creek, near Neihart. Contains green hornblende, biotite, and plagioclase, with some apatite, iron ore, calcite, kaolin, and muscovite, and a very little quartz and orthoclase. Analysis by Hillebrand.

	G.	H.	I.	J.	K.	L.
SiO ₂	62.58	67.44	67.04	64.95	62.18	55.13
Al ₂ O ₃	16.42	15.78	15.25	15.44	15.77	20.27
Fe ₂ O ₃	2.46	1.58	1.69	2.02	1.83	1.52
FeO.....	1.96	.85	1.13	1.60	2.44	4.29
MgO.....	1.84	1.43	1.75	2.65	3.55	1.80
CaO.....	2.47	2.38	2.17	3.07	4.13	7.05
Na ₂ O.....	4.57	4.11	4.09	4.25	3.92	4.31
K ₂ O.....	3.91	4.87	5.10	3.87	3.91	2.84
H ₂ O at 110°...	.38	.32	.56	.26	.30	.14
H ₂ O above 110°	1.40	.70	.51	.85	.70	.95
TiO ₂40	.32	.20	.39	.55	.74
P ₂ O ₅33	.21	.21	.25	.32	.40
MnO.....	.08	trace	.05	trace	trace	.13
SrO.....	.10	.09	.03	.10	.16	.06
BaO.....	.41	.24	.33	.35	.43	.11
Li ₂ O.....	trace	trace	trace?	-----	trace	trace
CO ₂77	-----	-----	-----	-----	.26
SO ₃	-----	trace	-----	.02	trace	-----
Cl.....	-----	trace	-----	.04	.04	-----
	100.08	100.32	100.11	100.11	100.23	100.00

M. Minette, intrusive sheet, head of Sheep Creek. Chiefly biotite augite, and orthoclase, with accessory apatite, plagioclase, quartz, and iron ore, and some secondary calcite, chlorite, and kaolin. Analysis by Hillebrand.

N. Monzonite, Yogo Peak. Described in Amer. Journ. Sci., 3d series, vol. 50, p. 473, and 4th series, vol. 1, p. 356. Contains apatite sphene, iron ore, pyroxene, hornblende, biotite, orthoclase, and oligoclase, and a little secondary kaolin. Analysis by Hillebrand.

O. Monchiquite, dike on Big Baldy Mountain. Contains much pyroxene, a few serpentinized olivines, iron ore, and apatite in a colorless base of analcite. Analysis by Hillebrand.

P. Monchiquite, dike on Bandbox Mountain. Contains olivine, augite, biotite, analcite, and apatite, with traces of serpentine and chlorite. Analysis by Stokes.

Q. Shonkinite, Yogo Peak. Described in Amer. Journ. Sci., 3d series, vol. 50, p. 474. Chiefly augite and orthoclase, with a considerable amount of accessory biotite, iron ore, and andesine, less apatite and olivine, and a trace of kaolin. Analysis by Hillebrand.

	M.	N.	O.	P.	Q.
SiO ₂	52.26	54.42	48.35	48.39	48.98
Al ₂ O ₃	13.96	14.28	13.27	11.64	12.29
Fe ₂ O ₃	2.76	3.32	4.38	4.09	2.88
FeO.....	4.45	4.13	3.23	3.57	5.77
MgO.....	8.21	6.12	8.36	12.55	9.19
CaO.....	7.06	7.72	9.94	7.64	9.65
Na ₂ O.....	2.80	3.44	3.35	4.14	2.22
K ₂ O.....	3.87	4.22	3.01	3.24	4.96
H ₂ O at 110°.....	1.53	.22	.90	.28	.26
H ₂ O above 110°.....	1.34	.38	2.89	2.56	.56
TiO ₂58	.80	.52	.73	1.44
P ₂ O ₅52	.59	.40	.45	.98
Cr ₂ O ₃	trace	trace	trace	.07	trace
NiO.....			.04		
MnO.....	.14	.10	.19	trace	.08
SrO.....	.05	.13	.09	.15	.08
BaO.....	.23	.32	.54	.32	.43
Li ₂ O.....	trace	trace	trace	trace	trace
SO ₃08	
Cl.....				trace	
CO ₂49		.30		
F.....			.25		.22
	100.25	100.19	100.01	99.90	99.99

5. CASTLE MOUNTAIN DISTRICT.

Rocks described by Weed and Pirsson in Bull. 139. Analyses made by Pirsson in the laboratory of the Sheffield Scientific School.

A. Rhyolite, between Fourmile and Fivemile creeks, near Smith River. Contains quartz, soda orthoclase, tourmaline, and a little iron ore and zircon. P. R. C. 560.

B. Quartz-tourmaline-porphyry, upper Fourmile Creek. Contains quartz, orthoclase, plagioclase, tourmaline, fluorite, and a little white mica, apatite, and zircon. P. R. C. 558.

C. Granite, Elk Peak. Contains quartz, orthoclase, oligoclase, biotite, hornblende, iron ore, apatite, sphene, and zircon. The last three minerals are rare. P. R. C. 551.

D. Quartz-porphyry, sheet on ridge between Fourmile and Checkerboard creeks. Contains quartz, plagioclase, an untwinned feldspar and biotite, with occasional grains of iron ore and crystals of apatite and zircon. P. R. C. 557.

E. Rhyolite-pitchstone, forks of Checkerboard Creek. Essentially glass, inclosing grains of iron ore. P. R. C. 561.

F. Aplitic granite, dike between Blackhawk and Robinson. Contains quartz and unstriated feldspar, some plagioclase, and occasional biotite. P. R. C. 553.

G. Quartz-porphyry, Musselshell Canyon. Contains quartz, biotite, plagioclase, apatite, iron ore, and zircon. P. R. C. 556.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	74.90	74.82	72.48	72.38	72.56	72.88	71.67
Al ₂ O ₃	13.64	13.80	13.14	14.71	12.33	12.90	15.82
Fe ₂ O ₃66	.37	1.66	1.09	.80	.74	1.18
FeO.....	.50	.30	1.02	.82	.82	1.05	.35
MgO.....	trace	.10	.15	.70	trace	.75	.13
CaO.....	.61	.17	1.04	.67	trace	.81	.25
Na ₂ O.....	4.22	4.33	4.22	4.28	5.36	3.72	4.46
K ₂ O.....	4.64	4.81	4.88	4.15	3.08	5.03	4.45
H ₂ O.....	.33	.83	.42	.92	4.59	1.22	1.21
TiO ₂15	.25	.32	.10	.20	.45	.10
MnO.....	trace	trace	trace	trace	trace	.05	trace
Li ₂ O.....	trace	trace	trace	trace	-----	-----	-----
	99.65	99.78	99.33	99.82	99.74	99.60	99.62
Sp. gr.....	2.61	2.59	2.62	2.61	2.37	2.64	2.60

H. Feldspar-porphry, dike below Castle. Contains orthoclase, less plagioclase, hornblende, biotite partly altered to chlorite, sometimes allanite, and occasional iron ore, apatite, and zircon. P. R. C. 555.

I. Rhyolitic tuff, near forks of Checkerboard Creek. Contains fragments of shale, quartz, plagioclase, sanidine, and hornblende, in a brownish material carrying puniceous glass. Much decomposed, with formation of kaolin. P. R. C. 562.

J. Syenitic mass included in granite, head of Cottonwood Creek. Contains orthoclase, plagioclase, quartz, hornblende, biotite, apatite, and iron ore. A hornblende-mica-syenite. P. R. C. 554.

K. Diorite, between Blackhawk and Robinson. Contains plagioclase, orthoclase, quartz, biotite, diallage, hypersthene, iron ore, apatite, and zircon. P. R. C. 559.

L. Basalt, Volcano Butte. Contains labradorite, augite, olivine, a little serpentine, occasional quartz grains, small patches of glass, iron ore, chiefly ilmenite, apatite, a trace of calcite, and a mineral which may be nephelite or analcite. P. R. C. 565.

M. Augite-vogesite, dike on west side of upper Fourmile Creek. Contains augite, hornblende, iron ore, a little plagioclase, orthoclase, calcite, and some decomposition products. P. R. C. 563.

N. Monchiquite-like dike rock, west side of upper Willow Creek. Contains augite, olivine, biotite, ilmenite, and a colorless base which appears to be partially zeolitized glass. Also a little serpentine, chlorite, and calcite. P. R. C. 564.

	H.	I.	J.	K.	L.	M.	N.
SiO ₂	65.87	61.21	61.87	56.80	46.52	45.15	42.46
Al ₂ O ₃	16.82	15.67	17.26	18.30	10.48	15.39	12.04
Fe ₂ O ₃	1.58	4.06	2.35	1.64	4.40	2.76	3.19
FeO.....	1.23	.62	2.43	5.58	7.79	5.64	5.34
MgO.....	1.54	1.58	1.82	3.63	10.58	6.38	12.40
CaO.....	2.65	2.18	3.23	5.31	9.49	8.83	12.14
Na ₂ O.....	4.72	1.57	5.18	4.35	3.12	2.67	1.21
K ₂ O.....	3.15	2.75	3.83	3.28	1.55	2.77	2.68
H ₂ O.....	1.43	10.20	1.07	.53	1.79	2.85	4.03
TiO ₂37	.56	.87	.46	2.98	2.80	2.47
P ₂ O ₅	trace	-----	trace	trace	.83	.56	.84
X ^a	-----	-----	-----	-----	.73	-----	-----
MnO.....	trace	.10	.03	trace	.11	.14	.16
Li ₂ O.....	-----	trace	trace	trace	trace	trace	trace
CO ₂	-----	-----	-----	-----	trace	4.27	.55
	99.36	100.50	99.94	99.88	100.37	100.21	99.51
Sp. gr.....	2.62	-----	2.67	2.83	2.99	2.70	2.94

^a Unseparated rare earths or metallic acids, such as columbic, tantallic, etc.

6. HIGHWOOD MOUNTAINS.

Rocks collected by W. H. Weed and L. V. Pirsson, who furnish the petrographic data. Analyses made in the laboratory of the Sheffield Scientific School, under the direction of Professor Pirsson.

A. Trachy-andesitic breccia, head of north fork of Willow Creek. Contains hornblende, biotite, iron ore, apatite, plagioclase, and orthoclase, with a little kaolin and limonite. E. B. Hurlbut and B. Barnes, analysts.

B. Trachyte, dike 1 mile north of divide in Highwood Gap. Contains abundant phenocrysts of orthoclase and augite in a groundmass of the same minerals; also a little biotite, apatite, and iron ore, and some secondary kaolin. E. B. Hurlbut, analyst.

C. Trachyte, dike north end of South Mountain. Contains phenocrysts of alkali feldspar, ægirine-augite, and melanite, in a groundmass of alkali feldspar. Apatite, iron ore, a trace of calcite, and some kaolin are also present. H. W. Foote, analyst.

D. Trachyte, dike on Aspen Creek. Contains alkali-hornblende and alkali-feldspar phenocrysts, in a groundmass of alkali feldspars; also some apatite and iron ore. Rock stained by limonite. H. W. Foote, analyst.

E. Trachyte, dike at head of Shonkin Creek. Contains augite, biotite, iron ore, and orthoclase, in a groundmass of alkali feldspar; also apatite and some kaolin. W. M. Bradley, analyst.

	A.	B.	C.	D.	E.
SiO ₂	59.24	58.04	57.18	55.23	51.94
Al ₂ O ₃	13.84	17.24	18.54	18.31	15.78
Fe ₂ O ₃	5.46	2.49	3.65	4.90	4.07
FeO.....	1.36	1.24	1.15	2.06	3.17
MgO.....	4.79	1.79	.69	1.85	3.48
CaO.....	5.60	3.50	2.31	3.62	6.04
Na ₂ O.....	3.13	3.37	4.48	4.02	3.44
K ₂ O.....	4.22	10.06	8.58	6.43	7.69
H ₂ O.....	2.02	1.95	2.10	1.84	2.17
TiO ₂22	.30	.30	.42	.39
P ₂ O ₅34	.22	.05	.58	.59
MnO.....	trace	trace	trace	trace	trace
SrO.....	none	undet	trace	trace	.28
BaO.....	trace	undet	.49	.46	.42
SO ₃08	trace	.06	.23	.29
CO ₂	none	-----	-----	none	-----
Cl.....	.04	.38	.77	.32	.08
Less O.....	100.34	100.58	100.35	100.27	99.83
	.01	.09	.17	.08	.02
	100.33	100.49	100.18	100.19	99.81

F. Syenite, top of Palisade Butte. Contains augite, iron ore, apatite, alkali feldspar, a little biotite, some limonite in cracks, much natrolite, and possibly other zeolites. H. W. Foote, analyst.

G. Leucite-syenite, head of Davis Creek. Contains augite, iron ore, olivine, biotite, apatite, alkali feldspar, leucite, and some zeolites, with traces of limonite and serpentine. E. B. Hurlbut, analyst.

H. Monchiquite, dike east side of Highwood Gap. Contains augite, olivine, biotite, iron ore, apatite, and analcite, with some serpentine and a little kaolin. H. W. Foote, analyst.

I. Leucite-basalt, saddle between Highwood and Pinewood peaks. Contains augite, iron ore, leucite, apatite, altered olivine, chlorite, calcite, and zeolites. H. W. Foote, analyst.

J. Leucite-basalt or leucite-monchiquite, dike on Arrow Peak. Phenocrysts of augite, olivine, and biotite, with some leucite, in a groundmass of the same minerals. Also analcite, iron ore, and apatite. H. W. Foote, analyst.

	F.	G.	H.	I.	J.
SiO ₂	50.11	49.59	47.82	47.98	46.04
Al ₂ O ₃	17.13	14.51	13.56	13.34	12.23
Fe ₂ O ₃	3.73	3.51	4.73	4.09	3.86
FeO.....	3.28	5.53	4.54	4.24	4.60
MgO.....	2.47	6.17	7.49	7.01	10.38
CaO.....	5.09	9.04	8.91	9.32	8.97
Na ₂ O.....	3.72	3.52	4.37	3.51	2.42
K ₂ O.....	7.47	5.60	3.23	5.00	5.77
H ₂ O.....	4.47	1.95	3.37	2.10	2.87
TiO ₂82	.36	.67	.58	.64
P ₂ O ₅67	.15	1.10	1.03	1.14
MnO.....	trace	trace	trace	trace	trace
SrO.....	.35	.21	.21	.14	.25
BaO.....	.63	.49	.16	.50	.48
SO ₃08	.02	trace	trace	trace
CO ₂				1.24	
Cl.....	.07	.13	.04	.21	.11
	100.09	100.78	100.20	100.29	99.76
Less O.....	.02	.03	.01	.07	.03
	100.07	100.75	100.19	100.22	99.73

K. Missouriite, head of Shonkin Creek. Described in Amer. Journ. Sci., 4th series, vol. 2, p. 315. Contains apatite, iron ore, olivine, biotite, augite, and leucite, the two latter being the chief minerals. Some zeolites and analcite are also present. E. B. Hurlbut, analyst.

L. Pseudo leucite-syenite, head of Shonkin Creek. Contains augite, olivine, biotite, iron ore, apatite, orthoclase, nephelite, zeolites, and traces of serpentine, limonite, and kaolin. Contains pseudomorphs after leucite. E. B. Hurlbut, analyst.

M. Monzonite, Highwood Peak. Contains augite, biotite, iron ore, apatite, plagioclase, and alkali feldspar. E. B. Hurlbut, analyst.

N. Monzonite, Middle Peak. Contains augite, olivine, biotite, iron ore, apatite, plagioclase, and orthoclase. The two latter, with augite, are the chief minerals. E. B. Hurlbut, analyst.

	K.	L.	M.	N.
SiO ₂	46.06	51.75	51.00	52.05
Al ₂ O ₃	10.01	14.52	17.21	15.02
Fe ₂ O ₃	3.17	5.08	2.41	2.65
FeO.....	5.61	3.58	4.23	5.52
MgO.....	14.74	4.55	6.19	5.39
CaO.....	10.55	7.04	9.15	8.14
Na ₂ O.....	1.31	2.93	2.88	3.17
K ₂ O.....	5.14	7.61	4.93	6.10
H ₂ O.....	1.44	2.25	.63	.35
TiO ₂73	.23	.13	.47
P ₂ O ₅21	.18	.33	.21
MnO.....	trace	trace	trace	trace
SrO.....	.20	.07	.14	.28
BaO.....	.32	.30	.34	.42
SO ₃05	trace	.03	.02
Cl.....	.03	.05	trace	.24
	99.57	100.14	99.60	100.03
Less O.....	.01	.01	-----	.06
	99.56	100.13	-----	99.97

The following rock and separations, from Square Butte, at the east end of the Highwood Mountains, are described by Lindgren in Amer. Journ. Sci., 3d series, vol. 45, p. 286. Analyses by W. H. Melville, record No. 1268.

A. Post-Cretaceous sodalite-syenite. Contains orthoclase, some albite, hornblende, sodalite, analcite, and apatite. Orthoclase predominates. The sodalite amounts to 8 per cent. P. R. C. 184.

B. Hornblende separated from A. Near barkevikite.

C. Sodalite from A. Sp. gr. 2.265.

D. Analcite from A. Sp. gr. 2.255.

In addition, the orthoclase gave 3.88 per cent Na_2O and 11.03 per cent K_2O . A separation of mixed feldspars, sp. gr. 2.56, gave 6.08 per cent Na_2O and 8.91 per cent K_2O .

	A.	B.	C.	D.
SiO_2	56.45	38.41	41.56	49.54
Al_2O_3	20.08	17.65	29.48	25.07
Fe_2O_3	1.31	3.75		
FeO	4.39	21.75	.49	.40
MgO63	2.54	.15	.20
CaO	2.14	10.52	.49	.22
Na_2O	5.61	2.95	19.21	15.32
K_2O	7.13	1.95	.91	.89
H_2O at 100°26		.45	undet.
H_2O above 100°	1.51	.24	3.73	undet.
TiO_229			
P_2O_513			
NiO	trace	trace		
MnO09	.15		
Cl43		4.79	1.67
	100.45	99.91	101.26	93.31
Less O.....	.10		1.08	.38
	100.35		100.18	92.93

7. LITTLE ROCKY MOUNTAINS.

Granite-syenite-porphry. Described by Weed and Pirsson, Journal of Geology, vol. 4, p. 399. Contains orthoclase, quartz, oligoclase, iron oxides, and a little muscovite. Analysis by H. N. Stokes, record No. 1558. P. R. C. 905.

SiO_2	68.65	P_2O_5	trace
Al_2O_3	18.31	MnO	trace
Fe_2O_356	SrO10
FeO08	BaO13
MgO12	Li_2O	trace
CaO	1.00	SO_3	trace
Na_2O	4.86	Cl03
K_2O	4.74	F	trace
H_2O at 110°27		99.88
H_2O above 110°83		
TiO_220		

S. BEARPAW MOUNTAINS.

Described by Weed and Pirsson, Amer. Journ. Sci., 4th series, vol. 1, pp. 283 and 351, and vol. 2, pp. 136 and 188. Analyses by H. N. Stokes, record Nos. 1558 and 1572.

A. Quartz-syenite-porphry, Gray Butte. Contains anorthoclase, microlites of plagioclase, ægirite, augite, quartz, and apatite, with an occasional zircon and very few biotite leaves. P. R. C., 897.

B. Quartz-syenite, Beaver Creek stock. Contains orthoclase, albite, quartz, augite, and iron oxides, with very little biotite, hornblende, and sphene. P. R. C., 900.

C. Basic syenite or monzonite (yogoite), Beaver Creek. Contains orthoclase, plagioclase, diopside, biotite, iron oxides, and apatite. P. R. C., 902.

	A.	B.	C.
SiO ₂	66.22	68.34	52.81
Al ₂ O ₃	16.22	15.32	15.66
Fe ₂ O ₃	1.98	1.90	3.06
FeO.....	.16	.84	4.76
MgO.....	.77	.54	4.99
CaO.....	1.32	.92	7.57
Na ₂ O.....	6.49	5.45	3.60
K ₂ O.....	5.76	5.62	4.84
H ₂ O at 110°.....	.08	.15	.16
H ₂ O above 110°.....	.24	.30	.93
TiO ₂22	.21	.71
P ₂ O ₅10	.13	.75
MnO.....	trace	.07	trace
SrO.....	.06	.04	.09
BaO.....	.29	.08	.24
Li ₂ O.....	trace	none	trace
SO ₃02	trace	trace
Cl.....	.04	.04	.07
F.....	trace	none?	trace
	99.97	99.95	100.24

D. Shonkinite, Beaver Creek. Contains anorthoclase, diopside, biotite, iron oxides, and apatite, with very little olivine and probably a trace of nephelite. P. R. C. 901.

E. Leucitite, Bearpaw Mountains. An olivine-free leucite-basalt. Contains leucite, augite, iron oxides, rarely biotite, and very little glassy base. P. R. C. 903.

F. Tinguaita, dike on Bear Creek. Contains orthoclase, nephelite, cancrinite, augite, ægirite, apatite, a little sodalite, and a doubtful fibrous hornblende. P. R. C. 899.

G. Pseudo leucite-sodalite-tinguaita, Beaver Creek. Contains orthoclase, nephelite, sodalite, nosean, ægirite, diopside, and fluorite. P. R. C. 904.

	D.	E.	F.	G.
SiO ₂	50.00	46.51	57.46	51.93
Al ₂ O ₃	9.87	11.86	15.40	20.29
Fe ₂ O ₃	3.46	7.59	4.87	3.59
FeO.....	5.01	4.39	.87	1.20
MgO.....	11.92	4.73	1.37	.22
CaO.....	8.31	7.41	2.59	1.65
Na ₂ O.....	2.41	2.39	5.48	8.49
K ₂ O.....	5.02	8.71	9.44	9.81
H ₂ O at 110°.....	.17	1.10	.09	.10
H ₂ O above 110°.....	1.16	2.45	.82	.99
TiO ₂73	.83	.60	.20
P ₂ O ₅81	.80	.21	.06
Cr ₂ O ₃11	none	-----	-----
NiO.....	.07	.04	-----	-----
CoO.....	-----	trace	-----	-----
MnO.....	trace	.22	trace	trace
CuO.....	-----	trace	-----	-----
SrO.....	.07	.16	.16	.07
BaO.....	.32	.50	.60	.09
Li ₂ O.....	trace	trace	trace	trace
SO ₃02	.05	.13	.67
CO ₂31	none	.13	.25
Cl.....	.08	.04	.20	.70
F.....	.16	trace-	trace	.27
Less O.....	100.01 .08	99.78 .02	100.42 .05	100.58 .27
	99.93	99.76	100.37	100.31

IDAHO.

Rocks collected by Waldemar Lindgren, who furnishes the petrographic data. Those from Silver City and Hailey will be described in 20th Ann., Part III.

A. Quartz-monzonite, Idaho-Democrat mine, Hailey. Contains quartz, orthoclase, microcline, oligoclase, biotite, apatite, titanite, and magnetite. Sp. gr. 2.672, 27.5°.

B. Same rock and locality as A, but in altered condition. Contains quartz, sericite, chlorite, calcite, pyrite, rutile, etc. Sp. gr. 2.472, 29°.

C. Gabbro, related to diorite, Ceresus mine, Hailey. Contains labradorite, biotite, diallage, hypersthene, hornblende, quartz, titanite, magnetite, orthoclase, and chlorite. Sp. gr. 2.826, 28°.

D. Same rock and locality as C, but in altered condition. Contains quartz, sericite, chlorite, calcite, pyrite, arsenopyrite, rutile, etc. Sp. gr. 2.898, 28°.

Analyses A to D by W. F. Hillebrand, record No. 1826. The metals which are bracketed with sulphur represent sulphides. Traces of lithia are present in all four.

	A.	B.	C.	D.
SiO ₂	68.42	71.93	57.78	58.01
Al ₂ O ₃	15.01	12.21	16.28	15.72
Fe ₂ O ₃97	.64	1.02	.64
FeO.....	1.93	2.99	4.92	3.87
MgO.....	1.21	.58	4.60	2.07
CaO.....	2.60	2.59	6.65	2.15
Na ₂ O.....	3.23	.23	3.25	.10
K ₂ O.....	4.25	3.29	2.22	4.79
H ₂ O at 105°.....	.54	.37	.34	.31
H ₂ O above 105°.....	.73	2.06	.92	2.71
TiO ₂50	.40	1.07	1.08
P ₂ O ₅13	.10	.30	.31
CoO, NiO.....	none	none	.02	none
MnO.....	.06	.18	.15	.17
SrO.....	.03	none	.07	none
BaO.....	.12	trace	.12	trace ?
CO ₂20	1.95	.15	2.86
S.....	.02	.18	.02	1.25
Fe.....		.13		1.52
Co, Ni.....		none		.12
Zn.....		.09		
Pb.....		trace		.86
Cu.....		none		.05
As.....				1.65
	99.95	99.92	99.88	100.24

E. Altered rhyolite, De Lamar mine, Silver City, contains quartz, sericite, pyrite, apatite, and rutile, with some undeterminable magnesian mineral. Analysis by H. N. Stokes, record No. 1731.

F. Altered rhyolite, De Lamar mine. Contains sericite, quartz, kaolinite, and pyrite. Sp. gr., 2.655, 23°.

G. Altered rhyolite, De Lamar mine. Contains quartz, sericite, kaolinite, and pyrite. Sp. gr., 2.576, 24°.

H. Diabasic basalt, Trade Dollar mine, Silver City. Contains labradorite, augite, chlorite, magnetite, secondary quartz, etc.

Analyses F, G, and H by Hillebrand, record No. 1826.

	E.	F.	G.	H.
SiO ₂	66.69	87.37	78.59	48.47
Al ₂ O ₃	15.40	7.44	12.13	16.07
Fe ₂ O ₃	1.84	.09	none	4.12
FeO	undet	.18	.09	7.47
MgO85	.12	.41	5.96
CaO09	.10	.16	4.84
Na ₂ O16	.14	.10	2.43
K ₂ O	3.50	1.79	2.55	1.41
H ₂ O at 105°83	.51	.82	2.30
H ₂ O above 105°	2.97	1.39	2.47	4.63
TiO ₂	2.11	.09	.12	1.51
P ₂ O ₅08	trace	trace	.44
CoO, NiO		none	none	trace
MnO	trace	trace?	trace?	.23
SrO		none	none	trace
BaO09	.02	.02	.03
Li ₂ O		trace	trace	trace
SO ₃11			
FeS ₂	3.99	1.00	2.61	.24
Cu				trace
	98.71	100.24	100.07	100.15

Heavy metals not looked for in E.

I. Quartz-monzonite, Schafer Butte, Boise County. Contains quartz, orthoclase, oligoclase, biotite, apatite, titanite, and magnetite. To be described in 20th Ann. Analysis by George Steiger, record No. 1802.

J. Granodiorite, Silver Wreath mine, Boise County. Contains orthoclase, oligoclase, apatite, biotite, titanite, and magnetite. Described in 18th Ann., Part III, p. 642. Sp. gr. 2.714, 23°.

K. Same as J, but in altered condition. Also described in 18th Ann. Contains sericite, quartz, titanite, apatite, and pyrite, with carbonates of calcium, magnesium, and iron. Sp. gr. 2.774, 23°.

Analyses J and K by George Steiger, record No. 1691.

	I.	J.	K.
SiO ₂	69.56	65.23	66.66
Al ₂ O ₃	15.29	16.94	14.26
Fe ₂ O ₃86	1.60	.67
FeO	2.06	1.91	2.41
MgO69	1.31	.95
CaO	2.81	3.85	3.37
Na ₂ O	3.97	3.57	none
K ₂ O	3.36	3.02	4.19
H ₂ O at 100°86	.18	.36
H ₂ O above 100°88	2.16
TiO ₂55	.66	.49
P ₂ O ₅16	.19	.17
MnO		trace	trace
BaO19	none
S		none	.95
CO ₂25	3.67
	100.17	99.78	100.31
Less O=S24
			100.07

COLORADO.

1. DENVER BASIN.

Rocks described by Cross in Mon. XXVII. Analyses A, B, D, and E by L. G. Eakins, C by W. F. Hillebrand. All but A were made in the Denver laboratory.

A. Dolerite, dike near Valmont. Contains augite, plagioclase, olivine, orthoclase, and biotite, with accessory magnetite and apatite. Record No. 1145. P. R. C. 534.

B. Augite separated from A. P. R. C. 105.

C. Basalt, Table Mountain, lower capping sheet. Contains plagioclase, orthoclase, augite, magnetite, and apatite, with olivine much serpentinized. Sp. gr. 2.83, 22.5°.

D. Basalt, earlier flow, south slope of North Table Mountain. Contains augite, olivine, plagioclase, probably orthoclase, magnetite, apatite, and a little biotite. P. R. C. 535.

E. Augite-mica-syenite, from north fork of Turkey Creek, Jefferson County. Contains orthoclase, augite, biotite, rhombic pyroxene, hornblende, plagioclase, quartz, apatite, and magnetite. Sp. gr. 2.857, 29.5°. P. R. C. 532.

	A.	B.	C.	D.	E.
SiO ₂	48.25	49.10	52.59	49.69	56.90
Al ₂ O ₃	16.73	7.95	17.91	18.06	18.50
Fe ₂ O ₃	3.99	-----	3.81	2.64	.17
FeO.....	6.28	8.30	5.18	6.19	4.61
MgO.....	5.77	12.37	4.11	5.73	5.10
CaO.....	8.32	22.54	7.24	8.24	6.17
Na ₂ O.....	3.24	trace	2.94	2.99	2.99
K ₂ O.....	4.08	trace	3.83	3.90	4.14
H ₂ O.....	1.72	-----	1.24	.91	.51
TiO ₂89	-----	.84	.85	.19
P ₂ O ₅68	-----	.14	.81	.79
MnO.....	trace	-----	trace	.13	trace
BaO.....	.01	-----	-----	-----	-----
Cl.....	.08	-----	.05	.13	trace
SO ₃12	-----	-----	-----	-----
	100.16	100.26	99.88	100.27	100.07

The following rocks from the Denver Basin were analyzed by L. G. Eakins in the Denver laboratory, but the analyses do not appear in the monograph. The subjoined data have been supplied by Whitman Cross.

A. Enstatite-diabase-porphry, Mount Morrison. Contains labradorite and enstatite in a groundmass of considerable amount, which is colorless and cryptocrystalline, probably feldspathic, and carries magnetite and other indistinct ferritic matter. P. R. C. 533.

B. Augite-andesite, Table Mountain. Contains plagioclase (andesine), with rare augite and biotite, in a groundmass of plagioclase, augite, magnetite, and minor accessories. This rock contained ptilolite in its vesicular equivalent. Described in Proc. Colorado Sci. Soc., 1886, p. 72.

C. Augite-andesite, a pebble from the Denver beds, Table Mountain. A few augite and andesite phenocrysts in a groundmass of oligoclase, orthoclase, and quartz (?) grains, with some augite and magnetite.

	A.	B.	C.
SiO ₂	56.74	59.26	59.29
Al ₂ O ₃	18.80	23.63	21.27
Fe ₂ O ₃15	.30	3.33
FeO.....	6.91	.57	1.04
MgO.....	5.57	.31	1.12
CaO.....	7.34	5.93	5.25
Na ₂ O.....	2.32	4.94	3.39
K ₂ O.....	.77	4.78	3.00
H ₂ O.....	1.09	.74	1.63
P ₂ O ₅2023
MnO.....	.0721
	99.96	100.46	99.76
Sp. gr.....	2.876, 27°	2.625, 31°	2.596, 14°.5

2. PIKES PEAK DISTRICT.

FIRST GROUP. GRANITE.

Described by E. B. Mathews. Analyses by W. F. Hillebrand, record No. 1470.

A. Granite, Sentinel Point, western part of Pikes Peak massif. Contains microcline, microcline-perthite, quartz, biotite, a little oligoclase, and accessory fluorite, apatite, zircon, sphene, magnetite, and allanite. P. R. C. 600.

B. Granite, near road between Florissant and the Platte River. Consists chiefly of microcline in perthitic intergrowth with albite, quartz, and biotite. P. R. C. 606.

C. Porphyritic granite, south side of Pikes Peak, ridge between Middle and North Beaver creeks. Contains microcline, perthite, orthoclase, oligoclase, quartz, biotite, and accessory apatite, fluorite, zircon, and magnetite. P. R. C. 602.

D. Sheared granite, Currant Creek Canyon, north of Twelvemile Park. Contains perthitic microcline, quartz, muscovite, and sericitic aggregates replacing plagioclase and a part of the microcline. P. R. C. 604.

E. Granite, Currant Creek Canyon, north of Twelvemile Park. Consists chiefly of perthitic microcline, quartz, greenish biotite, muscovite, and plagioclase altered to a sericitic mass. Also flakes of limonite. Accessory minerals rare. P. R. C. 603.

F. Granite, Middle Beaver Creek, south side of Pikes Peak. Contains microcline, orthoclase, perthitic albite, oligoclase, abundant quartz, biotite, and a little accessory magnetite, fluorite, and zircon. P. R. C. 601.

G. Granite-gneiss, north of Twin Creek. Contains microcline, orthoclase, quartz, biotite, abundant fluorite, and a little sphene and apatite. P. R. C. 605.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	77.03	75.92	75.17	74.40	73.90	73.51	66.90
Al ₂ O ₃	12.00	12.96	12.66	14.43	13.65	13.28	14.86
Fe ₂ O ₃76	.33	.23	.89	.28	.94	.93
FeO.....	.86	1.40	1.40	.22	.42	.97	3.41
MgO.....	.04	trace	.05	.07	.14	.05	.31
CaO.....	.80	.15	.82	.58	.23	1.11	1.23
Na ₂ O.....	3.21	4.60	2.88	1.76	2.53	3.79	5.56
K ₂ O.....	4.92	4.15	5.75	6.56	7.99	5.22	5.02
H ₂ O at 110°.....	.14	.16	.16	.15	.16	.16	.16
H ₂ O above 110°.....	.30	.32	.66	.92	.33	.62	.31
TiO ₂13	.05	.10	.12	.07	.18	.43
P ₂ O ₅	trace	trace	.03	.22	.05	trace	.12
MnO.....	trace	.04	trace	trace	trace	trace	.15
SrO.....	none	none	trace?	none	none	none	none
BaO.....	trace	trace	.03	trace	trace	trace	.14
Li ₂ O.....	trace	trace	trace	trace	trace	trace	.06
F.....	.36	.12	.31	.04	none	.55	1.00
CO ₂03					
Less O.....	100.55	100.23	100.26	100.36	99.75	100.38	100.59
	.15	.05	.13	.02	-----	.22	.42
	100.40	100.18	100.13	100.34	-----	100.16	100.17

SECOND GROUP. ROCKS OF THE CRIPPLE CREEK DISTRICT.

Described by Cross. Descriptions published, except when otherwise stated, in 16th Ann., Part II, pp. 38-50. Analyses, with two exceptions, which are properly noted, by W. F. Hillebrand, record Nos. 1448, 1453, and 1524.

A. Phonolite, dike in granite northeast of Big Bull Mountain. Contains anorthoclase, nepheline, sodalite, ægirine-augite, some ægirine, biotite, magnetite, and limonite. Nosite not distinguishable. P. R. C. 616.

B. Phonolite, Mitre Peak. Contains sanidine, nepheline, sodalite, ægirine, nosite, and colorless particles which may be lāvenite. P. R. C. 608.

C. Phonolite, hill 1 mile south of Straub Mountain. Contains anorthoclase, sanidine, nepheline, sodalite, nosite, analcite, and ægirine, with sometimes ægirine-augite. P. R. C. 612.

D. Phonolite, Rhyolite Mountain. Contains nepheline, nosite, analcite, ægirine, and some ægirine-augite, in a feldspathic groundmass. Sp. gr. 2.52, 23°. P. R. C. 618.

E. Phonolite, between Florissant and Manitou. Analysis by L. G. Eakins, made in the Denver laboratory, inserted here for comparison with the Cripple Creek samples. Described by Cross in Proc. Colorado Sci. Soc., vol. 2, p. 167. Contains sanidine, nepheline, hornblende with accessory pyroxene (?), magnetite, apatite, and sphene. Sp. gr. 2.576, 13°. P. R. C. 637.

F. The portion of E soluble in hydrochloric acid. This portion amounted to 25.39 per cent, and is reduced here to 100. Analysis by Eakins.

	A.	B.	C.	D.	E.	F.
SiO ₂	59.00	58.98	58.78	58.64	60.02	44.66
Al ₂ O ₃	20.07	20.54	20.03	19.62	20.98	31.59
Fe ₂ O ₃	1.58	1.65	1.87	2.17	2.21	.95
FeO.....	.65	.48	.49	.42	.51
MgO.....	.10	.11	.16	.37	trace
CaO.....	1.05	.67	.83	1.24	1.18	2.25
Na ₂ O.....	8.34	9.95	9.36	8.39	8.83	18.42
K ₂ O.....	5.63	5.31	5.50	5.26	5.72	2.13
H ₂ O at 100°.....	.24	.19	.31	.34	} .70
H ₂ O above 100°.....	2.03	.97	1.57	2.40	
TiO ₂29	.24	.29	.20
P ₂ O ₅05	.04	.03	.03	trace
ZrO ₂20	.20	.17	.09
MnO.....	.12	.26	.15	.20	trace
SrO.....	none	none	none	trace
BaO.....	trace	none	none	trace
Li ₂ O.....	trace	trace	trace	trace
SO ₃07	.20	.12	trace?
Cl.....	.24	.28	.58	.14	trace
CO ₂2623
	99.92	100.07	100.24	99.74	100.15	100.00

G. Trachytic phonolite, dike on west slope of Bull Cliff. Contains alkali feldspars, nepheline, nosite, sodalite, augite, scanty ægirine, brown hornblende, magnetite, sphene, apatite, and sometimes lāvenite. P. R. C. 624.

H. Altered phonolite, Washington shaft, Victor. Contains potash feldspars, with some alteration to muscovite. Crystals of pyrite and fluorite are visible, but nepheline and ægirine have disappeared. Analysis hitherto unpublished. P. R. C. 623.

I. Nepheline-syenite, near the Longfellow mine. Contains alkali feldspars, nepheline, sodalite, augite, some ægirine, hornblende, biotite, sphene, apatite, and magnetite. Sp. gr. 2.68, 23°. Contains 0.02 V_2O_5 .

J. Nepheline-basalt, Appie Ellen shaft. Much altered. Rich in olivine, augite, and magnetite. Also contains nepheline, feldspars in small amount, and biotite. Sp. gr. 2.99, 23°.

K. Altered nepheline-basalt, Anna Lee mine. Analysis hitherto unpublished and not discussed.

L. Local facies of a phonolitic mass, Bull Cliff. Contains abundant augite, plagioclase, alkali feldspar, magnetite, and a little red-brown biotite, with a colorless isotropic base in the darker spots. Analysis hitherto unpublished. P. R. C. 627. Contains 0.03 V_2O_5 .

	G.	H.	I.	J.	K.	L.
SiO ₂	59.38	56.74	54.34	35.03	48.61	49.84
Al ₂ O ₃	19.47	20.30	19.23	9.80	20.74	17.78
Fe ₂ O ₃	1.60	1.06	3.19	5.55	4.29	5.86
FeO.....	1.19	-----	2.11	4.98	.22	2.62
MgO.....	.36	.23	1.28	9.78	2.11	3.02
CaO.....	1.96	.57	4.53	15.09	.25	7.35
Na ₂ O.....	7.80	.62	6.38	2.04	.16	5.20
K ₂ O.....	5.83	13.36	5.14	2.16	.77	3.04
H ₂ O at 110°....	.11	.33	.14	.41	12.10	.34
H ₂ O above 110°	.69	1.15	1.17	2.05	7.07	2.02
TiO ₂58	.58	1.09	2.20	3.57	1.43
P ₂ O ₅08	.25	.27	1.99	.29	.76
ZrO ₂10	.07	.07	none	-----	.03
Cr ₂ O ₃	-----	-----	-----	-----	trace	-----
MnO.....	.15	none	.08	.06	none	.21
SrO.....	.03	trace	.16	.17	none	.18
BaO.....	.13	.19	.24	.14	none	.22
Li ₂ O.....	trace	trace	trace	trace	(?)	trace
SO ₃37	-----	.07	none	-----	none
Cl.....	.22	-----	.28	trace	-----	trace
F.....	-----	undet.	undet.	undet.	.63	undet.
CO ₂	-----	-----	none	7.83	-----	.52
FeS ₂	-----	4.65	-----	.38	-----	-----
	100.05	100.10	99.77	99.66	100.81	100.42

THIRD GROUP. MISCELLANEOUS ROCKS.

Analyses, hitherto unpublished, except when otherwise stated, by W. F. Hillebrand, record Nos. 1448, 1453, and 1669. Petrographic data supplied by Whitman Cross. Names marked with a query are provisional designations only.

A. Trachyte?, near Robbins's ranch. Has scattered phenocrysts of alkali feldspar, oligoclase, and biotite in a predominant trachytic groundmass of alkali feldspar, tridymite, particles of residual glass, and ferritic flakes and grains.

B. Trachyte?, Wicher Mountain. Shows biotite and a few glassy feldspars in a groundmass resembling that of A.

C. Andesite?, Bare Hills. Composed mainly of plagioclase and alkali feldspar, with small augites and much ferritic matter. Minute prisms of a yellowish brown amphibole (?) and scales of tridymite are also visible.

D. Pyroxene-andesite?, Wicher Mountain. Contains phenocrysts of plagioclase, a few of augite, minute specks of iddingsite, and flakes of limonite. In the groundmass are plagioclase, orthoclase (?), augite, iddingsite, and hypersthene. In the pores tridymite appears. Contains 0.02 V₂O₃.

E. Plagioclase-basalt?, mesa east of Mac Gulch. Contains plagioclase, augite, olivine, iddingsite, magnetite, biotite, and apatite.

F. Plagioclase-basalt, Saddle Mountain. Phenocrysts of augite and olivine in a groundmass of plagioclase, orthoclase, augite, magnetite, biotite, and apatite. Very fresh. See Cross, Journ. Geol., Vol. 5, p. 684.

	A.	B.	C.	D.	E.	F.
SiO ₂	69.52	66.12	62.64	57.48	52.97	48.76
Al ₂ O ₃	15.44	17.21	17.82	18.04	18.31	15.89
Fe ₂ O ₃	1.90	2.43	3.91	5.73	1.86	6.04
FeO.....	.09	trace	.31	.73	6.73	4.56
MgO.....	.17	.35	.47	1.17	3.04	5.98
CaO.....	1.70	2.11	3.22	5.03	6.51	8.15
Na ₂ O.....	4.54	4.70	4.47	4.28	3.74	3.43
K ₂ O.....	5.04	5.57	4.99	4.15	3.35	2.93
H ₂ O at 100°...	.33	.14	.58	.62	.44	.40
H ₂ O above 100°	.27	.71	.65	.55	.31	1.48
TiO ₂23	.29	.59	1.00	1.04	1.65
P ₂ O ₅14	.11	.25	.66	.81	.60
ZrO ₂05	.06	.08	.04	.05	none
MnO.....	.08	.08	.04	trace	.09	.13
SrO.....	.04	.05	.07	.12	.14	.06
BaO.....	.19	.25	.28	.20	.18	.17
SO ₃16		
CO ₂17					
	99.90	100.18	100.37	99.96	99.57	100.23

G. Analcite-basalt, from the Basin. Contains phenocrysts of augite, olivine, and analcite. Also magnetite, with subordinate amounts of alkali feldspars, biotite, and apatite.

H. Portion of the analcite-basalt soluble in hydrochloric acid.

I. Augite separated from G.

J. Analcite separated from G. After deduction of 4.22 per cent of substance insoluble in boiling dilute hydrochloric acid, and later removal of liberated silica with weak potash solution. Sixty-two one hundredths per cent of the water goes off over sulphuric acid. A trace of lithia was found.

The analcite-basalt and its fractions are described by Cross in Journ. Geol., vol. 5, p. 684.

	G.	H.	I.	J.
SiO ₂	45.59	44.44	49.26	51.24
Al ₂ O ₃	12.98	<i>a</i> 20.11	6.01	24.00
Fe ₂ O ₃	4.97	} 7.50	3.31	} 1.20
FeO.....	4.70		4.23	
MgO.....	8.36	5.81	12.40	.33
CaO.....	11.09	3.94	21.79	1.62
Na ₂ O.....	4.53	8.17	.79	11.61
K ₂ O.....	1.04	1.13	.41	1.25
H ₂ O at 100°.....	.51	} 7.91	undet.	} 9.09
H ₂ O above 100°.....	3.40		undet.	
TiO ₂	1.32	-----	1.53	-----
P ₂ O ₅91	-----	-----	-----
ZrO ₂03	-----	-----	-----
MnO.....	.14	trace	undet.	-----
SrO.....	.12	.16	.06	.06
BaO.....	.13	(?)	(?)	-----
Cl.....	.05	.10	-----	trace
	99.87	99.27	99.79	100.40

a Includes P₂O₅ and possible ZrO₂ and TiO₂.

3. SILVER CLIFF AND ROSITA.

Rocks described by Cross, mostly in 17th Ann., Part II, p. 263. Also, partly, in Proc. Colorado Scientific Soc., vol. 2, p. 228. Analyses by L. G. Eakins. Those with record numbers attached were made in the Washington laboratory; all others in the laboratory at Denver.

A. Peridotite, Cottonwood Gulch. Contains hornblende, biotite, hypersthene, olivine, a little plagioclase, apatite, pyrrhotite, and sillimanite (?). P. R. C. 519.

B. Augite-diorite, Mount Fairview, Rosita Hills. Contains augite, biotite, labradorite, and accessory orthoclase, olivine, magnetite, and apatite. Sp. gr. 2.870, 32°. Olivine a subordinate constituent. P. R. C. 526.

C. The same as B, but with orthoclase in much larger amount. Sp. gr. 2.768, 34°. Record No., 1091. P. R. C. 529.

D. Trachyte, Game Ridge. Contains sanidine, plagioclase, a little biotite, magnetite, apatite, and zircon in a groundmass of orthoclase, with a little quartz. Sp. gr., 2.592, 29°. P. R. C. 524.

E. Trachyte, dark-colored dike. Contains more magnetite than D, but otherwise the two are practically identical. Sp. gr. 2.621, 24°.

	A.	B.	C.	D.	E.
SiO ₂	46.03	50.47	53.80	66.03	65.41
Al ₂ O ₃	9.27	18.73	20.13	18.49	18.78
Fe ₂ O ₃	2.72	4.19	3.57	2.18	.94
FeO.....	9.94	4.92	2.63	.22	.72
MgO.....	25.04	3.48	2.26	.39	.16
CaO.....	3.53	8.82	5.60	.96	1.58
Na ₂ O.....	1.48	4.62	5.20	5.22	5.91
K ₂ O.....	.87	3.56	4.49	5.86	5.41
H ₂ O.....	.64	.58	.90	.85	1.38
Ti ₂ O.....51	.43
P ₂ O ₅17	.10	.56	.04	trace
MnO.....	.40	.11	.29	trace	trace
CO ₂	trace	trace
Cl.....	trace
	100.09	100.09	99.86	100.24	100.29

F. Andesite, Pringle Hill. Contains plagioclase, orthoclase, quartz, biotite, augite, magnetite, and apatite, with sometimes a little hornblende. Sp. gr. 2.651, 17.8°. Record No. 1124. P. R. C. 528.

G. Pringle andesite, dike. Like F, but often contains sphene. Sp. gr. 2.690, 28°. P. R. C. 527.

H. Bunker andesite, Lookout Mountain. Contains plagioclase, orthoclase, augite, biotite, hornblende, quartz, magnetite, and apatite. Sp. gr. 2.699, 34°. Record No. 1091.

I. Altered Bunker andesite, Robinson Plateau. Biotite gone, magnetite altered to limonite, feldspars not much attacked. Record No. 1124. Sp. gr. 2.580, 19.7°.

J. Much-decomposed Bunker andesite, ridge near Knickerbocker Hill. Augite, hornblende, and biotite all replaced by decomposition products, plagioclase much muscovitized. Record No. 1124. P. R. C. 592.

	F.	G.	H.	I.	J.
SiO ₂	58.94	63.49	57.01	63.88	67.13
Al ₂ O ₃	17.19	18.40	18.41	19.96	18.41
Fe ₂ O ₃	2.63	2.44	3.69	2.21	.45
FeO	1.98	1.09	2.36	.57	.07
MgO	1.52	.66	2.34	.58	.44
CaO	4.45	2.30	4.29	2.03	.55
Na ₂ O	4.20	5.70	4.95	4.19	4.17
K ₂ O	3.90	4.62	3.72	3.88	5.28
H ₂ O	4.53	1.04	2.29	2.63	2.98
TiO ₂27	trace	.27	-----	.30
P ₂ O ₅23	trace	.42	-----	trace
MnO10	.16	.21	trace	trace
CO ₂	-----	trace	-----	-----	-----
	99.94	99.90	99.96	99.93	99.78

K. Dacite, Bald Mountain, near Rosita. Contains plagioclase, biotite, hornblende, sometimes augite, magnetite, and quartz. Sp. gr. 2.574, 24° . Record No. 1068. P. R. C. 530.

L. Mica-dacite. Contains plagioclase, sanidine, biotite, and quartz. Sp. gr. 2.563, 24° . Record No. 1068.

M. Rhyolite, Pennsylvania Hill. Sp. gr. 2.470, 26° . Record No. 1079. P. R. C. 525.

N. Rhyolite, Round Mountain. Contains sanidine, quartz, some glass, and accessory garnet. P. R. C. 521.

O. Rhyolite, Silver Cliff. Sp. gr. 2.560, 15° . Record No. 1125. P. R. C. 522.

	K.	L.	M.	N.	O.
SiO ₂	66.46	67.49	70.87	75.20	75.39
Al ₂ O ₃	17.91	17.76	15.18	12.96	13.65
Fe ₂ O ₃	2.42	2.54	2.18	.37	.38
FeO.....	.35	.08	.12	.27	.18
MgO.....	.49	.35	.60	.12	.15
CaO.....	2.89	1.67	1.58	.29	.51
Na ₂ O.....	4.79	5.03	3.47	2.02	1.84
K ₂ O.....	3.74	4.40	5.04	8.38	6.81
H ₂ O.....	1.01	.52	1.08	.58	1.13
TiO ₂	-----	-----	trace	-----	trace
P ₂ O ₅	-----	trace	trace	trace	trace
MnO.....	trace	trace	trace	.03	.14
	100.06	99.84	100.12	100.22	100.18

P. Rhyolitic tuff, a lake-bed deposit east of the Blue Mountains. Mainly composed of glassy rhyolite dust.

Q. Devitrified pitchstone, the gangue of ptilolite, 3 miles southeast of Silver Cliff. Shows feldspar, quartz, and barite. Described by Cross and Eakins in Amer. Journ. Sci., 3d series, vol. 44, p. 96. Record No. 1342. P. R. C. 590.

R Pitchstone, Rosita. Record No. 1033. P. R. C. 531.

S. Pitchstone, Fleetwood tunnel, Silver Cliff. P. R. C. 523.

T. Alteration product of S. Consists mainly of silica and kaolin. Record No. of S and T, 1034. P. R. C. 591.

	P.	Q.	R.	S.	T.
SiO ₂	71.02	65.67	73.11	71.56	71.71
Al ₂ O ₃	14.27	13.48	13.16	13.10	12.36
Fe ₂ O ₃	1.22	1.51	.62	.66	1.10
FeO.....			.23	.28	
MgO.....	trace	.31	.19	.14	1.21
CaO.....	1.38	2.41	.54	.74	1.11
Na ₂ O.....	2.28	1.52	2.85	3.77	.17
K ₂ O.....	3.97	2.42	5.10	4.06	.36
H ₂ O.....	6.12	12.27	4.05	5.52	11.97
P ₂ O ₅		trace			
MnO.....		trace	.14	.16	.17
BaO.....		.32			
SO ₃28			
	100.26	100.19	99.99	99.99	100.16

U. Syenite, Silver Cliff. Contains orthoclase and plagioclase in nearly equal amounts, colored by ferric hydroxide, with amphibole, a little biotite, and secondary epidote, calcite, and chlorite. Sp. gr. 2.689, 20°. P. R. C. 520.

V. Quartz-alunite rock, Democrat Hill. About two-thirds quartz and one-third alunite. Record No. 1126. P. R. C. 596.

W. Thesane, Mount Robinson. About one-fourth alunite. Record No. 1248. P. R. C. 593.

X. Quartz-diaspore rock, Mount Robinson. About 18 per cent diaspore, the rest quartz. Record No. 1167. P. R. C. 594.

Rocks V, W, and X are also described in Amer. Journ. Sci., 3d series, vol. 41, p. 471. These three rocks are decomposition products of rhyolite.

	U.	V.	W.	X.
SiO ₂	59.78	65.94	69.67	76.22
Al ₂ O ₃	16.86	12.95	13.72	19.45
Fe ₂ O ₃	3.08	.33	-----	trace
FeO.....	3.72	.07	-----	-----
MgO.....	.69	.05	trace	-----
CaO.....	2.96	.10	.07	trace
Na ₂ O.....	5.39	1.19	.34	trace
K ₂ O.....	5.01	2.32	2.44	trace
H ₂ O.....	1.58	4.47	4.73	3.82
TiO ₂	-----	-----	-----	.11
P ₂ O ₅	-----	-----	-----	.13
MnO.....	.14	trace	-----	-----
BaO.....	-----	trace	-----	-----
SO ₃	-----	12.47	9.27	.29
CO ₂75	-----	-----	-----
	99.96	99.89	100.24	100.02

The following analyses, all by L. G. Eakins, do not appear in the published memoir just cited. A and B were made in the Denver laboratory. The petrographic details have been supplied by Whitman Cross.

A. Spherulite in rhyolite, ridge west of Mount Tyndall. Mainly composed of quartz and feldspar needles, with some ferritic coloring due to decomposition of trichites.

B. Rhyolitic residual glass, same locality as A.

C. Interspherulitic mass, Rosita. Record No. 1285.

D. Spherulite, Rosita. Record No. 1286.

E, F, G. Spherulites, Silver Cliff. Record Nos. 1285, 1286.

These spherulites, C, D, E, F, and G, are made up of orthoclase needles, with free silica in fibers or grains, or rarely as tridymite.

The "soluble silica" is that which is dissolved by sodium carbonate solution.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	71.27	85.50	74.47	80.61	83.91	79.21	78.77
Al ₂ O ₃	16.02	7.42	13.87	10.94	9.54	12.24	12.46
Fe ₂ O ₃	1.41	1.23	-----	-----	-----	-----	-----
FeO17	.34	-----	-----	-----	-----	-----
MgO	trace	.82	trace	.09	trace	.11	.09
CaO35	.37	.51	.26	.19	.43	.34
Na ₂ O	5.76	.74	2.10	2.90	.62	2.58	2.12
K ₂ O	4.08	2.64	7.46	3.02	5.06	5.26	5.84
H ₂ O	1.14	1.22	1.88	2.20	.69	.66	.70
P ₂ O ₅	trace	none	-----	-----	-----	-----	-----
MnO	trace	.08	-----	-----	-----	-----	-----
	100.20	100.36	100.29	100.02	100.01	100.49	100.32
Soluble SiO ₂	-----	-----	12.72	11.12	1.06	1.27	1.25

4. BUFFALO PEAKS.

Hypersthene-andesite, described by Cross in Bull. 1. Contains hypersthene, augite, plagioclase, magnetite, and apatite. Sp. gr. 2.742, 16°. Analyses made by W. F. Hillebrand in the Denver laboratory.

A. Hypersthene-andesite. P. R. C. 588.

B, C, D. Hypersthene separated from the rock. Sp. gr. of D, 3.307, 23°. In B and C alkalis were not tested for. In C and D all the iron is given as FeO.

	A.	B.	C.	D.
SiO ₂	56.19	51.70	51.16	50.04
Al ₂ O ₃	16.12	1.72	2.15	2.91
Fe ₂ O ₃	4.92	.30
FeO.....	4.43	18.00	18.36	17.81
MgO.....	4.60	25.09	24.25	21.74
CaO.....	7.00	2.87	3.81	6.70
Na ₂ O.....	2.9627
K ₂ O.....	2.37
H ₂ O.....	1.03
P ₂ O ₅27
MnO.....	trace	.36	.36	.12
SrO.....	trace
BaO.....	trace
Cl.....	.02
	99.91	100.04	100.10	99.59

5. LEADVILLE REGION.

Rocks described by Cross in Mon. XII, Appendix A. Analyses made in the Denver laboratory.

A. Porphyry, Mount Zion. Contains orthoclase, plagioclase, quartz, biotite, apatite, magnetite, and zircon. Analysis by L. G. Eakins. P. R. C. 504.

B. White, or Leadville, porphyry. Contains orthoclase, plagioclase, quartz, muscovite, magnetite, apatite, and zircon, with crystals which appear to be rutile and anatase. Sp. gr. 2.680, 16°. Analysis by W. F. Hillebrand. P. R. C. 587.

C. Porphyry, summit of Mount Lincoln. Contains quartz, orthoclase, plagioclase, biotite, apatite, sphene, magnetite, zircon, and allanite. The sample analyzed showed some muscovite, chlorite, and calcite. Sp. gr. 2.670, 16°. Analysis by W. F. Hillebrand. P. R. C. 505.

	A.	B.	C.
SiO ₂	73.50	70.74	66.45
Al ₂ O ₃	14.87	14.68	15.84
Fe ₂ O ₃95	.69	2.59
FeO.....	.42	.58	1.43
MgO.....	.29	.28	1.21
CaO.....	2.14	4.12	2.90
Na ₂ O.....	3.46	2.29	3.92
K ₂ O.....	3.56	2.59	2.89
H ₂ O.....	.90	2.09	.84
TiO ₂10
P ₂ O ₅	none		.36
MnO.....	.03	.06	.09
SrO.....	trace	trace	.07
BaO.....		.03	
Li ₂ O.....			trace
CO ₂		2.14	1.35
Cl.....		trace	.05
	100.12	100.29	100.09

D. Gray porphyry, Johnson Gulch, near Leadville. Slightly altered. Contains orthoclase, plagioclase, biotite, and quartz, with decomposition products probably derived from original hornblende. Sp. gr. 2.736, 16° . Analysis by Hillebrand.

E. Pink orthoclase crystals from D. Analysis by Hillebrand.

F. Hornblendic porphyrite, lower Buckskin Gulch. Contains plagioclase, orthoclase, quartz, hornblende, biotite, magnetite, apatite, and zircon, with a little secondary calcite and chlorite. Sp. gr. 2.768, 16° . Analysis by Hillebrand. P. R. C. 506.

G. Biotite-porphyrite, dike in gneiss in the North Mosquito amphitheater. Composition like F, but with no hornblende, much biotite, and some pyrite. Sp. gr. 2.740, 16° . Analysis by Hillebrand. P. R. C. 507.

	D.	E.	F.	G.
SiO ₂	68.10	62.22	56.62	64.81
Al ₂ O ₃	14.97	20.33	16.74	15.73
Fe ₂ O ₃	2.78	4.94	1.68
FeO.....	1.10	3.27	2.91
MgO.....	1.10	4.08	2.82
CaO.....	3.04	2.95	7.39	4.22
Na ₂ O.....	3.46	3.45	3.50	3.98
K ₂ O.....	2.93	8.31	1.97	1.43
H ₂ O.....	1.28	1.90	.92	.62
TiO ₂0708
P ₂ O ₅16	trace	.23
MnO.....	.0915	.08
SrO.....	.08	trace	trace
CO ₂92	1.15	1.08
Cl.....	.0304
FeS ₂90
	100.11	99.16	100.73	100.61

H. Rhyolite (nevadite), from Chalk Mountain. Mainly quartz and feldspar, the latter being sanidine and plagioclase. A little biotite, magnetite, apatite, and zircon are present. Analysis by Hillebrand. P. R. C. 64.

I. Sanidine from H. Analysis by Hillebrand. P. R. C. 64.

	H.	I.
SiO ₂	74.45	65.04
Al ₂ O ₃	14.72	20.40
Fe ₂ O ₃	none
FeO.....	.56
MgO.....	.37
CaO.....	.83	.79
Na ₂ O.....	3.97	4.11
K ₂ O.....	4.53	9.74
H ₂ O.....	.66	.29
P ₂ O ₅01
MnO.....	.28
Li ₂ O.....	trace
	100.38	100.37

6. TENMILE DISTRICT.

Rocks described by Cross in 14th Ann., page 165. Analyses made in the Denver laboratory, A and C by W. F. Hillebrand, B by L. G. Eakins.

A. Quartz-hornblende-mica-porphyrity, Gold Hill. Contains plagioclase, hornblende, biotite, and quartz, in a groundmass of quartz, orthoclase, and a little plagioclase. P. R. C. 510.

B. Quartz-porphyrity, Sugar Loaf. Contains plagioclase, biotite, and quartz, in a groundmass of quartz and orthoclase. P. R. C. 509.

C. Quartz-porphyrity, Chicago Mountain. Contains plagioclase, orthoclase, biotite, and quartz. P. R. C. 508.

	A.	B.	C.
SiO ₂	63.66	67.29	68.30
Al ₂ O ₃	17.05	15.78	16.24
Fe ₂ O ₃	1.97	1.86	1.60
FeO.....	2.62	1.97	1.63
MgO.....	1.99	.72	1.05
CaO.....	3.89	2.36	2.79
Xa ₂ O.....	4.13	3.77	3.90
K ₂ O.....	3.09	3.55	3.52
H ₂ O at 110°.....	1.19	2.10	.71
H ₂ O above 110°.....			
TiO ₂	undet.	none	undet.
P ₂ O ₅27	.28	.13
MnO.....	.14	.21	.12
SrO.....	.08	none	.04
BaO.....			trace
Li ₂ O.....	none	trace	trace
CO ₂27	
	100.08	100.16	100.03

The following rocks were also analyzed in the Denver laboratory, but the analyses are hitherto unpublished. Petrographic data furnished by Whitman Cross.

A. Granite-prophyry, McNulty Gulch. Contains phenocrysts of orthoclase, oligoclase, andesine, quartz, biotite, and altered hornblende, in a groundmass of quartz, orthoclase, and magnetite. Accessory sphene, allanite, apatite, and zircon, and a little secondary chlorite are also present. Analysis by W. F. Hillebrand. P. R. C. 586.

B. Granite-porphyry, Jefferson tunnel. Contains orthoclase, oligoclase, quartz, and biotite, in a groundmass of mainly quartz and orthoclase; also accessory magnetite, apatite, zircon, and allanite. Chlorite appears as a decomposition product of biotite, and calcite and magnetite are present in small amounts. Analysis by Hillebrand.

C. Same as B. Analysis by L. G. Eakins. P. R. C. 583.

D. Diorite-porphyry, Copper Mountain. Contains oligoclase, hornblende, and biotite, in a groundmass of quartz, orthoclase, plagioclase, and magnetite; also accessory zircon, sphene, and apatite, and a little secondary chlorite and epidote. Analysis by Eakins. P. R. C. 585.

E. Diorite-porphyry, McNulty type. Contains oligoclase, andesine, hornblende, biotite, and magnetite, in a groundmass of orthoclase, plagioclase, quartz, magnetite, apatite, allanite, and sphene; also secondary chlorite, epidote, and calcite. Analysis by Eakins. P. R. C. 584.

	A.	B.	C.	D.	E.
SiO ₂	68.60	65.94	65.51	67.01	63.02
Al ₂ O ₃	16.21	16.00	17.01	18.03	17.61
Fe ₂ O ₃	1.67	.60	none	.66	1.78
FeO.....	1.57	1.74	2.79	.72	2.76
MgO.....	1.05	1.02	.90	.84	1.63
CaO.....	2.61	2.87	3.16	3.99	3.30
Na ₂ O.....	3.29	3.85	3.82	4.42	4.72
K ₂ O.....	3.88	4.56	4.67	3.53	3.23
H ₂ O.....	.92	1.13	1.78	.91	2.03
P ₂ O ₅21	.23	.13	.10	.16
MnO.....	.09	.1409	trace
SrO.....	trace	trace
BaO.....10	.08
CO ₂19	1.55
Cl.....	.03	.03	trace
S.....38
FeS ₂60
	100.32	100.26	100.15	100.40	100.32
Sp. gr.....	2.640, 27°	2.672, 21°	2.666, 26°	2.689, 16°5

7. ELK MOUNTAINS.

Analyses made by L. G. Eakins in the Denver laboratory, and hitherto unpublished. Petrographic data supplied by Whitman Cross.

A. Rhyolite, East Mountain, Crested Butte district. Phenocrysts of orthoclase, oligoclase, quartz, and biotite, in a microspherulitic and cryptocrystalline groundmass. P. R. C. 514.

B. Rhyolite, Round Mountain, Crested Butte district. Small phenocrysts of quartz, sanidine, biotite, and oligoclase, in a groundmass of quartz and orthoclase. P. R. C. 513.

C. Diorite, Brush Creek, Gunnison County. Contains several varieties of plagioclase, with orthoclase, quartz, hornblende, biotite, augite, sphene, apatite, and magnetite. P. R. C. 93.

	A.	B.	C.
SiO ₂	74.84	71.56	62.71
Al ₂ O ₃	14.05	14.91	17.06
Fe ₂ O ₃17	1.47	3.79
FeO.....	.31	1.04	2.74
MgO.....	trace	.08	1.78
CaO.....	1.57	1.98	5.51
Na ₂ O.....	3.66	3.78	3.54
K ₂ O.....	3.14	4.94	2.96
H ₂ O.....	2.33	.44	.24
P ₂ O ₅		trace	none
MnO.....			trace
	100.07	100.20	100.33
Sp. gr.....	2.38, 17°	2.59, 18°	2.791, 30°

8. WEST ELK MOUNTAINS.

Rocks described by Cross in 14th Ann., p. 165.

A. Hornblende-mica-porphyrity, Cliff Creek. Contains plagioclase, hornblende, and biotite, in a groundmass of quartz, feldspar, and mica. Analysis by W. F. Hillebrand, record No. 1429.

B. Porphyrite, Storm Ridge. Contains plagioclase, biotite, hypersthene, hornblende, and augite, in a groundmass of quartz and orthoclase. Analysis by L. G. Eakins, record No. 1238. P. R. C. 517.

C. Porphyrite-diorite, Mount Marcellina. Contains plagioclase, hornblende, and a little biotite, in a groundmass of quartz and orthoclase. Analysis by T. M. Chatard, record No. 1238. P. R. C. 516.

D. Quartz-porphyrity, Mount Carbon. Contains plagioclase, orthoclase, biotite, hornblende, augite, and quartz. Analysis by Chatard, record No. 1238. P. R. C. 518.

E. Quartz-porphyrity, Crested Butte. Contains plagioclase, orthoclase, hornblende, biotite, quartz, and a little augite. Analysis made by Eakins in the Denver laboratory. P. R. C. 515.

	A.	B.	C.	D.	E.
SiO ₂	63.05	61.42	62.85	65.36	65.71
Al ₂ O ₃	15.58	17.69	16.21	15.48	18.30
Fe ₂ O ₃	2.92	4.24	3.08	3.09	1.19
FeO	2.11	1.74	1.46	1.21	1.53
MgO	1.70	1.81	1.47	1.53	.98
CaO	4.15	5.29	4.72	4.14	2.17
Na ₂ O	3.77	3.14	3.49	3.58	5.00
K ₂ O	3.66	3.19	3.10	3.41	3.95
H ₂ O at 100°55	} .97	.29	.82	} 1.39
H ₂ O above 100°	1.38		2.03	.70	
TiO ₂60	.37	.41	.52	undet.
P ₂ O ₅27	.14	.48	.25	-----
MnO12	.19	.15	.19	.02
SrO07	-----	-----	-----	-----
BaO13	.09	.11	.08	-----
Li ₂ O	trace	-----	-----	-----	-----
	100.06	100.28	99.85	100.36	100.24

9. SAN JUAN REGION.

Rocks collected by Whitman Cross, who supplies the petrographic data. Hitherto unpublished.

A. Rhyolitic vitrophyre, near Del Norte, Rio Grande County. Reported by Cross as containing phenocrysts of oligoclase, quartz, biotite, and augite in a dark, fresh, glassy groundmass, the latter being predominant. Analysis made by Eakins in the Denver laboratory. Sp. gr. 2.423, 14°. P. R. C. 598.

B. Rhyolite, Summit district, Rio Grande County. Large phenocrysts of sanidine, with smaller ones of oligoclase and biotite, in a groundmass of orthoclase, quartz, oligoclase, biotite, and magnetite. Analysis by Eakins, made in the Denver laboratory. Sp. gr. 2.489 14°. P. R. C. 164.

C. Augite-diorite, Sultan Mountain, San Juan County. Contains plagioclase, orthoclase, quartz, augite, biotite, magnetite, and apatite. Some chlorite and epidote as alteration products. Analysis by L. G. Eakins in the Denver laboratory. Sp. gr. 2.751, 14°.

D. Augite-diorite, Stony Mountain, Ouray County. Analysis by Eakins in the Denver laboratory. Sp. gr. 2.891, 13.5°. P. R. C. 199.

	A.	B.	C.	D.
SiO ₂	68.61	68.85	63.91	52.05
Al ₂ O ₃	16.43	17.01	17.07	17.96
Fe ₂ O ₃73	1.78	4.39	4.09
FeO.....	1.52	.65	1.51	6.33
MgO.....	.05	trace	.81	5.03
CaO.....	1.79	1.62	4.47	8.64
Na ₂ O.....	2.82	3.44	3.48	2.99
K ₂ O.....	4.65	5.11	3.74	1.61
H ₂ O.....	3.35	1.79	.33	.97
P ₂ O ₅21	.31
MnO.....		trace		.43
	99.95	100.25	99.92	100.41

E. Diorite, La Plata Mountains. Contains augite, hornblende, plagioclase, and orthoclase in large amount, with biotite, quartz, sphene, apatite, and magnetite as subordinate constituents. Also secondary chlorite, muscovite, and calcite. Analysis by W. F. Hillebrand, record No. 1640. Sp. gr. 2.79, 21°.

F. Diorite-porphry, La Plata Mountains. Contains phenocrysts of hornblende, plagioclase, occasional quartz, sphene, apatite, and magnetite, in a groundmass of orthoclase, plagioclase, and quartz. Also secondary epidote, chlorite, and calcite. Analysis by Hillebrand, record No. 1636. Sp. gr. 2.677, 24°. A trace of sulphur is present.

G. Syenite, between Tirbircio and Schurman gulches, La Plata Mountains. Contains much alkali feldspar, some oligoclase, augite, biotite, and hornblende, with a little titanite, magnetite, and apatite. Analysis by H. N. Stokes, record No. 1764. Sp. gr. 2.704, 25°.

H. Monzonite, Babcock Peak, La Plata Mountains. Contains orthoclase and plagioclase in about equal amounts, with augite and hornblende, and a little quartz, titanite, magnetite, and apatite. Analysis by Stokes, record No. 1764. Sp. gr. 2.767, 26°.

I. Porphyritic lamprophyre, allied to camptonite, Snowstorm Peak, La Plata Mountains. Contains numerous phenocrysts of green hornblende, augite, and plagioclase, in a groundmass of plagioclase, orthoclase, augite, magnetite, and apatite. Some secondary calcite. Analysis by Hillebrand, record No. 1640. Sp. gr. 2.906, 21°.

	E.	F.	G.	H.	I.
SiO ₂	55.53	60.44	59.79	57.42	47.25
Al ₂ O ₃	16.78	16.65	17.25	18.48	15.14
Fe ₂ O ₃	4.06	2.31	3.60	3.74	5.05
FeO.....	3.35	3.09	1.59	2.10	4.95
MgO.....	3.00	2.18	1.24	1.71	6.87
CaO.....	6.96	4.22	3.77	6.84	9.98
Na ₂ O.....	4.31	5.18	5.04	4.52	2.39
K ₂ O.....	3.57	2.71	5.05	3.71	2.60
H ₂ O at 110°.....	.09	.36	.19	.08	.40
H ₂ O above 110°.....	.55	1.07	.39	.28	2.12
TiO ₂95	.60	.67	.86	1.22
P ₂ O ₅47	.29	.35	.36	.25
V ₂ O ₃02	.0205
NiO, CoO.....	trace	none02
MnO.....	.16	.13	.20	.09	.17
SrO.....	.11	.11	.11	.08	.05
BaO.....	.13	.12	.14	.15	.08
Li ₂ O.....	trace	trace	trace	trace
CO ₂09	.48	.72	none	1.24
SO ₃04	none
Cl.....	trace	.03
FeS ₂04	none
	100.17	99.96	100.14	100.45	100.46

J. Porphyritic lamprophyre, allied to camptonite, Indian Trail Ridge, La Plata quadrangle. Contains phenocrysts of green hornblende and colorless dioxide in a subordinate groundmass of plagioclase, orthoclase (?), augite, magnetite, and apatite. Much secondary calcite and some serpentine. Analysis by W. F. Hillebrand, record No. 1640. Sp. gr. 2.912, 19.5°.

K. Lamprophyre, allied to camptonite, Black Face, Telluride quadrangle. Consists of a fine felt of plagioclase, augite, and brown hornblende microlites, with flakes of biotite, and a cryptocrystalline part, which is probably in large degree orthoclase. Some magnetite and apatite. Analysis by Hillebrand, record No. 1719. Sp. gr. 2.783, 22°.

L. Quartz monzonite, northeast of San Miguel Peak, Telluride quadrangle. Contains orthoclase and plagioclase in about equal amounts, with abundant quartz and much less augite, hornblende, biotite, magnetite, and apatite. Analysis by H. N. Stokes, record No. 1764. Sp. gr. 2.720, 34°.

M. Gabbroitic facies of a diorite-monzonite stock, Ophir Needles, Telluride quadrangle. Contains abundant labradorite, with augite, hypersthene, biotite, orthoclase, magnetite, apatite, and a very little quartz. Analysis by Stokes, record No. 1764. Sp. gr. 2.860, 33°.

N. Gabbro-porphry, pass south of Mount Sneffels, Telluride quadrangle. Contains numerous phenocrysts of labradorite or bytownite, in a groundmass of plagioclase, orthoclase (?), augite, hypersthene, biotite, magnetite, and apatite. Analysis by Stokes, record No. 1764. Sp. gr. 2.949, 26.5°.

	J.	K.	L.	M.	N.
SiO ₂	43.98	55.65	65.70	56.93	47.32
Al ₂ O ₃	13.30	17.04	15.31	17.03	16.71
Fe ₂ O ₃	3.67	2.81	2.54	3.67	6.92
FeO.....	6.92	5.17	1.62	4.54	5.94
MgO.....	7.03	3.42	1.62	3.30	5.69
CaO.....	10.66	6.82	2.56	6.51	8.51
Na ₂ O.....	2.15	3.27	3.62	3.19	2.70
K ₂ O.....	1.64	2.29	4.62	2.58	2.02
H ₂ O at 110°.....	.42	.46	.17	.13	.24
H ₂ O above 110°.....	1.52	1.49	.42	.45	1.04
TiO ₂	1.18	.90	.72	1.03	1.50
P ₂ O ₅32	.37	.33	.44	.96
MnO.....	.22	.20	trace	.10	.08
NiO, CoO.....	.03	none			
SrO.....	.05	.05	.03	.06	.06
BaO.....	.06	.08	.12	.08	.07
Li ₂ O.....	trace	trace	trace	none	trace
CO ₂	6.46		none	none	none
SO ₃		none	.12	none	.19
Cl.....			.03	trace	trace
FeS ₂54				
	100.15	100.02	99.53	100.04	99.95

10. MISCELLANEOUS ROCKS.

A. Rhyolite, east bank of Arkansas River, Nathrop. Described by Cross in Proc. Colorado Sci. Soc., vol. 2, p. 69. Contains quartz and sanidine in a groundmass mainly of quartz and alkali feldspar. This rock carries topaz and spessartite in its lithophysae. Analysis made by L. G. Eakins in the Denver laboratory. Sp. gr. 2.602, 29°.

B. Granite, Platte Canyon. Described by E. B. Mathews in Bull. 150, p. 172. Contains microcline, quartz, biotite, oligoclase, and fluorite. Apatite, zircon, magnetite, hematite, limonite, epidote, and rutile (?) are sometimes present. Analysis by H. N. Stokes, record No. 1314.

C. Hornblende-porphyrity, Hermano Peak, Sierra El Late. Described by Cross in 14th Ann., p. 165. Contains plagioclase, hornblende, rare quartz, and a little biotite. Analysis by W. F. Hillebrand, record No. 1429.

D. Hornblende-porphyrity, Ute Peak, Sierra El Late. Described by Cross in 14th Ann., p. 165. Contains plagioclase, hornblende, and very little augite, in a groundmass of quartz, orthoclase, and plagioclase. Analysis by Hillebrand, No. 1429.

E. Porphyritic augite-diorite, Lone Cone, San Miguel Mountains. Described by Cross in 14th Ann., p. 165. Contains plagioclase, augite, hornblende, and biotite, in a groundmass of quartz, orthoclase, and plagioclase. Inclusions of magnetite and apatite in the augite. Analysis by Hillebrand, No. 1429.

	A.	B.	C.	D.	E.
SiO ₂	69.89	77.02	62.65	59.42	59.19
Al ₂ O ₃	17.94	11.63	16.68	16.79	18.00
Fe ₂ O ₃39	.32	2.35	3.23	3.07
FeO.....	.52	1.09	2.63	3.29	2.32
MgO.....	.14	.14	1.43	2.24	1.41
CaO.....	trace	1.24	4.96	5.57	6.55
Na ₂ O.....	4.21	2.85	4.45	4.15	4.01
K ₂ O.....	4.38	5.21	2.75	2.82	2.74
H ₂ O at 100°.....	2.07	.35	.27	.27	.46
H ₂ O above 100°.....			.66	.79	1.06
TiO ₂42	.68	.58
P ₂ O ₅	trace		.28	.35	.29
MnO.....	.23	trace	.16	.13	.19
SrO.....			.11	.07	.13
BaO.....			.13	.14	.18
Li ₂ O.....	trace		trace	trace	trace
CO ₂44	
	99.77	99.85	99.93	100.38	100.18

F. Tinguaité?, Two Buttes. Collected by G. K. Gilbert; petrographic data supplied by Whitman Cross. Consists chiefly of pale-green augite, hornblende, apatite, magnetite, and occasional crystals of alkali feldspar, in an obscure, largely isotropic groundmass. Sp. gr., 2.79, 25°.

G. Pyroxene from F. Sp. gr., 3.43, 28°.

H. Portion of F soluble in $\frac{1}{40}$ nitric acid.

I. Syenitic lamprophyre?, Two Buttes. Collected by Gilbert; description by Cross. Chief constituents, diopside, alkali feldspar, considerable biotite, magnetite, and olivine. The ferromagnesian minerals predominate. Sp. gr., 2.88, 29°. Also contains 0.03 V₂O₃.

J. Pyroxene from I. Sp. gr., 3.45, 25°.

K. Portion of I soluble in $\frac{1}{40}$ nitric acid.

Analyses F to K by W. F. Hillebrand, record No. 1604.

	F.	G.	H.	I.	J.	K.
SiO ₂	47.61	47.54	13.27	50.41	51.27	1.58
Al ₂ O ₃	14.26	4.14	9.40	12.30	3.05	1.00
Fe ₂ O ₃	4.90	5.64	-----	5.71	3.08	none
FeO.....	4.07	6.42	trace	3.06	4.34	.87
MgO.....	2.62	10.05	trace	8.69	14.21	1.22
CaO.....	8.71	21.57	1.70	7.08	22.58	.68
Na ₂ O.....	6.70	1.38	5.41	.97	.67	undet.
K ₂ O.....	4.08	.12	.66	7.53	.06	undet.
H ₂ O at 110°...	.26	none	undet.	.46	none	none
H ₂ O above 110°	1.89	undet.	undet.	1.80	undet.	undet.
TiO ₂	1.38	3.00	-----	1.47	.70	-----
P ₂ O ₅	1.38	-----	1.34	.46	-----	.46
ZrO ₂18	none	-----	none	none	-----
Cr ₂ O ₃	trace?	trace?	-----	trace	none	-----
NiO.....	trace	trace	-----	.04	.03	-----
MnO.....	.30	.36	trace	.15	.28	trace
SrO.....	.36	none	.11	.06	none	(?)
BaO.....	.41	none	none	.23	none	none
Li ₂ O.....	trace	trace	(?)	trace	-----	-----
SO ₂	1.17	-----	.77	none	-----	-----
S.....	.03	-----	.03	none	-----	-----
Cl.....	.37	-----	.37	trace	-----	trace
F.....	trace	-----	trace	trace?	-----	-----
	100.68	100.21	33.06	100.42	100.27	5.81

UTAH.

I. ROCKS FROM THE TINTIC DISTRICT.

Described by Tower and Smith in 19th Ann., Part III, pp. 609 et seq. Analyses by Stokes, record No. 1746.

A. Gray, porphyritic rhyolite, south of Pinyon Creek. Contains phenocrysts of sanidine, quartz, biotite, plagioclase, and hornblende; the last mineral sparingly. Also tridymite, magnetite, apatite, zircon, and a small amount of glassy residue.

B. Quartz-porphyry, Swansea mine. Phenocrysts of feldspar and quartz, the orthoclase somewhat altered. Microscopic biotite occurs sparingly. Also contains magnetite, apatite, zircon, a little chlorite, and secondary pyrite.

C. Andesite, Tintic Mountain. Phenocrysts of biotite and feldspar. Contains plagioclase, chiefly labradorite, augite, biotite, hypersthene, magnetite, and apatite, in a dark-gray glass.

D. Granular monzonite, Iron Duke mine. Contains orthoclase, plagioclase, quartz, hornblende, biotite, magnetite, apatite, zircon, and titanite, with a little chlorite and epidote.

E. Altered monzonite, near Tintic mine. Feldspar and ferromagnesian minerals completely altered. Rutile is present; quartz seems to have been added.

	A.	B.	C.	D.	E.
SiO ₂	69.18	71.56	60.17	59.76	71.14
Al ₂ O ₃	14.36	14.27	15.77	15.77	16.22
Fe ₂ O ₃	2.52	.89	3.42	3.77	.94
FeO.....	.57		2.95	3.30	.16
MgO.....	.70	.42	2.52	2.16	1.12
CaO.....	1.88	1.18	4.69	3.88	.25
Na ₂ O.....	3.58	3.00	2.96	3.01	.07
K ₂ O.....	5.00	4.37	4.16	4.40	4.96
H ₂ O at 110°.....	.35	.36	.25	.31	.49
H ₂ O above 110°.....	.25	.79	1.23	1.11	2.74
TiO ₂69	.38	.87	.87	.75
P ₂ O ₅26	.13	.40	.42	.32
Cr ₂ O ₃	trace	trace	none	none	none
V ₂ O ₃01	.01	.01	.02	.02
MnO.....	.10	trace	.11	.12	trace
BaO.....	.09	.28	.14	.09	.05
SrO.....	trace	trace	.09	trace	trace
Li ₂ O.....	trace	none	trace	trace	trace
CO ₂	none	none	none	.78	none
S.....	none	-----	none	none	-----
SO ₃	none	-----	none	none	.26
FeS ₂	-----	2.29	-----	-----	-----
Cl.....	trace	.06	.04	.04	trace
	99.54	99.99	99.78	99.81	99.49

Fluorine was not looked for in the analyses.

2. ROCKS FROM THE HENRY MOUNTAINS.

A. Porphyry. Not hitherto published. Reported by J. S. Diller as containing prominent plagioclase with some augite and hornblende in a crystalline groundmass of quartz and orthoclase. Analysis by R. B. Riggs, record No. 728.

B. Augite-porphyrite, dike, north spur of Mount Pennell. Contains hornblende, augite, and plagioclase in a feldspathic groundmass.

C. Hornblende-porphyrite, Mount Hillers. Contains plagioclase, hornblende, quartz, and magnetite.

Analyses B and C by W. F. Hillebrand, record No. 1428. Rocks described by Cross in 14th Ann., p. 165.

	A.	B.	C.
SiO ₂	63.16	60.98	62.88
Al ₂ O ₃	17.21	19.09	17.13
Fe ₂ O ₃	2.43	1.76	1.86
FeO	2.30	1.15	2.58
MgO	1.27	.65	1.48
CaO	6.27	3.67	5.39
Na ₂ O	4.70	6.70	4.50
K ₂ O	1.84	3.53	2.25
H ₂ O at 100°48	.16
H ₂ O above 100°69	.44	.42
TiO ₂21	.36	.51
P ₂ O ₅12	.10	.26
MnO	trace	.15	.16
SrO	trace?	.28	.12
BaO09	.43	.16
Li ₂ O	trace	trace	trace
CO ₂52	
SO ₃	trace		
	100.29	100.29	99.86

3. MISCELLANEOUS ROCKS.

A. Rhyolite, Thomas Range. Analysis made by L. G. Eakins in the Denver laboratory. Described by Cross in Proc. Colo. Sci. Soc., vol. 2, p. 69. Contains phenocrysts of quartz and sanidine in a ground-mass mainly made up of quartz and alkali feldspar.

B. Rhyolitic glass or pitchstone, edge of Gold Mountain mining district, 8 miles north of west from Marysvale. Identified by Whitman Cross, but not published. Sp. gr. 2.25 at 23.5°. Analysis by W. F. Hillebrand, record No. 1833.

	A.	B.
SiO ₂	74.49	70.17
Al ₂ O ₃	14.51	11.83
Fe ₂ O ₃57	.93
FeO32	none
MgO	trace	.06
CaO	1.03	.76
Na ₂ O	3.79	3.85
K ₂ O	4.64	3.74
H ₂ O64	8.72
TiO ₂17
MnO	trace
Li ₂ O	trace
	99.99	100.23

NEW MEXICO.

1. BASALTS FROM RIO GRANDE CANYON.

Described by Iddings, Bull. 66; also in Amer. Journ. Sci., 3d series, vol. 36, p. 220. Contain plagioclase, augite, olivine, and magnetite. A, B, and C contain quartz also; D is quartzless. Analyses by L. G. Eakins, record Nos. 847, 850.

A. Light gray, dense. P. R. C. 548.

B. Greenish black, dense.

C. Dark red, vesicular. P. R. C. 549.

D. Gray dense. P. R. C. 547.

	A.	B.	C.	D.
SiO ₂	52.27	52.37	51.57	52.38
Al ₂ O ₃	17.68	17.01	17.72	18.79
Fe ₂ O ₃	2.51	1.44	6.24	2.88
FeO.....	5.00	5.89	1.78	4.90
MgO.....	6.05	6.86	4.91	4.91
CaO.....	8.39	7.59	8.82	7.70
Na ₂ O.....	4.19	3.51	3.59	3.99
K ₂ O.....	1.58	1.59	1.99	1.76
H ₂ O.....	.82	1.29	.64	.53
TiO ₂	1.49	1.60	1.43	1.22
P ₂ O ₅56
MnO.....	.23	.32	.45	.18
BaO.....	.06	.06	.16	.11
CO ₂	trace	.37	.58
Cl.....	trace	trace
	100.27	99.90	99.88	99.91

2. MOUNT TAYLOR REGION.

Rocks received from J. S. Diller. Analyses by T. M. Chatard, record Nos. 219, 227, 228, 235, 271, 268, and 269. Not hitherto published. Petrographic details furnished by Diller.

A. Lava, canyon on east side of San Mateo Mountain.

B. Phonolitic andesite, canyon on east side of San Mateo Mountain. Contains feldspar and corroded augite, with sometimes olivine, in a groundmass of feldspar and a green ferromagnesian silicate.

C. Mica-andesite, canyon on east side of San Mateo Mountain. Shows prominent plagioclase, with augite and biotite.

D. Mica-andesite, 6 miles northeast of Grant's. Principally plagioclase and biotite, with less epidote, quartz, sphene, and carbonates.

E. Basalt, 6 miles northeast of Grant's. Contains chiefly plagioclase, augite, and olivine, with much magnetite.

F. Augite from E.

G. Feldspar from E. Analysis on three-fourths gramme of material.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	68.40	65.51	65.78	49.80	47.54	47.06	52.54
Al ₂ O ₃	17.99	16.89	17.32	15.33	16.73	7.77	31.26
Fe ₂ O ₃	2.66	1.41	3.68	-----	6.69	1.30	-----
FeO.....	1.63	2.52	.46	7.44	6.67	8.15	-----
MgO.....	.49	.39	.47	6.61	6.38	13.52	.28
CaO.....	.67	1.19	1.66	7.19	8.74	19.33	12.34
Na ₂ O.....	4.54	6.42	5.23	2.71	2.81	.33	3.55
K ₂ O.....	3.54	5.02	4.64	4.36	1.10	.11	.42
H ₂ O.....	.52	.16	.14	1.38	.36	.20	.26
TiO ₂	-----	.92	.27	2.67	2.76	1.82	undet.
P ₂ O ₅	-----	.07	.13	.73	.51	.06	-----
Cr ₂ O ₃	-----	-----	-----	-----	-----	trace	-----
MnO.....	.21	.31	.32	.30	.19	.20	-----
(CoNi)O.....	-----	-----	-----	-----	-----	traces	-----
BaO.....	-----	-----	-----	-----	.03	-----	trace ?
CO ₂	-----	-----	-----	2.56	(?)	-----	-----
	100.65	100.81	100.10	101.08	100.51	99.85	100.65

3. COLFAX COUNTY.

Descriptions, hitherto unpublished, supplied by Whitman Cross. Analyses by W. F. Hillebrand, record No. 1719.

A. Phonolite, Pleasant Valley. Contains much nephelite, some aegirite, alkali feldspar, a scanty dust of magnetite, and a few decomposed grains of noselite or sodelite. Sp. gr. 2.619, 22°; 40.8 per cent soluble in dilute (1:40) nitric acid, of which soluble portion 43.5 per cent is silica.

B. Pyroxene-andesite, Sierra Grande. Contains augite, less hypersthene, microliths of plagioclase, apatite, magnetite, and a smoky-brown glassy base. Sp. gr. 2.635, 21°.

C. Plagioclase-basalt, end of San Rafael flow. Contains plagioclase, augite, olivine, with much iddingsite, magnetite, and apatite. Sp. gr. 2.970, 21.5°.

D. Nepheline-basanite, Ciruella. Contains augite, olivine, nephelite, plagioclase, magnetite, apatite, and a little biotite. Sp. gr. 3.122, 22°. 0.55 per cent K_2O and 2.10 per cent Na_2O soluble in 1:40 nitric acid.

	A.	B.	C.	D.
SiO_2	56.24	60.16	48.35	42.35
Al_2O_3	21.43	15.34	15.47	12.29
Fe_2O_3	2.01	3.07	4.80	3.89
FeO55	2.18	7.58	7.05
MgO15	3.41	8.15	13.09
CaO	1.38	5.79	8.81	12.49
Na_2O	10.53	3.88	3.09	2.74
K_2O	5.74	2.59	.95	1.04
H_2O at 110°.....	.12	.25	.28	.32
H_2O above 110°.....	.86	1.79	.73	1.50
TiO_226	.84	1.33	1.82
P_2O_506	.46	.33	.99
ZrO_209	.01	none	none
Cr_2O_3	none	trace?	trace	.10
V_2O_504
NiO, CoO	none	trace	.02	.03
MnO08	.08	.21	.21
SrO03	.08	.03	.09
BaO08	.14	.06	.10
Li_2O	trace	trace	trace	trace
SO_310	.08	.07	.05
S.....	.03	trace	trace	trace
Cl.....	.12	undet.	undet.	undet.
F.....	trace	undet.	undet.	undet.
	99.86	100.15	100.26	100.19

4. MISCELLANEOUS ROCKS.

A. Obsidian, Obsidian Hill camp, Tewan Mountains. Described by Iddings, 7th Ann., p. 292. A rhyolitic obsidian, containing grains of iron oxide and a few microscopic feldspars. Resembles that from Obsidian Cliff in the Yellowstone National Park. Analysis by L. G. Eakins, record No. 851. Sp. gr., 2,352, $23^{\circ}.5$.

B. Trachyte (?), from Los Cerillos. Described by Diller, Bull. 42, p. 39. Analysis by F. W. Clarke, record No. 346. Rock composed chiefly of orthoclase, with a considerable amount of biotite, epidote, pyrite, and limonite, and some amorphous substance. It is the matrix or gangue rock of the Los Cerillos turquois.

	A.	B.
SiO ₂	76.20	56.68
Al ₂ O ₃	13.17	16.62
Fe ₂ O ₃34	6.28
FeO.....	.73
MgO.....	.19	.79
CaO.....	.42	.59
Na ₂ O.....	4.31	1.03
K ₂ O.....	4.46	11.18
H ₂ O.....	.33	3.28
TiO ₂	trace	.22
P ₂ O ₅73
FeS ₂	2.21
MnO.....	.10	1.02
CuO.....	trace
	100.25	100.63

ARIZONA.

A, B, C, D. Mica-basalt, Santa Maria Basin. See Iddings, Bull. Phil. Soc. Washington, vol. 12, p. 212. Not fully described. Analyses by W. F. Hillebrand, record No. 1261.

E. Hornblende-porphyrite, Sierra Carrizo. Described by Cross, 14th Ann., p. 165. Contains plagioclase and hornblende, in a ground-mass of quartz and orthoclase. Analysis by Hillebrand, No. 1429.

	A.	B.	C.	D.	E.
SiO ₂	49.36	55.35	57.04	57.48	63.18
Al ₂ O ₃	16.35	12.91	13.66	14.09	16.47
Fe ₂ O ₃	2.93	4.67	4.96	5.21	2.36
FeO.....	8.55	2.06	1.77	1.35	2.28
MgO	7.06	6.29	4.43	3.49	1.33
CaO.....	10.08	5.77	6.23	6.05	4.77
Na ₂ O	2.67	2.65	3.08	3.00	4.40
K ₂ O.....	.82	4.86	4.95	4.69	2.93
H ₂ O at 100°.....	.22	2.67	1.11	1.20	.27
H ₂ O above 100°.....	.65	1.18	1.10	1.37	.60
TiO ₂98	.87	.94	.94	.60
P ₂ O ₅30	.58	.63	.65	.28
MnO19	.08	.17	.09	.15
NiO, CoO.....	.05	.05	.07	.08	-----
SrO	none	trace	trace	trace	.09
BaO.....	.04	.19	.22	.23	.15
Li ₂ O	none	trace	trace?	none	trace
	100.25	99.98	100.36	99.92	99.86

F. Typical hypersthene-andesite, San Francisco Mountains. Not described. Analysis by T. M. Chatard, record No. 270.

G. Recent lava, 2 miles south of Mount Trumbull. Not described. Analysis by L. G. Eakins, record No. 1024.

	F.	G.
SiO ₂	64.82	45.30
Al ₂ O ₃	18.27	14.95
Fe ₂ O ₃	3.48	1.98
FeO56	9.32
MgO85	8.29
CaO	2.89	8.87
Na ₂ O	5.05	4.27
K ₂ O	2.67	1.27
H ₂ O20	.85
TiO ₂56	2.66
P ₂ O ₅23	2.23
MnO20	trace
	99.78	99.99

NEVADA.

1. ROCKS FROM WASHOE.

Described by Hague and Iddings, Bull. 17. Analyses by F. A. Gooch, record Nos. 119, 129. FeO not separately determined.

A. Rhyolite, south-southeast of McClellan Peak. Contains feldspars, orthoclase predominating over plagioclase, quartz, mica, and hornblende.

B. Dacite, spur northeast of McClellan Peak. More plagioclase than orthoclase, much mica, less hornblende, little quartz.

	A.	B.
SiO ₂	73.07	69.96
Al ₂ O ₃	11.78	15.79
Fe ₂ O ₃	2.30	2.50
MgO39	.64
CaO	2.02	1.73
Na ₂ O	1.19	3.80
K ₂ O	6.84	4.12
Ignition	2.24	1.53
	99.83	100.07

2. ROCKS FROM EUREKA.

Described by Hague and Iddings, Mon. XX.

A. Basalt, summit of Richmond Mountain. Red, porous. Contains augite, less hypersthene, feldspars, and magnetite, in a glassy base, with accessory olivine and quartz. Analysis by J. E. Whitfield, record No. 424.

B. Andesitic perlite, south of Carbon Ridge. Contains plagioclase, hornblende, biotite, quartz, hypersthene, augite, magnetite, apatite, and zircon, with a glassy base. Analysis by W. H. Melville, record No. 1240.

	A.	B.
SiO ₂	50.38	65.13
Al ₂ O ₃	19.83	15.73
Fe ₂ O ₃	6.05	2.24
FeO.....	2.00	1.86
MgO.....	5.36	1.49
CaO.....	10.03	3.62
Na ₂ O.....	2.15	2.93
K ₂ O.....	1.76	3.96
H ₂ O at 105°.....	1.37	.52
H ₂ O above 105°.....		1.91
TiO ₂	none	.58
P ₂ O ₅	none	.23
NiO.....	-----	.07
MnO.....	.38	trace
Li ₂ O.....	trace	-----
SO ₃83	-----
	100.14	100.27

CALIFORNIA.

1. MOUNT SHASTA.

Description of rocks furnished by J. S. Diller, who also describes A, B, and C in Bull. 150, pp. 221, 227.

A. Hornblende-andesite, Black Butte, west base of Shasta. Contains plagioclase and hornblende in a microlitic groundmass. The latter carries hypersthene, magnetite, and amorphous matter. Analysis by W. H. Melville, record No. 1346. P. R. C. 84.

B. Hypersthene-andesite, older flow, west base of Shasta. Contains plagioclase, hypersthene, magnetite, and glass. Analysis by Melville, No. 1346.

C. Hypersthene-andesite, like B, same locality, later flow. Analysis by Melville, No. 1346. P. R. C. 87.

D. Hornblende-andesite, late flow, eastern side of Shasta. Contains small crystals of plagioclase and hornblende in a dark groundmass. Thin section not examined. Analysis by H. N. Stokes, record No. 1532.

E. Ophitic basalt from near the McCloud River, south of Mount Shasta. A gray cellular rock. Thin section not examined. Analysis by Stokes, No. 1532.

F. Andesite basalt, Delta, Shasta County. Much plagioclase and a few hypersthene crystals in a groundmass chiefly of feldspar, pyroxene, magnetite, and trace of olivine. Analysis by Melville, No. 1346.

	A.	B.	C.	D.	E.	F.
Si ₂ O.....	64.48	64.52	63.03	61.58	47.94	55.08
Al ₂ O ₃	19.28	18.31	17.72	16.96	18.90	18.93
Fe ₂ O ₃	1.40	.90	2.27	1.75	2.21	2.02
FeO.....	1.78	2.51	1.92	2.85	8.59	5.56
MgO.....	1.64	2.35	3.63	3.67	8.21	5.17
CaO.....	5.06	5.11	5.97	6.28	9.86	8.40
Na ₂ O.....	4.41	4.64	3.92	3.94	2.81	4.23
K ₂ O.....	1.12	1.25	1.06	1.28	.29	.74
H ₂ O at 110°...	.06	.20	.44	.24	.39	.29
H ₂ O above 110°				1.06	.74	
TiO ₂	undet.	undet.	undet.	.49	.57	trace
P ₂ O ₅22	.15
MnO.....				trace	trace
SrO.....				trace	none
BaO.....				.03	none
Li ₂ O.....				trace	trace
	99.23	99.79	99.96	100.35	100.66	100.42

The following rocks are from Shasta County, but not within either the Mount Shasta or the Lassen Peak areas. Collected by J. S. Diller, who furnishes the petrographic data.

A. Dacite-porphyry, east fork of Clear Creek, 9 miles above French Gulch. Contains conspicuous phenocrysts of plagioclase, a few of quartz, and smaller ones of biotite and pyroxene, in a groundmass of quartz and feldspar. Analysis by J. E. Whitfield, record No. 970.

B. Dacite-porphyry, Smiths Gulch, 6 miles up Clear Creek from French Gulch. Contains phenocrysts of plagioclase, quartz, biotite, and hornblende, in a groundmass chiefly of quartz and feldspar. Analysis by Whitfield, No. 971. Also described by J. P. Iddings in Bull. 150, p. 233. P. R. C. 90.

C. Diorite, from Ono. Not described. Analysis by T. M. Chatard, record No. 1107.

	A.	B.	C.
SiO ₂ -----	66.30	64.24	68.10
Al ₂ O ₃ -----	17.55	18.67	15.18
Fe ₂ O ₃ -----	2.19	1.40	1.34
FeO-----	.55	1.96	1.70
MgO-----	.97	1.48	2.06
CaO-----	3.12	4.11	4.66
Na ₂ O-----	5.15	4.14	3.71
K ₂ O-----	2.45	1.71	1.48
H ₂ O-----	1.25	1.18	.55
TiO ₂ -----	trace	.76	.35
P ₂ O ₅ -----	.15	.08	.18
MnO-----	trace	trace	.20
BaO-----			.06
SO ₃ -----	.28	.22	
Cl-----		.25	
Less O-----	99.96	100.20	99.57
		.05	
		100.15	

2. LASSEN PEAK REGION.

Rocks collected by J. S. Diller, who has furnished the petrographic data. Nearly all are from the area covered by the Lassen Peak atlas sheet of the U. S. Geological Survey. The quartz basalts have been described by Diller in Bull. 79, and partly in Amer. Journ. Sci., 3d series, vol. 33, p. 49. The analyses are so numerous that it seems best to divide them into subordinate groups.

1ST. RHYOLITES.

A. Rhyolite, a short distance northwest of Willow Lake, near the Geyser, Plumas County. A light-gray rock with occasional phenocrysts of quartz and feldspar in a granular groundmass of the same materials. Analysis by W. F. Hillebrand, record No. 414.

B. Rhyolite, 2 miles northwest of Deer Creek Meadows, Tehama County. Shows many small crystals of quartz, feldspar, and biotite in a spherulitic groundmass. Analysis by Hillebrand, No. 415.

C. Rhyolite, Slate Creek, west of Deer Creek Meadows. Composed wholly of spherulites, some of which include crystals of feldspar or biotite. Analysis by Hillebrand, No. 416.

D. Rhyolite, summit of Mount Stover, Plumas County. Composed almost wholly of spherulites, with a few crystals of feldspar and hornblende. Analysis by Hillebrand, No. 417.

E. Rhyolite, near Slate Creek, west of Deer Creek Meadows. A pearlite, composed chiefly of little glass balls, with a few spherulites, and crystals of feldspar and biotite. Analysis by Hillebrand, No. 418.

F. Rhyolite, $1\frac{1}{2}$ miles northeast of Clipper Mills, on tramway to Rock Creek, Shasta County. A spherulitic rock containing a few microscopic particles of feldspar and hornblende. Analysis by Hillebrand, No. 678.

G. Rhyolite-tuff, divide between the west fork of Willards Creek and the stage road, in Lassen County. Elevation, 5,800 feet. Composed almost wholly of fine angular particles of clear glass. Analysis by George Steiger, record No. 1427.

Materials for A to F dried at 110° previous to analysis.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂	74.24	74.65	73.62	72.40	73.64	74.60	70.01
Al ₂ O ₃	14.50	14.11	14.24	14.81	13.44	13.41	12.61
Fe ₂ O ₃	1.27	1.08	.93	.81	.60	1.28	1.47
FeO.....	.67	.29	.67	.88	.74	.30	.50
MgO.....	.25	.20	.33	.47	.26	.26	.72
CaO.....	.11	.80	1.07	1.94	1.26	1.08	1.06
Na ₂ O.....	3.00	2.81	3.25	3.91	3.51	3.38	1.94
K ₂ O.....	3.66	4.59	4.28	3.90	4.50	4.50	5.12
H ₂ O at 100°.....							2.37
H ₂ O above 100°.....	2.04	1.40	1.29	.59	1.99	.85	4.68
TiO ₂20	.21	.21	.18	.11	.16	-----
P ₂ O ₅07	trace	.02	.03	.06	.03	.04
MnO.....	.06	.11	.08	.07	.06	.06	trace
SrO.....	trace	trace	trace	.04	.02	none	-----
BaO.....	.18	.08	.10	.10	.11	.11	-----
Li ₂ O.....	none	none	none	trace	trace	trace	-----
SO ₃03	-----	-----	-----	-----	-----	-----
	100.28	100.33	100.09	100.13	100.30	100.02	100.52

2D DACITES AND ANDESITES.

A. Gray dacite, Lassen Peak. Contains hornblende, biotite, plagioclase, quartz, scarce pyroxene, magnetite, apatite, and a glassy base. Analysis by T. M. Chatard, record No. 111. P. R. C. 82.

B. Secretion in dacite, Lassen Peak. Composed chiefly of plagioclase and hornblende. Analysis by Chatard, No. 110.

C. Reddish dacite, Lassen Peak. Essentially like A. Analysis by Chatard, No. 110. P. R. C. 82.

D. Dacite, near the timber line, west base of Lassen Peak. Small phenocrysts of plagioclase and hornblende, with a few of quartz, in a reddish-gray groundmass containing much amorphous matter. Analysis by W. F. Hillebrand, record No. 668. P. R. C. 82.

E. Secretion in D. Composed essentially of plagioclase and hornblende. Analysis by Hillebrand, No. 669. P. R. C. 82.

Rocks A, C, D, and E described by Diller in Bull. 150, p. 217.

	A.	B.	C.	D.	E.
SiO ₂	69.51	58.97	68.20	68.32	55.14
Al ₂ O ₃	15.75	18.60	16.98	15.26	19.10
Fe ₂ O ₃	3.34	5.94	3.75	1.66	6.16
FeO.....				1.26	.54
MgO.....	2.09	6.89	2.07	1.32	4.23
CaO.....	1.71	2.84	4.33	3.26	8.36
Na ₂ O.....	3.89	3.05	2.98	4.27	3.71
K ₂ O.....	3.34	2.24	1.52	2.81	1.04
H ₂ O.....	.56	1.35	.44	1.37	.91
TiO ₂31	.52
P ₂ O ₅	trace	undet.		.12	.18
MnO.....				.04	.11
SrO.....				trace	.07
BaO.....				.07	trace
Li ₂ O.....				trace	trace
	100.19	99.88	100.27	100.07	100.07

F. Dacite, east end of Chaos, northwest base of Lassen Peak. The youngest dacite of the region. Contains quartz, feldspar, biotite, and hornblende, embedded in a clear puniceous glass. Analysis by W. F. Hillebrand, record No. 670. Described by Diller in Bull. 150, p. 218.

G. Secretion in F. Consists chiefly of plagioclase and hornblende, with some olivine and clear glass. Analysis by Hillebrand, No. 671.

H. Dacite (?), west side of old crater rim near the Thumb, at the head of Mill Creek, Shasta County. Shows phenocrysts of hornblende, plagioclase, and pyroxene, and apparently of quartz, in a gray, micro-litic groundmass. Analysis by Hillebrand, No. 674.

I. Streaked dacite, falls of south fork of Bear Creek, Shasta County. Contains plagioclase with a little sanidine, hornblende, quartz, magnetite, some pyroxene inclusions, and glass base. Analysis by R. B. Riggs, record No. 524. P. R. C. 80. Described by Diller in Bull. 150, p. 213.

J. Dacite-tuff, Rice's quarry, 6 miles southeast of Paskenta, Tehama County. Clear glass, with fragments of quartz, feldspar, and hornblende. Analysis by George Steiger, record No. 1427.

	F.	G.	H.	I.	J.
SiO ₂	68.72	53.35	63.81	68.10	65.78
Al ₂ O ₃	15.15	19.22	17.07	15.50	14.87
Fe ₂ O ₃	1.16	3.28	2.11	3.20	1.27
FeO.....	1.76	4.48	2.15	none	1.60
MgO.....	1.28	4.86	2.28	.10	1.89
CaO.....	3.30	9.76	4.97	3.02	2.41
Na ₂ O.....	4.26	2.89	4.08	4.20	2.58
K ₂ O.....	2.78	.99	1.96	3.13	2.71
H ₂ O at 100°.....					2.87
H ₂ O above 100°.....	.74	.77	1.03	2.72	4.32
TiO ₂31	.56	.38	.15	
P ₂ O ₅09	.10	.10	.03	.08
MnO.....	.11	.15	.09	trace	trace
SrO.....	.03	.03	.03	trace	
BaO.....	.07	trace ?	.04	.06	
Li ₂ O.....	trace	trace	trace	none	
	99.76	100.44	100.10	100.21	99.78

K. Pyroxene-andesite, west end of Butte Mountain, Plumas County. Prominent phenocrysts of pyroxene and minute ones of plagioclase, in a dark groundmass containing much globulitic matter. Analysis by W. F. Hillebrand, record No. 411.

L. Pyroxene-andesite, south base of Burney Butte, Shasta County. Numerous small phenocrysts of plagioclase and a few of pyroxene, in a gray groundmass containing much amorphous matter. Analysis by R. B. Riggs, record No. 684.

M. Hornblende-andesite, Tuscan Buttes, 7 miles east of Red Bluff. A few small phenocrysts or fragments of hornblende, in a groundmass consisting mainly of plagioclase and gray microlitic matter. Analysis by Hillebrand, No. 412.

N. Hornblende-andesite, near Buntingville, Lassen County. A few phenocrysts of hornblende in a groundmass consisting mainly of small feldspars. Analysis by T. M. Chatard, record No. 413.

O. Hornblende-andesite, northwest summit, head of Burney Creek, Shast. County. Inconspicuous plagioclase and, rarely, olivine, in a groundmass of plagioclase and pyroxene. Numerous dark spots are due to altered hornblende. Analysis by Riggs, No. 683.

Rocks in this group dried at 105° before analysis.

	K.	L.	M.	N.	O.
SiO ₂	55.53	62.44	60.93	67.89	60.04
Al ₂ O ₃	17.63	16.39	18.56	17.29	17.43
Fe ₂ O ₃	2.81	4.66	2.68	2.39	5.39
FeO.....	3.59	1.00	2.19	.21	.53
MgO.....	5.85	2.65	2.37	.66	3.51
CaO.....	8.74	6.22	6.63	3.01	6.65
Na ₂ O.....	3.09	3.16	3.79	5.11	4.15
K ₂ O.....	.92	2.25	1.33	1.69	1.24
H ₂ O.....	1.24	1.02	.90	1.34	.90
TiO ₂56	.31	.61	.21	.49
P ₂ O ₅21	.05	.18	.12	.04
MnO.....	.08	trace	.10	.12	.08
SrO.....	.06	trace	.12	.04	?
BaO.....	.02	.03	.02	.03	.04
Li ₂ O.....	none	trace	none	-----	trace
SO ₃	-----	trace	-----	-----	trace
	100.33	100.18	100.41	100.11	100.49

P. Hypersthene-andesite, 1 mile west of summit on Bidwell's road, Butte County. Rich in small phenocrysts of plagioclase and pyroxene, mostly hypersthene, in a groundmass of the same minerals, with magnetite, and probably some amorphous matter. Analysis by W. F. Hillebrand, record No. 410.

Q. Hypersthene-andesite, old crater at head of Mill Creek. Contains small phenocrysts of plagioclase and hypersthene, in a groundmass of plagioclase, pyroxene, magnetite, etc. Some greenish pseudomorphs suggest former olivine. Analysis by T. M. Chatard, record No. 409.

R. Hypersthene-andesite, 2 miles south of Suppans Mountain, Tehama County. Abundant plagioclase and hypersthene, with traces of hornblende, in a microlitic groundmass. Analysis by Hillebrand, No. 672.

S. Secretion in R. Composed chiefly of plagioclase and hypersthene, with some quartz and amorphous matter. Analysis by Hillebrand, No. 673.

T. Hypersthene-andesite, west base of Suppans Mountain, near Lassen Peak, Tehama County. Contains numerous microscopic crystals of plagioclase and hypersthene, in a microlitic groundmass. Analysis by Hillebrand, No. 676.

Rocks dried at 100° to 110° before analysis.

	P.	Q.	R.	S.	T.
SiO ₂	55.20	57.11	63.47	57.04	58.08
Al ₂ O ₃	18.68	17.78	16.75	19.11	18.37
Fe ₂ O ₃	3.14	3.54	2.15	4.37	2.92
FeO.....	4.42	2.74	2.75	2.48	3.38
MgO.....	4.59	3.41	3.04	3.94	3.35
CaO.....	8.02	7.21	5.72	7.34	7.05
Na ₂ O.....	3.66	3.81	3.94	3.48	3.66
K ₂ O.....	1.01	1.86	1.62	1.16	1.33
H ₂ O.....	.51	.98	.55	1.09	1.09
TiO ₂92	.95	.37	.47	.44
P ₂ O ₅24	.26	.13	.08	.16
MnO.....	.14	.33	.09	.12	.13
SrO.....	.02	trace?	.04	.02	.02
BaO.....	.03	.03	.04	trace?	.03
Li ₂ O.....	none	-----	trace	trace	trace
	100.58	100.01	100.66	100.70	100.01

U. Hypersthene-andesite, 1 mile southwest of Thumb, head of Bailey Creek, near Lassen Peak. Abundant but inconspicuous plagioclase and hypersthene, in a microlitic groundmass containing many small crystals of plagioclase. Analysis by W. F. Hillebrand, record No. 675.

V. Hypersthene-andesite, west summit of Crater Peak, Shasta County. Microphenocrysts of feldspar and hypersthene in a groundmass consisting largely of the same minerals, with some amorphous matter. Analysis by Hillebrand, No. 679.

W. Hypersthene-andesite, north slope of Crater Peak. Phenocrysts of plagioclase and hypersthene in a groundmass containing much dark amorphous matter. Analysis by Hillebrand, No. 680.

X. Secretion in W. Composed chiefly of plagioclase, hypersthene, and a globulitic base. Analysis by Hillebrand, No. 681.

Y. Andesitic tuff, Stillwater Creek, 8 miles northeast of Redding. Contains plagioclase, hornblende, rare hypersthene, magnetite, and glass. Fragments of andesite are inclosed. Analysis by W. H. Melville, record No. 1346. Described by Diller in Bull. 150, p. 211.

Rocks dried at 100° to 110° before analysis.

	U.	V.	W.	X.	Y.
SiO ₂	59.84	68.12	61.17	53.85	69.51
Al ₂ O ₃	16.81	16.24	17.74	18.53	15.61
Fe ₂ O ₃	1.88	1.26	1.78	1.96	.56
FeO	3.60	2.08	3.51	5.30	1.27
MgO	3.85	1.35	2.76	5.88	.61
CaO	6.30	3.80	5.90	9.66	2.80
Na ₂ O	3.63	3.89	3.79	2.98	3.43
K ₂ O	2.13	2.54	1.71	.74	2.81
H ₂ O	1.04	.40	.83	.45	3.63
TiO ₂57	.25	.45	.50	trace
P ₂ O ₅19	.14	.14	.05
Cr ₂ O ₃	trace?	none	none	trace
MnO14	.10	.12	.12
SrO02	.02	.04	.04
BaO07	.09	.06	.03
Li ₂ O	trace	trace	trace	trace
	100.07	100.28	100.00	100.09	100.23

3D. BASALTS.

The quartz-basalts are described by Diller in Bull. 79. That from Mitylene was analyzed for comparison with the Cinder Cone series.

A. Quartz-basalt, Cinder Cone, 10 miles northeast of Lassen Peak. Contains plagioclase, pyroxene (mostly hypersthene), olivine, quartz, and much unindividualized base; the latter about 25 per cent. Magnetite is also present; augite occurs sparingly. Analysis by W. F. Hillebrand, record No. 407. P. R. C. 96. Also described in Bull. 150, p. 252.

B. Volcanic bomb from quartz-basalt, Cinder Cone. Analysis by Hillebrand, No. 665.

C. Lapilli from quartz-basalt, Cinder Cone. Analysis by Hillebrand, No. 667. P. R. C. 96. Also described in Bull. 150, p. 249.

D. Volcanic sand, one-half mile northeast of Cinder Cone. Analysis by Hillebrand, No. 663.

E. White pumiceous inclosure from quartz-basalt, Cinder Cone. Analysis by Hillebrand, No. 664. Mainly glass.

Rocks dried at 100° to 110° before analysis.

	A.	B.	C.	D.	E.
SiO ₂	57.25	56.70	56.53	55.93	79.49
Al ₂ O ₃	16.45	15.75	17.50	17.34	11.60
Fe ₂ O ₃	1.67	1.29	1.35	1.50	.33
FeO.....	4.72	5.32	5.03	5.20	.49
MgO.....	6.74	7.16	5.94	7.29	.09
CaO.....	7.65	7.67	8.07	8.04	1.64
Na ₂ O.....	3.00	3.36	3.51	3.32	4.04
K ₂ O.....	1.57	1.56	1.55	1.35	1.52
H ₂ O.....	.40	.30	.27	.26	.68
TiO ₂60	.65	.54	undet.	undet.
P ₂ O ₅20	.20	.15	undet.	undet.
Cr ₂ O ₃		trace	trace		
MnO.....	.10	.19	.12	undet.	none
SrO.....	trace	trace	trace?	(?)	(?)
BaO.....	.03	.03	trace	(?)	(?)
Li ₂ O.....	none	trace	trace	(?)	(?)
	100.38	100.18	100.56	100.23	99.88

F. Quartz-basalt, one-half mile south of Cinder Cone, on border of lava field. Analysis by Hillebrand, No. 666.

G. Quartz-basalt, west end of Lake Bidwell, on border of Cinder Cone lava field. Contains a few grains of quartz, much olivine and plagioclase, less pyroxene, and a globulitic base. Analysis by Hillebrand, No. 661.

H. Quartz-basalt, Silver Lake, near Lassen Peak. Contains occasional grains of quartz, much feldspar and olivine, less pyroxene, and brownish base. Analysis by Hillebrand, No. 662.

I. Quartz-basalt, resting on dacite, near west base of Lassen Peak. Analysis by Hillebrand, No. 677.

J. Quartz-basalt, island of Mitylene, coast of Asia Minor. Analysis for comparison with the Cinder Cone series, by T. M. Chatard, record No. 845.

Rocks F to I dried at 100° to 110° before analysis.

	F.	G.	H.	I.	J.
SiO ₂	54.56	56.18	57.59	56.51	56.58
Al ₂ O ₃	16.04	16.59	16.49	18.10	14.88
Fe ₂ O ₃95	1.51	1.22	4.26	2.31
FeO.....	6.07	5.51	4.89	2.68	3.04
MgO	8.71	7.26	7.72	4.52	3.76
CaO.....	8.89	7.64	7.40	8.15	8.69
Na ₂ O	3.05	3.58	3.62	3.23	3.36
K ₂ O.....	1.18	1.47	.99	1.15	2.18
H ₂ O at 105°.....					.69
H ₂ O above 105°28	.42	.86	.69	1.43
TiO ₂53	undet.	undet.	.48	.77
P ₂ O ₅18	undet.	undet.	.14	.15
Cr ₂ O ₃	trace			trace	trace?
MnO17	undet.	undet.	.11	.16
SrO	trace	(?)	(?)	.04	
BaO.....	.03	(?)	(?)	.04	.07
Li ₂ O	trace	(?)	(?)	trace	
CO ₂					2.32
	100.64	100.16	100.78	100.10	100.39

K. Recent basalt, Pit River. Rich in feldspar and augite, poor in olivine. Partial analysis by F. W. Clarke, record No. 109.

L. Basalt, 1 mile southeast of Paynes Creek, on the road from Red Bluff to Lassen Peak. A normal basalt, rather rich in olivine. Analysis by T. M. Chatard, record No. 405.

M. Basalt, summit of Inskip Crater, 25 miles east of Red Bluff. Mainly feldspar and augite, with a few phenocrysts of olivine. Analysis by Hillebrand and Chatard, record No. 406.

N. Basalt from the cone at south base of Burney Butte, Shasta County. Composed of plagioclase and augite, with some olivine and a globulitic base. Analysis by R. B. Riggs, record No. 685.

O. Basalt, near eastern end of rim of Crater Peak, Shasta County. Contains plagioclase and pyroxene, some of the latter being hypersthene with a trace of olivine. Analysis by R. B. Riggs, record No. 682.

P. Hornblende-basalt, Kosk Creek, near its mouth, by the great bend of Pit River, Shasta County. Contains abundant phenocrysts of hornblende, with a few of plagioclase, pyroxene, and olivine, in a ground-mass of plagioclase, augite, and magnetite. The hornblendes are deeply corroded, and some have disappeared, leaving groups of magnetite grains to mark their former presence. Analysis by L. G. Eakins, record No. 1022. Described by Diller in Amer. Geologist, vol. 19, p. 253.

Rocks dried at 105° to 110° before analysis, except in the case of the rock marked L.

	K.	L.	M.	N.	O.	P.
SiO ₂	51.92	47.93	50.89	52.63	52.95	44.77
Al ₂ O ₃	19.76	18.51	16.76	17.62	18.25	17.82
Fe ₂ O ₃	11.21	2.07	3.86	6.49	4.36	5.05
FeO.....		7.25	4.69	3.10	4.19	6.95
MgO.....	3.38	9.03	8.49	5.64	4.93	8.22
CaO.....	9.30	11.14	11.72	8.62	8.73	10.36
Na ₂ O.....	2.16	2.28	2.61	3.38	3.57	2.13
K ₂ O.....	.60	.24	.32	1.73	.77	.92
H ₂ O.....	1.54	.76	.41	.79	1.47	2.64
TiO ₂73	.79	.07	.66	.53
P ₂ O ₅11	.09	.47	trace	.72
MnO.....		.20	.13	trace	.12	trace
SrO.....				trace	trace
BaO.....			trace	.04	.01
Li ₂ O.....				trace	(?)
CO ₂		none			
SO ₃				trace	trace
	99.87	100.25	100.76	100.58	100.01	100.11

3. PLUMAS COUNTY.

Other rocks from this county are described under the heading of the Lassen Peak area. The following rocks, with two exceptions, were collected by H. W. Turner, who supplies the descriptions:

A. Granite, dike in serpentine, south slope of Grizzly Hill. Described by Turner in Amer. Geologist, vol. 17, p. 375. Contains quartz, albite, and muscovite. Analysis by H. N. Stokes, record No. 1562. P. R. C. 757.

B. Meta-rhyolite, near Tower Rock, Grizzly Mountains. Described by Turner in 14th Ann., p. 441. Contains porphyritic quartz, feldspar, and pyrite, in a fine groundmass. Analysis by W. F. Hillebrand, record No. 1273. P. R. C. 741.

C. Meta-rhyolite, near Greenville. Collected by Diller, who finds phenocrysts of quartz in a groundmass chiefly of quartz and feldspar. Analysis by Hillebrand, record No. 1458.

D. Granodiorite, southwest base of Mount Ingalls. Description supplied by Turner. Contains plagioclase, quartz, orthoclase, brown mica, green hornblende, iron oxide, and a little apatite, sphene, and epidote. Analysis by Hillebrand, record No. 1456. P. R. C. 727.

E. Granodiorite, Spanish Peak. Description supplied by Turner. Contains plagioclase, quartz, orthoclase, biotite, hornblende, iron ore, and apatite; also abundant secondary epidote and chlorite. Analysis by Stokes, record No. 1562. P. R. C. 756.

	A.	B.	C.	D.	E.
SiO ₂	76.00	73.25	72.77	67.33	59.68
Al ₂ O ₃	14.88	13.25	13.00	15.93	17.09
Fe ₂ O ₃65	-----	1.28	1.90	2.85
FeO.....	.10	1.74	2.65	1.59	2.75
MgO.....	.06	.28	.67	1.63	3.54
CaO.....	.19	2.23	2.47	4.09	6.62
Na ₂ O.....	3.52	2.69	4.95	3.76	3.87
K ₂ O.....	2.77	3.79	.34	2.46	1.31
H ₂ O at 100°.....	.20	.07	.07	.19	.15
H ₂ O above 100°.....	1.42	1.03	1.16	.66	1.00
TiO ₂04	trace	.22	.36	.65
P ₂ O ₅11	trace	.04	.11	.25
MnO.....	trace	trace	.08	.09	trace
SrO.....	-----	trace?	trace	trace	trace
BaO.....	trace	trace	trace	.08	.04
Li ₂ O.....	-----	trace	trace	trace	trace
CO ₂	-----	1.05	.47	-----	.20
SO ₃	trace	-----	-----	-----	trace
Cl.....	trace	-----	-----	-----	.03
F.....	trace	-----	-----	-----	-----
FeS ₂	-----	.58	-----	-----	-----
	99.94	99.96	100.17	100.18	100.03

F. Rhyolite, $3\frac{1}{2}$ miles southwest of Grizzly Peak. Description furnished by Turner. Contains sanidine, with less quartz and biotite in a glassy groundmass. Analysis by Hillebrand, record No. 1461 P. R. C. 776.

G. Hornblende-andesite, 4 miles from Pilot Peak. Described by Turner in 14th Ann., p. 441. Contains plagioclase and hornblende in a groundmass carrying grains of magnetite. Analysis by Hillebrand, record No. 1432. P. R. C. 716.

H. Hornblende-pyroxene-andesite, southwest base of Mount Ingalls. Description supplied by Turner. Contains plagioclase, rhombic pyroxene, augite, brown hornblende, and magnetite, with much glass in the groundmass. Analysis by Hillebrand, record No. 1456. P. R. C. 728.

I. Hypersthene-andesite, Franklin Hill. Description supplied by Turner. Contains plagioclase, rhombic pyroxene, augite, and magnetite. Probably no glass. Analysis by Hillebrand, record No. 1548 P. R. C. 754.

	F.	G.	H.	I.
SiO ₂	71.39	60.20	58.47	56.88
Al ₂ O ₃	14.13	17.21	18.80	18.25
Fe ₂ O ₃63	3.12	3.34	2.35
FeO37	2.69	2.64	4.45
MgO08	3.18	2.69	4.07
CaO	1.01	6.04	6.60	7.53
Na ₂ O	2.89	3.35	3.58	3.29
K ₂ O	5.69	1.44	2.01	1.42
H ₂ O at 100°42	1.12	.14	.24
H ₂ O above 100°	3.32	1.18	.92	.50
TiO ₂17	.57	.51	.45
P ₂ O ₅03	.17	.22	.30
MnO	trace	.12	.13	.18
SrO	trace	trace	.05	.04
BaO09	.11	.09	.11
Li ₂ O	trace	trace	trace	trace
	100.22	100.50	100.19	100.06

J. Dolerite, Mount Ingalls. Described by Turner in 14th Ann., p. 441. Contains plagioclase, augite, hypersthene, magnetite, and a few olivines. Analysis by W. F. Hillebrand, record No. 1273. P. R. C. 739.

K. Dolerite, Mount Ingalls. Also in 14th Ann., p. 441. Like J, but with scarcely any olivine. Analysis by Hillebrand, record No. 432. P. R. C. 740.

L. Basalt, 4 miles southeast of Mount Ingalls. Also in 14th Ann., p. 441. Contains plagioclase, olivine, augite, and magnetite. Analysis by Hillebrand, record No. 1273.

M. Olivine-basalt, $1\frac{1}{2}$ miles from Franklin Hill. Contains plagioclase, augite, partly altered olivine, magnetite, and probably some glass. Description supplied by Turner. Analysis by George Steiger, record No. 1596. P. R. C. 755.

N. Serpentine, Greenville. Described by Diller in Bull. 150, p. 372. Besides serpentine, the rock contains some magnetite and less chromite, with remnants of the pyroxene from which the serpentine was in great part derived. Analysis by W. H. Melville, record No. 1346. P. R. C. 145.

	J.	K.	L.	M.	N.
SiO ₂	53.91	52.81	50.56	51.21	39.14
Al ₂ O ₃	17.95	16.60	14.71	17.59	2.08
Fe ₂ O ₃	2.21	2.66	3.54	4.71	4.27
FeO.....	4.80	6.13	8.90	4.42	2.04
MgO.....	5.52	6.12	4.07	7.12	39.84
CaO.....	10.40	10.14	7.58	10.36	trace
Na ₂ O.....	2.90	2.79	2.94	2.49	-----
K ₂ O.....	1.34	1.05	2.10	.91	-----
H ₂ O at 100°.....	.20	.38	1.06	.58	} 12.70
H ₂ O above 100°.....	.20	.54	1.12	1.07	
TiO ₂52	.84	1.71	.31	-----
P ₂ O ₅21	.23	1.14	.09	-----
MnO.....	.10	undet.	.13	trace	-----
SrO.....	trace	trace	trace ?	-----	-----
BaO.....	.05	.03	.25	none	-----
Li ₂ O.....	trace	trace	trace ?	none	-----
Chromite.....	-----	-----	-----	-----	.11
	100.31	100.32	99.81	100.86	100.18

4. BUTTE COUNTY.

Rocks from this county are also to be found under the heading of the Lassen Peak region. The following rocks were collected by H. W. Turner, to whom the petrographic data are due. Analyses, with two exceptions, by W. F. Hillebrand, record Nos. 1432, 1456, 1461, and 1548. Analysis G is by H. N. Stokes, record No. 1562.

A. Granodiorite, north side of south fork of Feather River, opposite Enterprise. Described in 14th Ann., p. 441. Contains plagioclase, potash feldspar, quartz, hornblende, brown mica, and accessory minerals. The ferromagnesian minerals are largely altered to chlorite. P. R. C. 720.

B. Granodiorite, 2 miles east of Bangor. Composition like A. The mica is largely altered to chlorite. See 14th Ann., p. 441. P. R. C. 717.

C. Diorite, South Honcut Creek. Description supplied by Turner. Contains feldspar, probably all plagioclase, brown hornblende, and a little chlorite. P. R. C. 775.

D. Quartz-diorite, 4.6 miles south of Table Mountain, on ridge between Butte and Plumas counties. Described in 17th Ann., Part I, p. 521. Contains hornblende, feldspar, quartz, rutile, and a little secondary chlorite and epidote. P. R. C. 758.

E. Amphibole separated from E. Analysis by William Valentine, record No. 1723. Cr_2O_3 determination by Hillebrand.

	A.	B.	C.	D.	E.
SiO_2	70.36	63.43	57.87	54.64	50.08
Al_2O_3	15.47	14.20	16.30	12.06	7.97
Fe_2O_398	1.54	1.71	1.81	2.69
FeO	1.17	4.56	3.86	5.03	6.71
MgO87	2.35	5.50	11.86	16.31
CaO	3.18	5.51	5.53	7.74	11.21
Na_2O	4.91	3.49	5.01	2.35	1.22
K_2O	1.71	2.19	.75	1.01	.46
H_2O at 100°06	.15	.26	.12
H_2O above 100°	1.00	1.50	2.40	2.44	1.40
TiO_220	.73	.53	.61	.76
P_2O_511	.11	.27	.08	trace
V_2O_303
Cr_2O_316
NiO05
MnO	trace	.03	.08	.13	.49
SrO	trace	trace	trace	trace
BaO06	.06	.05	.05	none
Li_2O	trace	none	trace	trace
	100.08	99.85	100.12	100.01	100.46

F. Meta-andesite-tuff. Described in 14th Ann., p. 441. Contains plagioclase, augite, epidote, chlorite, and secondary hornblende. P. R. C. 719.

G. Uralite-diorite, 1 mile southeast of Forbestown. Described in 17th Ann., Part I, p. 521. Contains plagioclase, hornblende, and magnetite. P. R. C. 751.

H. Basalt, Oroville, Table Mountain. Described in 14th Ann., p. 441. Contains plagioclase, olivine, augite, and magnetite. P. R. C. 718.

I. Altered peridotite, 5 miles northeast of Strawberry Valley. Largely serpentine, with olivine, hornblende, magnetite, and calcite or dolomite. P. R. C. 742.

	F.	G.	H.	I.
SiO ₂	54.66	51.07	50.66	44.81
Al ₂ O ₃	15.85	14.93	13.97	<i>a</i> 1.88
Fe ₂ O ₃	1.82	6.44	2.55	1.98
FeO.....	5.12	5.98	10.20	4.52
MgO.....	5.64	4.84	4.45	30.91
CaO.....	8.75	7.89	8.08	6.58
Na ₂ O.....	3.46	5.04	3.32	} .15
K ₂ O.....	.47	.16	1.95	
H ₂ O at 100°.....	.25	.24	.27	.15
H ₂ O above 100°.....	2.48	1.73	.43	6.88
TiO ₂67	1.65	2.39	-----
P ₂ O ₅15	.19	1.01	.02
Cr ₂ O ₃	-----	-----	-----	.29
MnO.....	.18	.22	.29	.13
NiO.....	trace?	-----	trace	.09
SrO.....	trace	-----	trace	none
BaO.....	.04	-----	.22	none
Li ₂ O.....	none	-----	none	-----
CO ₂39	-----	-----	1.79
Cl.....	-----	trace	.02	-----
F.....	-----	trace	-----	-----
SO ₃	-----	trace	-----	-----
FeS ₂09	-----	-----	-----
	100.02	100.38	99.81	100.18

a Includes possible TiO₂.

5. SIERRA COUNTY.

Rocks collected and described by H. W. Turner. See paper in 17th Ann., Part I, p. 521. Additional details, supplied by Turner, are given here. Analyses A to H by W. F. Hillebrand, record Nos. 1456 and 1548. Analysis I by H. N. Stokes, record No. 1514.

A. Granulite (aplite), Yuba Gap, road east of Sierra Buttes. Contains orthoclase, microcline, quartz, plagioclase, some shreds of greenish mica, and a little iron ore, chlorite, and apatite. P. R. C. 730.

B. Granulite (aplite), dike east of Milton. Contains orthoclase, quartz, plagioclase, a little microcline, brown mica, and iron ore. P. R. C. 734.

C. Biotite-quartz-monzonite, Indian Valley. Contains plagioclase, orthoclase, quartz, brown mica, apatite, and iron ore. P. R. C. 737.

D. Quartz-diorite-porphry, dike in Indian Valley granite. Contains plagioclase, hornblende, biotite, and quartz. P. R. C. 738.

E. Quartz-mica-diorite, large area east of Milton. Contains plagioclase, a turbid feldspar which is apparently not orthoclase, quartz, green hornblende, brown mica, iron ore, and apatite. P. R. C. 732.

	A.	B.	C.	D.	E.
SiO ₂	76.03	75.97	68.65	66.65	57.26
Al ₂ O ₃	13.39	13.07	16.34	17.61	16.51
Fe ₂ O ₃48	.61	.93	.93	3.27
FeO.....	.31	.39	1.48	1.67	5.19
MgO.....	.05	.14	1.29	1.26	3.41
CaO.....	1.28	1.49	3.07	4.44	6.69
Na ₂ O.....	2.98	2.51	4.85	4.59	2.65
K ₂ O.....	5.18	5.62	1.85	1.70	2.93
H ₂ O at 100°.....	.15	.14	.24	.03	.20
H ₂ O above 110°.....	.34	.24	.62	.41	.95
TiO ₂07	.09	.28	.33	.53
P ₂ O ₅03	trace	.15	.18	.30
MnO.....	trace	trace	.08	.07	.18
SrO.....	trace	.03	.0706
BaO.....	.04	.14	.09	.12	.10
Li ₂ O.....	none	trace	trace	trace	trace
	100.33	100.44	99.99	99.99	100.23

F. Diabase-porphry, dike east of Milton. Contains labradorite and other plagioclase, augite, and hornblende, the last mineral being perhaps secondary. P. R. C. 733.

G. Hypersthene-andesite, point northeast of Goodyears Bar. Contains plagioclase and rhombic pyroxene, a little augite, and scales which seem to represent former biotite, now replaced by magnetite. P. R. C. 731.

H. Hornblende-pyroxene-andesite, dike southeast of Poker Flat. Contains plagioclase, augite, hornblende, magnetite, some glass, and occasional quartz. P. R. C. 736.

I. Quartz-bearing andesite, northwest of Downieville. Contains plagioclase, augite, enstatite, magnetite, occasional quartz, and probably glass. P. R. C. 753.

	F.	G.	H.	I.
SiO ₂	51.27	66.94	50.34	60.02
Al ₂ O ₃	12.14	16.49	17.61	16.07
Fe ₂ O ₃	2.51	1.41	3.63	2.17
FeO	6.71	1.87	2.28	3.46
MgO	10.88	1.98	3.50	4.57
CaO	10.32	4.77	6.45	7.01
Na ₂ O	2.00	3.88	3.40	3.55
K ₂ O	1.63	1.65	1.94	1.59
H ₂ O at 100°17	.35	.64	.24
H ₂ O above 100°	1.16	.22	.74	.45
TiO ₂60	.30	.32	.42
P ₂ O ₅21	.12	.25	.17
Cr ₂ O ₃				trace
NiO04			
MnO21	.13	.12	.10
SrO	trace?	.05	.04	trace
BaO07	.07	.11	.08
Li ₂ O	trace	trace	trace	none
SO ₃06
	99.92	100.23	100.37	99.96

6. NEVADA CITY AND GRASS VALLEY.

Rocks of a mining district in Nevada County, described by Lindgren in 17th Ann., Part II, p. 1.

A. Granodiorite, 1 mile southeast of Nevada City. Contains hornblende, biotite, quartz, plagioclase, orthoclase, magnetite, apatite, sphene, and pyrite. Analysis by W. F. Hillebrand, record No. 1478.

B. Granodiorite, Kate Hayes Hill, Grass Valley. Contains plagioclase, orthoclase, quartz, hornblende, pyrite, magnetite, apatite, sphene, and zircon. Analysis by Hillebrand, No. 1478.

C. Hornblende-porphyrity, Nevada City. Contains feldspar, hornblende, quartz, epidote, sericite, and biotite. Analysis by H. N. Stokes, record No. 1531.

D. Quartz-porphyrity, New Ophir claim, Grass Valley. Contains plagioclase, quartz, uralite, epidote, and augite, and hornblende altered into chlorite. Analysis by Stokes, No. 1531.

E. Diabase, near Maryland mine, Grass Valley. Contains feldspar, augite, hornblende, ilmenite, pyrrhotite, pyrite, and some chlorite. Analysis by Stokes, No. 1522.

F. Diabase, Grass Valley. Contains feldspar, pyroxene, hornblende, ilmenite, pyrrhotite, pyrite, and chlorite, and probably a little quartz. Analysis by Stokes, No. 1522.

	A.	B.	C.	D.	E.	F.
SiO ₂	66.65	63.85	62.09	63.39	51.01	53.19
Al ₂ O ₃	16.15	15.84	16.69	16.58	11.89	17.12
Fe ₂ O ₃	1.52	1.91	1.45	1.41	1.57	4.35
FeO.....	2.36	2.75	3.76	3.08	6.08	5.16
MgO.....	1.74	2.07	1.93	2.15	8.87	3.98
CaO.....	4.53	4.76	6.08	4.76	10.36	9.39
Na ₂ O.....	3.40	3.29	3.36	3.47	4.17	2.79
K ₂ O.....	2.65	3.08	1.84	2.79	.15	.28
H ₂ O at 110°...	.18	.28	.19	.22	.24	.17
H ₂ O above 110°	.72	1.65	1.47	1.87	2.09	1.21
TiO ₂38	.58	.32	.44	.98	1.34
P ₂ O ₅10	.13	.39	.14	.17	.13
Cr ₂ O ₃04	none
MnO.....	.10	.07	trace	trace	trace	trace
SrO.....	trace	trace				
BaO.....	.07	.06	.10	.11	none	trace
Li ₂ O.....	trace	trace				
SO ₃10			
FeS ₂02	.04			1.73	.94
CuS (?).....					trace	
	100.57	100.36	99.77	100.41	99.35	100.05

G. Wall rock, Federal Loan mine. A siliceous argillite, of sedimentary origin. Contains quartz, feldspar, biotite, pyrrhotite, and a little calcite. Analysis by Hillebrand, No. 1478.

H. Altered wall rock, Providence mine. Derived from granodiorite. Analysis by Hillebrand, No. 1478.

I. Altered wall rock, Providence mine, back vein. Derived from granodiorite and schist. Analysis by Hillebrand, No. 1478.

J. Altered wall rock, North Star mine. Derived from uralite-diorite. Contains quartz, sericite, calcite, pyrite, and sphene. Analysis by Hillebrand, No. 1478.

K. Altered country rock, Idaho mine. Derived from serpentine. Analysis by Hillebrand, No. 1478.

	G.	H.	I.	J.	K.
SiO ₂	73.63	60.26	59.76	45.74	36.19
Al ₂ O ₃	10.54	15.73	14.45	5.29	4.93
Fe ₂ O ₃	1.87	1.25	1.04	.13	.21
FeO.....		2.68	3.52	2.06	5.36
MgO.....	1.84	1.82	2.26	.94	22.94
CaO.....	2.47	5.44	6.09	23.85	4.60
Na ₂ O.....	1.81	1.92	1.12	.11	.16
K ₂ O.....	1.89	3.71	3.73	1.29	.06
H ₂ O at 110°.....	.11	.33	.26	.22	.18
H ₂ O above 110°.....	1.07	2.54	2.58	1.07	2.87
TiO ₂52	.42	.46	.36	.16
P ₂ O ₅13	.12	.16	.07	.05
MnO.....	trace?	.04	.09	.26	.12
NiO.....					.10
SrO.....	trace	trace	trace?	none	trace
BaO.....	.12	.07	.05	trace	trace
Li ₂ O.....	trace	trace	trace	trace	trace
CO ₂62	3.99	4.47	18.91	21.82
FeS ₂08	.24	.49	.22
Fe ₇ S ₈	3.16				
Organic C.....	.59				
	100.37	100.40	100.28	100.79	99.97

- L. Bleached country rock, next to vein, Osborne Hill mine. Derived from sandstone. Analysis by George Steiger, record No. 1541.
- M. Altered wall rock, Empire mine. Derived from granodiorite. Analysis by Steiger, No. 1541. Sp. gr., 2.782, 20°.
- N. Altered wall rock, Ebaugh tunnel. Derived from granodiorite. Mainly quartz and sericite, with pyrite, apatite, sphene, and carbonates. Analysis by Steiger, No. 1541. Sp. gr., 2.747, 20°.
- O. Altered wall rock, Federal Loan mine. Derived from siliceous argillite. Analysis by Steiger, No. 1541.

	L.	M.	N.	O.
SiO ₂	71.97	58.43	56.25	34.91
Al ₂ O ₃	15.75	17.40	17.65	15.55
Fe ₂ O ₃77	.77	.76	.17
FeO.....	.45	2.19	2.64	4.96
MgO.....	.80	1.50	1.69	4.58
CaO.....	.80	5.25	4.46	11.10
Na ₂ O.....	.33	1.76	.30	.19
K ₂ O.....	4.88	4.03	6.01	4.28
H ₂ O at 100°.....	.30	.30	.30	.30
H ₂ O above 100°.....	2.16	2.61	2.36	1.86
TiO ₂88	none	.25	1.65
P ₂ O ₅15	.13	.21	.82
MnO.....	none	none	none	none
BaO.....	trace	none	.03	none
SO ₃	trace	none	none	none
CO ₂38	4.04	4.82	15.57
FeS ₂56	1.59	2.87	4.20
	100.18	100.00	100.60	100.14

7. PLACER COUNTY.

First, a series of rocks from the Ophir mining district, described by Lindgren in 14th Ann., p. 249. Analyses by W. F. Hillebrand, record Nos. 1419, 1433, 1434.

A. Granodiorite, quarries at Lincoln, 8 miles west of Ophir. Contains feldspars, quartz, biotite, and hornblende.

B. Pyritiferous amphibolite, Conrad tunnel. Partly altered. Contains pyrite, hornblende, magnetite, feldspars, quartz, epidote, chlorite, a few scales of mica, rutile, and carbonates. Sp. gr. 2.901, 23°.

C. Dike rock, near camptonite, Casey's tunnel, Flat Ledge, Duncan Hill. Contains hornblende, feldspars, pyrite, and apatite, with secondary epidote and quartz.

D. Altered wall rock, Mina Rica vein. Sp. gr. 2.979, 20°.

E. Altered wall rock, Plantz vein. These rocks, D and E, contain quartz, muscovite, a little chlorite, pyrite, and sphene, with carbonates of calcium, magnesium, and iron.

	A.	B	C.	D.	E.
SiO ₂	65.54	45.56	60.09	37.01	46.13
Al ₂ O ₃	16.52	14.15	16.43	12.99	15.82
Fe ₂ O ₃	1.40	1.20	2.28	.43	.89
FeO.....	2.49	9.83	3.01	3.57	2.27
MgO.....	2.52	6.76	4.37	5.49	2.13
CaO.....	4.88	2.30	5.76	9.78	10.68
Na ₂ O.....	4.09	1.57	4.52	.13	.17
K ₂ O.....	1.95	1.18	.70	4.02	5.30
H ₂ O at 100°.....	.12	.23	.20	.13	.12
H ₂ O above 100°.....	.59	4.84	1.16	1.92	2.42
TiO ₂39	1.11	.63	.85	.67
P ₂ O ₅18	.14	.12	.06	.10
MnO.....	.06	.25	.12	.24	.09
Ni, Zn.....	-----	traces	-----	traces	traces
SrO.....	trace	trace	trace	trace	trace
BaO.....	trace	trace	trace	trace	trace
Li ₂ O.....	trace	trace	none	trace	trace
SO ₃	-----	.03	trace	.04	.04
CO ₂	-----	3.04	.07	15.04	11.24
FeS ₂	-----	7.86	.34	7.99	1.61
Cu ₂ S (?).....	-----	.10	-----	-----	-----
	100.73	100.15	99.80	99.69	99.68

Second, rocks from other localities in Placer County. Studied also by Lindgren, who furnishes the petrographic data. Analysis A by W. H. Melville, record No. 1346; B, C, D, and E by W. F. Hillebrand, record No. 1419.

A. Granite, Rocklin. A normal granite, containing quartz, orthoclase, plagioclase, biotite, muscovite, magnetite, apatite, and zircon, with some secondary chlorite and epidote derived from the biotite. Described by Lindgren in Bull. 150, p. 170. P. R. C. 66.

B. Granodiorite, Donner Pass. Contains plagioclase, orthoclase, quartz, hornblende, biotite, and sphene.

C. Gabbro, 2 miles south of Emigrant Gap, on road to Onion Valley. Contains biotite, hypersthene, diallage, plagioclase, and orthoclase.

D. Gabbro, same locality as C. Contains hypersthene, diallage, plagioclase, and orthoclase.

E. Augite-granite, southeast spur of English Mountain. Contains "basic" plagioclase, augite, and quartz.

	A.	B.	C.	D.	E.
SiO ₂	73.00	59.48	55.40	55.87	64.67
Al ₂ O ₃	16.38	17.25	15.32	13.52	16.62
Fe ₂ O ₃	none	2.15	2.70	2.70	.51
FeO.....	.99	4.06	5.49	5.89	.76
MgO.....	.48	2.67	5.75	6.51	2.26
CaO.....	2.42	6.50	9.90	8.87	9.50
Na ₂ O.....	4.53	3.53	2.89	2.42	4.10
K ₂ O.....	1.87	2.27	1.52	1.72	.34
H ₂ O at 100°.....	.52	.09	.03	.09	.08
H ₂ O above 100°.....		.71	.38	1.56	.37
TiO ₂93	.60	.56	.51
P ₂ O ₅33	.22	.25	.12
MnO.....		.11	.11	.10	trace
SrO.....		trace	none	none	trace
BaO.....		.09	.07	.02	.02
Li ₂ O.....		trace	trace	trace	trace
	100.19	100.17	100.38	100.08	99.86

8. ELDORADO COUNTY.

A. Granite, Placerville canal, one-third mile north of Ditch Camp No. 7. Collected by W. Lindgren, who reports it as containing biotite, orthoclase, plagioclase, and quartz. Analysis by George Steiger, record No. 1591.

B. Granodiorite, 2 miles south of Silver Lake Hotel. Collected by Lindgren, who reports it as containing hornblende, biotite, plagioclase, and quartz. Analysis by Steiger, No. 1591. Analyses A and B are published by Lindgren in Amer. Journ. Sci., 4th ser., vol. 3, p. 306.

C. Porphyrite, 1 mile southwest of Latrobe. Published by Turner in 17th Ann., Part I, p. 521. Contains abundant plagioclase, less augite, calcite or dolomite, iron disulphide, a little chlorite, and secondary greenish mica. Analysis by W. F. Hillebrand, record No. 1432. P. R. C. 721.

	A.	B.	C.
SiO ₂	77.68	67.45	68.58
Al ₂ O ₃	11.81	15.51	13.04
Fe ₂ O ₃72	1.76	.26
FeO.....	.51	2.21	3.40
MgO.....	.18	1.10	1.01
CaO.....	.72	3.60	3.22
Na ₂ O.....	2.96	3.47	4.94
K ₂ O.....	5.00	3.66	1.90
H ₂ O at 100°.....	.04	.14	.16
H ₂ O above 100°.....	.27	.63	1.00
TiO ₂14	.58	.57
P ₂ O ₅10	.12	.20
MnO.....	trace15
SrO.....	trace
BaO.....10
CO ₂	1.31
FeS ₂15
	100.13	100.23	99.99

9. AMADOR COUNTY.

Rocks collected by H. W. Turner, and analyses published in 14th Ann., p. 441, and 17th Ann., Part I, p. 521. Additional data supplied by Turner. Analyses by W. F. Hillebrand, record Nos. 1432, 1456, and 1597.

A. Rhyolite, south point of Buena Vista Peak. Contains sanidine, quartz, and biotite in a glassy groundmass. P. R. C. 729.

B. Quartz-monzonite, north fork of Mokelumne River. Contains plagioclase, microcline, quartz, abundant biotite, iron ore, sphene, apatite, and perhaps rutile. P. R. C. 770.

C. Quartz-monzonite, north fork of the Mokelumne River. Like B. P. R. C. 765.

D. Quartz-porphyrity-schist, $2\frac{1}{2}$ miles southeast of Buena Vista Peak. Contains porphyritic quartz and hornblende, also calcite and other carbonates. See 14th Ann. P. R. C. 723.

E. Quartz-diorite-gneiss, north fork of Mokelumne River. Contains plagioclase, hornblende, quartz, brown mica, accessory biotite, and iron oxide. P. R. C. 764.

F. Diorite-porphry, north fork of Mokelumne River. Contains plagioclase, brown hornblende, epidote, and a little sulphide of iron and chlorite. P. R. C. 769.

	A.	B.	C.	D.	E.	F.
SiO ₂	73.23	70.75	70.43	70.29	57.41	55.18
Al ₂ O ₃	12.73	15.13	15.51	11.83	17.71	17.35
Fe ₂ O ₃99	.98	.96	1.30	2.16	2.77
FeO.....	.16	1.43	1.28	2.08	5.01	3.90
MgO.....	.22	.73	.37	1.24	3.38	4.80
CaO.....	.61	3.09	2.76	2.30	6.73	7.98
Na ₂ O.....	1.91	3.05	2.75	2.68	3.12	3.42
K ₂ O.....	5.17	3.62	5.14	3.05	1.82	1.42
H ₂ O at 100°...	.53	.10	.08	.10	.20	.16
H ₂ O above 100°	4.51	.51	.40	1.35	1.14	1.52
TiO ₂09	.42	.24	.29	1.04	.83
P ₂ O ₅02	.10	.11	.07	.24	.20
NiO.....	none	(?)02	.03
MnO.....	trace	trace	trace	.12	.15	.15
SrO.....	none	.04	.05	trace?	.04	.06
BaO.....	.02	.12	.20	.07	.09	.04
Li ₂ O.....	trace	trace	trace	none	trace	trace
CO ₂	none	none	3.25	none	none
FeS ₂06	trace	none	.28
	100.19	100.13	100.28	100.02	100.26	100.09

G. Diorite, north fork of Mokelumne River. Contains quartz, feldspar, biotite, sphene, epidote, and secondary chlorite. P. R. C. 771.

H. Diorite, north fork of Mokelumne River. Contains plagioclase, quartz, hornblende, biotite, apatite, iron ore, epidote, and chlorite, P. R. C. 772.

I. Plagioclase-gneiss, north fork of Mokelumne River. Contains plagioclase, hornblende, biotite, and apatite. P. R. C. 768.

J. Plagioclase-gneiss, north fork of Mokelumne River. Contains plagioclase, hornblende, brown mica, apatite, epidote, and grains of iron ore. P. R. C. 767.

	G.	H.	I.	J.
SiO ₂	69.66	55.86	52.21	46.63
Al ₂ O ₃	17.57	19.30	18.79	19.47
Fe ₂ O ₃21	.91	2.71	3.26
FeO	1.04	4.78	5.30	6.63
MgO58	2.94	5.11	5.37
CaO	4.54	7.31	8.01	9.15
Na ₂ O	4.91	3.52	3.31	3.19
K ₂ O71	1.52	1.60	1.55
H ₂ O at 110°05	.19	.12	.10
H ₂ O above 110°50	1.23	1.35	1.61
TiO ₂21	1.20	1.16	1.82
P ₂ O ₅03	.38	.36	.66
V ₂ O ₃02
NiO	none	trace	trace	.02
MnO	trace	.16	.06	.21
SrO05	.04		.06
BaO03	.13	.08	.14
Li ₂ O	none	trace	trace	trace
CO ₂	none	none	none	none
FeS ₂	trace?	.39	.06	.19
	100.09	99.86	100.23	100.08

K. Wollastonite-gneiss, north fork of Mokelumne River. Mainly wollastonite, but garnet, quartz, and sphene are also present. P. R. C. 766.

L. Melaphyr-tuff, altered basalt, west of Jackson. Contains augite and plagioclase, with secondary quartz, chlorite, and chrysotile. Originally glassy in part, but devitrified. See 14th Ann. P. R. C. 722.

M. Reddish-brown mica separated from pyroxenic-gneiss, north fork of Mokelumne River, about 1 kilometer above mouth of Bear River. Described by Turner in Amer. Journ. Sci., 4th series, vol. 7. p. 294. Analysis by William Valentine, record No. 1736.

	K.	L.	M.
SiO ₂	50.67	49.24	36.62
Al ₂ O ₃	6.37	14.79	14.37
Fe ₂ O ₃31	1.36	4.04
FeO.....	.50	8.00	17.09
MgO.....	.58	6.89	9.68
CaO.....	40.34	10.74	1.48
Na ₂ O.....	.14	2.76	.45
K ₂ O.....	.22	.88	8.20
H ₂ O at 110°.....	.08	.20	.90
H ₂ O above 110°.....	.31	2.97	3.26
TiO ₂20	.96	3.03
P ₂ O ₅	none	.17	none
NiO.....	none		
MnO.....	trace	.18	.40
SrO.....	none	trace	trace
BaO.....	none	.04	.33
Li ₂ O.....	none	trace	trace
F.....			.10
CO ₂52	.90	
	100.24	100.08	99.95
Less O.....			.04
			99.91

10. CALAVERAS COUNTY.

Rocks collected by H. W. Turner, and described in 14th Ann., p. 441. Additional data supplied by Turner relative to analysis B. Analyses by W. F. Hillebrand, record No. 1432.

A. Meta-dacite, $1\frac{3}{4}$ miles southeast of Milton. Contains quartz, feldspar, and hornblende. P. R. C. 777.

B. Meta-dacite, $1\frac{1}{2}$ miles northeast of Milton. Contains feldspar, quartz, epidote, chlorite, and iron ore, in a groundmass made up probably of feldspar and quartz. P. R. C. 752.

C. Meta-andesite, $1\frac{1}{2}$ miles northward from Jenny Lind. Contains quartz, plagioclase, epidote, and chlorite derived from augite.

	A.	B.	C.
SiO ₂	72.24	71.19	61.37
Al ₂ O ₃	13.84	13.81	15.41
Fe ₂ O ₃	1.45	1.45	3.15
FeO.....	1.86	1.68	3.89
MgO.....	1.10	.74	3.48
CaO.....	3.40	2.87	4.42
Na ₂ O.....	4.43	4.24	3.76
K ₂ O.....	.39	1.82	.34
H ₂ O at 100°.....	.17	.15	.29
H ₂ O above 100°.....	.69	.92	2.70
TiO ₂41	.35	.60
P ₂ O ₅10	.08	.08
MnO.....	.12	.07	.47
SrO.....	trace	trace	trace
BaO.....	.08	.16	.08
CO ₂82	
	100.28	100.35	100.04

11. TUOLUMNE COUNTY.

Rocks collected by H. W. Turner, and partly described in his papers in the 14th and 17th Annuals. The latites were named and described by Ransome in Bull. 89. Some additional data have been furnished by Turner.

A. Soda-syenite-porphyry, dike east of Moccasin Creek. Consists mainly of albite, with a greenish mineral which is probably ægirite. Analysis by H. N. Stokes, record No. 1563. P. R. C. 773.

B. Augite-syenite, dike on Turnback Creek, about 1 mile north of Carter post-office. Hitherto unpublished. Contains orthoclase and augite, with less plagioclase and quartz. Analysis by Stokes, No. 1642.

C. Diorite, dike $1\frac{1}{2}$ miles southeasterly from Sonora. Contains feldspar largely altered to hornblende. A few black grains are probably iron ore. Analysis by W. F. Hillebrand, record No. 1548. P. R. C. 759.

D. Quartz-pyroxene-diorite, large area east of Sonora. Contains plagioclase, quartz, biotite, augite, rhombic pyroxene, and a trace of iron ore. Analysis by Hillebrand, No. 1548. P. R. C. 760.

E. Diorite, dike about $1\frac{1}{2}$ miles southeasterly from Sonora. Contains altered plagioclase and hornblende, with epidote, chlorite, and iron disulphide as secondary products. Analysis by Hillebrand, No. 1548. P. R. C. 761.

	A.	B.	C.	D.	E.
SiO ₂	67.53	61.28	58.05	57.80	53.46
Al ₂ O ₃	18.57	14.71	15.46	16.43	14.81
Fe ₂ O ₃	1.13	1.21	1.69	1.62	2.60
FeO08	2.85	5.09	6.51	5.15
MgO24	1.69	4.84	4.14	7.27
CaO55	5.61	6.94	7.21	8.44
Na ₂ O	11.50	2.99	2.86	2.35	2.64
K ₂ O10	7.70	2.14	2.29	1.30
H ₂ O at 110°15	.28	.10	.11	.12
H ₂ O above 110°31	.43	2.02	.38	2.13
TiO ₂07	.41	.72	.70	.70
P ₂ O ₅11	.16	.16	.19	.16
MnO	trace	trace	.14	.18	.18
NiO			none	.03	.05
SrO	trace	.04	trace	trace?	trace
BaO72	.07	.09	.05
Li ₂ O			trace	trace	trace
SO ₃	trace	.08			
CO ₂			none	none	.44
F	trace				
FeS ₂			none	none	.26
	100.34	100.16	100.28	100.03	99.76

F. Biotite-augite-latite, 4 miles southwest of Clover Meadow. Called "trachyte-andesite-tuff" in former edition of this bulletin. Contains plagioclase, biotite, augite, magnetite, apatite, and glass. Analysis by W. F. Hillebrand, record No. 1597. P. R. C. 762.

G. Augite-latite, Dardanelle flow, near Clover Meadow. Called "trachyte-andesite" in former edition. Contains plagioclase, in part labradorite, augite, iron ore, some olivine, apatite, and brown glass. The potassium is probably in the glass, as no potash mineral was observed. Analysis by H. N. Stokes, record No. 1645. P. R. C. 785.

H. Augite-latite, Table Mountain. Called "basalt" in former edition. Contains labradorite, olivine, augite, and magnetite. Analysis by Hillebrand, record No. 1273. P. R. C. 724.

I. Augite-latite, Table Mountain, near Clover Meadow. Contains labradorite, augite, olivine, magnetite, apatite, and glass. Analysis by George Steiger, record No. 1697.

	F.	G.	H.	I.
SiO ₂	62.33	59.43	56.19	56.78
Al ₂ O ₃	17.30	16.68	16.76	16.86
Fe ₂ O ₃	3.00	2.54	3.05	3.56
FeO.....	1.63	3.48	4.18	2.93
MgO.....	1.05	1.84	3.79	3.41
CaO.....	3.23	4.09	6.53	6.57
Na ₂ O.....	4.21	3.72	2.53	3.19
K ₂ O.....	4.46	5.04	4.46	3.48
H ₂ O at 110°.....	.44	.27	.34	.15
H ₂ O above 110°.....	.75	.72	.66	1.21
TiO ₂	1.05	1.38	.69	1.15
P ₂ O ₅29	.58	.55	.42
ZrO ₂04	.08
V ₂ O ₃01
MnO.....	.08	trace	.10	none
SrO.....	.05	trace	trace
BaO.....	.24	.14	.19	trace
Li ₂ O.....	trace	none	trace
CO ₂18
Cl.....05
F.....	trace
C.....	.11
FeS ₂06
	100.33	100.04	100.02	99.89

J. Amphibole-gabbro, Beaver Creek, Big Trees quadrangle. Contains labradorite and amphibole, with a little pyrite and pyrrhotite. Analysis by H. N. Stokes, record No. 1752.

K. Amphibole separated from J. Analysis by William Valentine, record No. 1733.

L. Olivine-gabbro, just east of south end of Phoenix reservoir. Contains plagioclase, a few grains of alkali feldspar, augite, rhombic pyroxene, amphibole, olivine, magnetite, and iron sulphide, with a little secondary chlorite and epidote. Analysis by H. N. Stokes, record No. 1750.

For description of J and K see Turner, Amer. Jour. Sci., 4th series, vol. 7. p. 294. The description of L is hitherto unpublished.

	J.	K.	L.
SiO ₂	47.27	46.08	43.41
Al ₂ O ₃	20.82	10.52	23.15
Fe ₂ O ₃	1.85	2.81	3.72
FeO	4.26	8.30	4.39
MgO	6.44	14.40	7.65
CaO	13.02	12.64	14.27
Na ₂ O	2.75	1.62	.82
K ₂ O22	.34	.22
H ₂ O at 110°08	.17	.18
H ₂ O above 110°	1.27	1.97	1.53
TiO ₂92	.77	.39
P ₂ O ₅74	.18	.02
V ₂ O ₃02	.04	-----
Cr ₂ O ₃	trace	-----	none
MnO	trace	.15	.08
SrO	trace	-----	-----
Li ₂ O	none	none	trace
CO ₂	-----	-----	.10
FeS ₂20	-----	.14
Cl	trace	-----	trace
	99.86	99.99	100.07

12. MARIPOSA COUNTY.

Rocks collected by H. W. Turner, and partly described in his papers in the 14th and 17th Annuals. Additional data supplied by Turner.

A. Soda-granulite or aplite, about 4 miles west of Mariposa. See 7th Ann., Part I, p. 721. Contains plagioclase (albite?) and micropegmatite, with less epidote, quartz, sphene, and apatite. Analyses by W. F. Hillebrand, record No. 1461. P. R. C. 748.

B. Micropegmatite, Agua Fria Creek. See 17th Ann., Part I, p. 91. Contains quartz, plagioclase, brown mica, epidote, and a little iron ore. Analysis by Hillebrand, No. 1461. P. R. C. 746.

C. Soda-granite-porphry, Merced River, below the mouth of the north fork. Published in 17th Ann. Contains feldspar, largely albite, hornblende, muscovite, abundant epidote, apatite, and a little iron ore. Analysis by George Steiger, record No. 1573. P. R. C. 774.

D. Granite-porphry, about one-fourth of a mile north of Lake Tenaya, Yosemite National Park. See 14th and 17th Annuals. Contains orthoclase, quartz, plagioclase, and biotite, with a little iron ore and sphene. Analysis by Hillebrand, record No. 1432. P. R. C. 726.

E. Granite, west of Lake Tenaya, Yosemite National Park. See 14th and 17th Annuals. Contains quartz, orthoclase, plagioclase, and biotite, with some hornblende, iron ore, sphene, and apatite. Analysis by Hillebrand, No. 1432. P. R. C. 725.

	A.	B.	C.	D.	E.
SiO ₂	74.21	73.18	71.88	72.48	66.28
Al ₂ O ₃	14.47	13.66	15.57	14.06	16.03
Fe ₂ O ₃35	.21	1.07	.89	1.80
FeO.....	.50	2.24	.30	1.05	1.88
MgO.....	.28	.93	.68	.62	1.12
CaO.....	1.71	2.10	2.03	2.17	3.75
Na ₂ O.....	7.62	3.70	5.81	3.30	4.10
K ₂ O.....	.10	2.72	1.80	4.75	3.49
H ₂ O at 110°.....	.15	.10	.11	.16	.10
H ₂ O above 110°.....	.23	.57	.68	.35	.39
TiO ₂30	.25	.17	.28	.54
P ₂ O ₅07	.09	.08	.09	.30
MnO.....	none	.07	none	trace	.05
SrO.....	trace	trace	.08	trace	trace
BaO.....	none	.10	.02	.08	.08
Li ₂ O.....	trace	trace	none	trace	trace
CO ₂17	none		
	99.99	100.09	100.28	100.28	99.91

F. Amphibole-biotite granite, Nevada Falls trail, Yosemite Valley. Contains alkali feldspar, plagioclase, quartz, amphibole, biotite, magnetite, and apatite.

G. Biotite-granite, base of El Capitan, Yosemite Valley. Contains alkali feldspar, plagioclase, quartz, biotite, titanite, apatite, and iron oxides.

H. Brown mica separated from G.

Analyses F, G, and H by William Valentine, record Nos. 1732, 1733.

I. Amphibole separated from quartz-monzonite, Tioga road, south-east of Mount Hoffman. Sp. gr. 3.203, 2.15°.

J. Mica separated from the same rock as I. Sp. gr. 3.05, 21°.

Analyses I and J by W. F. Hillebrand, record No. 1774.

Samples G to J are described by Turner in Amer. Journ. Sci., 4th series, vol. 7, p. 294.

	F.	G.	H.	I.	J.
SiO ₂	66.83	71.08	35.64	47.49	35.75
Al ₂ O ₃	15.24	15.90	18.62	7.07	14.70
Fe ₂ O ₃	2.73	.62	5.54	4.88	4.65
FeO.....	1.66	1.31	14.60	10.69	14.08
MgO.....	1.63	.54	9.72	13.06	12.37
CaO.....	3.59	2.60	.90	11.92	.17
Na ₂ O.....	3.10	3.54	.38	.75	.32
K ₂ O.....	4.46	4.08	9.22	.49	9.19
H ₂ O at 110°.....	none	none	.48	1.03
H ₂ O above 110°.....	.56	.30	2.54	1.86	3.64
TiO ₂54	.22	1.12	1.21	3.16
P ₂ O ₅18	.10	.20	none	.03
ZrO ₂04	.08
V ₂ O ₃04	.05
Cr ₂ O ₃	none	trace
MnO.....	.10	.15	.79	.51	.45
NiO, CoO.....02	.02
SrO.....	.03	.02	none	(?)
BaO.....	.11	.04	trace	none	.12
Li ₂ O.....	trace	trace	trace	trace
CO ₂	trace	trace
Cl.....	.02	.02
F.....26	.06	.17
Less O.....	100.82	100.60	100.01	100.05	99.90
			.11	.02	.07
			99.90	100.03	99.83

K. Quartz-mica-diorite, Chowchilla River. See 17th Ann., Part I, p. 691. Contains plagioclase, quartz, a little orthoclase (?), brown mica, hornblende, rather abundant apatite, a little iron ore; one zircon-like crystal was noted. Analysis by W. F. Hillebrand, record No. 1461. P. R. C. 745.

L. Quartz-mica-diorite, Yaqui Creek. For the Educational Series of Rocks. Contains plagioclase, quartz, biotite, hornblende, a little pyroxene, iron ore, and apatite. Analysis by George Steiger, record No. 1643. P. R. C. 135. Described by Turner in Bull. 150, p. 339.

M. Diabase, dike $1\frac{1}{2}$ miles northeast of Hornitos. See 17th Ann., Part I, p. 694. Contains plagioclase, partly labradorite, augite, brown hornblende, and iron ore. Analysis by Hillebrand, No. 1461. P. R. C. 750.

N. Igneous rock, near Cathay Hill. See 17th Ann., Part I, p. 694. Contains two minerals unidentified; neither is olivine. Analysis by Hillebrand, No. 1461. P. R. C. 749.

	K.	L.	M.	N.
SiO ₂	62.62	58.09	51.32	47.75
Al ₂ O ₃	17.51	17.46	15.28	10.56
Fe ₂ O ₃49	1.12	.47	.74
FeO.....	4.06	5.08	8.59	8.34
MgO.....	2.84	4.06	7.25	19.09
CaO.....	5.49	6.24	11.58	9.62
Na ₂ O.....	3.49	2.94	2.92	1.32
K ₂ O.....	1.76	2.02	.22	.12
H ₂ O at 110°.....	.22	.29	.06	.05
H ₂ O above 110°.....	.92	1.45	.95	2.06
TiO ₂55	.95	1.23	.37
P ₂ O ₅12	.17	.25	.03
Cr ₂ O ₃24
NiO.....				.07
MnO.....	.05	none	.16	.10
SrO.....	trace	.04	trace	trace
BaO.....	trace	.07	none	none
Li ₂ O.....	trace	none	trace	trace
CO ₂21		
SO ₃05		
Cl.....		.02		
F.....		trace		
C.....		.11		
	100.12	100.37	100.28	100.46

O. Feldspathic mica-schist, Chowchilla River. See 17th Ann., Part I, p. 691. Contains quartz, feldspar, biotite, muscovite, apatite, and specular iron. Analysis by W. F. Hillebrand, record No. 1461. P. R. C. 744.

P. Andalusite-hornfels, Yaqui Gulch. Principally quartz, andalusite, brown and white mica, black graphite-like grains, a little iron ore, and probably feldspar. Analysis by George Steiger, record No. 1643. Described by Turner in Bull. 150, p. 342. P. R. C. 135.

Q. Andalusite-schist, Chowchilla River. See 17th Ann., Part I, p. 691. Contains quartz, biotite, andalusite, sericite, a little muscovite, probably graphite, iron ore, a few garnets, and apparently chlorite. Analysis by Hillebrand, No. 1461. P. R. C. 743.

R. Chiastolite-schist, Yaqui Gulch. Contains chiastolite, sillimanite, brown mica, sericite (?), probably graphite, and clear grains which appear to be quartz and feldspar. Analysis by Steiger, No. 1643. Described by Turner in Bull. 150, p. 342. P. R. C. 135.

S. Hornfels, Agua Fria Creek. See 17th Ann., Part I, p. 691. Contains quartz, brown mica, iron ore, and plagioclase. Analysis by Hillebrand, No. 1461. P. R. C. 747.

	O.	P.	Q.	R.	S.
SiO ₂	70.40	65.10	64.28	62.15	68.27
Al ₂ O ₃	14.70	17.77	17.28	19.34	14.03
Fe ₂ O ₃65	1.95	1.10	4.23	.46
FeO.....	2.57	3.29	5.34	2.25	4.68
MgO.....	1.47	1.43	2.57	1.88	2.23
CaO.....	1.63	1.38	1.19	1.50	3.89
Na ₂ O.....	3.17	2.25	.91	1.60	2.29
K ₂ O.....	3.46	2.45	2.93	3.07	3.35
H ₂ O at 110°.....	.19	.47	.20	.19	.08
H ₂ O above 110°.....	.91	2.49	2.72	1.79	.98
TiO ₂51	.72	.65	.80	.57
P ₂ O ₅05	.14	.27	.15	.21
MnO.....	.08	none	.09	trace	.04
SrO.....	trace	none	trace	none	trace
BaO.....	.09	none	.10	.04	.08
Li ₂ O.....	trace	none	trace	none	trace
SO ₃0313
Cl.....	trace	none
F.....1222
C.....	.15	1.21	.43	1.12
	100.03	100.80	100.06	100.55	100.16
Less O.....0610
		100.74	100.45	

13. THE QUICKSILVER REGION.

Rocks described by Becker in Mon. XIII. Analyses made by W. H. Melville in the San Francisco laboratory. With one exception (the serpentine from New Idria) all the rocks are from the districts north of San Francisco.

A. Pseudo-diabase, near Mount St. Helena. Contains augite, hornblende, oligoclase, albite, zoisite, ilmenite, leucoxene, and a little chlorite.

B. Pseudo-diabase, Sulphur Bank. Contains oligoclase, a little quartz, pyroxene, hornblende, ilmenite, sphene, serpentine, and chlorite.

C. Pseudo-diorite, Knoxville. Mainly actinolite, with a little white mica, chlorite, serpentine, sphene, rutile, and zircon.

D. Glaucophane-schist, Sulphur Bank. Mainly glaucophane and zoisite. Quartz, albite, muscovite, and sphene are also present.

	A.	B.	C.	D.
SiO ₂	49.08	51.28	50.44	49.68
Al ₂ O ₃	14.68	15.05	8.18	13.60
Fe ₂ O ₃	1.95	2.42	1.06	1.86
FeO.....	9.63	8.01	6.29	8.61
MgO.....	6.69	6.07	17.63	6.26
CaO.....	10.09	7.08	11.55	10.97
Na ₂ O.....	4.60	4.43	2.98	3.09
K ₂ O.....	.20	.12	.50	.12
H ₂ O at 100°.....	.27	.39	.07	-----
H ₂ O above 100°.....	1.18	2.96	.92	3.84
TiO ₂	1.72	1.33	-----	1.31
P ₂ O ₅23	.13	-----	.21
Cr ₂ O ₃	-----	-----	.48	-----
MnO.....	.15	.25	.21	.04
NiO.....	-----	.10	-----	-----
	100.47	99.62	100.31	99.59

E. Andesitic obsidian, Clear Lake. Shows grains of plagioclase, augite, and hypersthene. Sp. gr. 2.391.

F. Andesite (asperite), Clear Lake. Contains pyroxene, plagioclase, magnetite, and sometimes biotite. Pyroxene mostly rhombic. Sp. gr. 2.664.

G. Obsidian, south of Borax Lake. Sp. gr. 2.390.

H. Basalt, south of Burns Valley. Rich in olivine, with a microlitic groundmass of plagioclase and augite. Sp. gr. 2.380.

I. Ordinary basalt, Knoxville.

	E.	F.	G.	H.	I.
SiO ₂	74.01	65.43	75.40	57.37	51.66
Al ₂ O ₃	12.95	17.10	7.72	15.66	11.22
Fe ₂ O ₃		2.39	1.41	2.06	} 7.62
FeO.....	1.42	1.19		4.46	
MgO.....	.48	1.48	1.26	8.84	13.61
CaO.....	.99	3.88	1.55	4.94	7.72
Na ₂ O.....	5.34	3.66	8.09	3.05	5.98
K ₂ O.....	4.65	2.83	4.52	1.51	.89
H ₂ O at 100°.....		.20	} .43	.61	} 1.06
H ₂ O above 100°.....	.29	.36		.12	
TiO ₂24	.83		.60	trace
P ₂ O ₅01	trace		.02	
Cr ₂ O ₃25
MnO.....	trace	.70	.12	.27	.12
NiO.....		.20		.41	
Cl.....	.07		.12		
	100.45	100.25	100.62	99.92	100.13

J. Light-green, marmolitic serpentine, New Idria.

K. Black serpentine, Sulphur Bank.

L. Light-green serpentine, Sulphur Bank.

	J.	K.	L.
SiO ₂	41.54	39.64	41.86
Al ₂ O ₃	2.48	1.30	.69
FeO.....	1.37	7.76	4.15
MgO.....	40.42	37.13	38.63
H ₂ O.....	14.18	13.81	14.16
Cr ₂ O ₃29	.24
MnO.....		.12	.20
NiO.....	.04	.33	trace
	100.03	100.38	99.93

14. MOUNT DIABLO.

Described by Turner and Melville in Bull. Geol. Soc. Amer., vol. 2, pp. 383-414. Analyses by W. H. Melville. Those with record numbers were made in the Washington laboratory; the others were made in the laboratory at San Francisco.

A. Diabase, Mitchell Canyon. Composed of augite, plagioclase, and ilmenite, with uraltite and chlorite secondary.

B. The same rock, partly altered and partly uraltitic. Called "diabase-diorite" by Turner.

C. Pyroxenite, near Bagley Creek. Composed of bronzite and diallage. Equivalent to the websterite of North Carolina. P. R. C., 735.

D. Glaucophane-schist, Pine Canyon. Contains numerous cinnamon garnets.

A, B, and C have the record No. 1247.

	A.	B.	C.	D.
SiO ₂	52.06	51.58	53.25	47.84
Al ₂ O ₃	14.34	14.99	2.80	16.88
Fe ₂ O ₃	2.11	2.04	.69	4.99
FeO.....	7.74	8.36	5.93	5.56
MgO.....	9.26	6.51	19.91	7.89
CaO.....	8.05	8.59	16.22	11.15
Na ₂ O.....	1.74	3.08	.19	3.20
K ₂ O.....	.73	.31	trace	.46
H ₂ O at 105°.....	.59	.34	.05	.17
H ₂ O above 105°.....	2.90	2.67	.24	1.81
TiO ₂47	1.05	-----	-----
P ₂ O ₅13	.24	-----	.14
Cr ₂ O ₃	-----	-----	.54	-----
NiO.....	-----	-----	.07	-----
MnO.....	trace	trace	.09	.56
	100.12	99.76	99.98	100.65

E. Crystalline gabbro, Bagley Creek. Contains plagioclase and diallage. Record No. 1166.

F. Shaly gabbro. Friable, containing carbonates and sulphates. Somewhat resembles serpentine.

G. Shaly gabbro, like F, much resembling a true serpentine.

According to Turner, F and G are merely weathered layers of the gabbro.

	E.	F.	G.
SiO ₂	47.49	45.43	45.69
Al ₂ O ₃	15.81	12.55	13.30
Fe ₂ O ₃	1.07	-----	1.85
FeO.....	4.50	6.50	4.72
MgO.....	10.39	13.41	13.06
CaO.....	15.53	12.39	13.50
Na ₂ O.....	1.16	1.71	1.36
K ₂ O.....	trace	.11	trace
H ₂ O at 105°.....	1.20	2.41	2.29
H ₂ O above 105°.....	1.83	2.74	2.47
P ₂ O ₅	trace	.04	.06
NiO.....	.06	-----	-----
MnO.....	.41	.21	.24
CO ₂	-----	2.35	1.89
SO ₃	-----	.24	.43
Organic matter.....	-----	trace	-----
	99.45	100.09	100.86

The following analyses are of serpentines derived from a peridotite-pyroxenite dike:

H. Almost black. Possibly derived from adjacent shale.

I. Almost black. Derived from pyroxenite.

J. Bastite. Derived from pyroxenite. Record No. 1166.

K. Friable serpentine.

	H.	I.	J.	K.
SiO ₂	38.53	40.50	36.57	36.96
Al ₂ O ₃	14.55	.78	.95	.39
Fe ₂ O ₃	2.65	4.01	7.29	5.00
FeO.....	4.01	2.04	.37	2.34
MgO.....	21.79	37.43	40.27	33.84
CaO.....	3.13	.39	.14	3.81
Na ₂ O.....	.07	.28	.31	.34
K ₂ O.....	.88	.16	trace	.14
H ₂ O at 100°.....	4.51	2.81	.94	2.16
H ₂ O above 100°.....	9.56	10.94	12.43	14.02
P ₂ O ₅	trace	trace	-----	.02
Cr ₂ O ₃	trace	.41	.33	.78
NiO.....	trace	.11	.31	trace
MnO.....	.32	.13	.10	.09
	100.00	99.99	100.01	99.89

Accidental organic matter was deducted from analyses H and I, with subsequent recalculation of the data to 100 per cent.

L. Serpentine.

M. Talc-like, yellowish-green serpentine.

N. Weathered serpentine.

O. Olive colored, probably serpentine.

Samples H, I, and J are from near Bagley Creek; K, L, M, and N from near Arroyo del Cerro; O from Ferguson ravine. M and O carry considerable chromite.

	L.	M.	N.	O.
SiO ₂	34.84	32.27	41.52	30.98
Al ₂ O ₃42	11.45	1.57	1.04
Fe ₂ O ₃	6.08	trace	3.50	4.88
FeO	1.85	5.05	1.07	2.01
MgO	30.74	33.30	36.84	38.44
CaO	7.02	.41	.44	.22
Na ₂ O42	trace	-----	.40
K ₂ O07	trace	-----	.16
H ₂ O at 100°.....	1.67	.44	3.32	.39
H ₂ O above 100°	15.72	12.40	12.51	20.43
P ₂ O ₅04	trace	-----	trace
Cr ₂ O ₃68	5.19	-----	.34
NiO	trace	.19	-----	-----
MnO01	trace	.29	.42
SO ₃	-----	-----	-----	.44
	99.56	100.70	101.06	100.15

15. MISCELLANEOUS ROCKS.

A. Rhyolitic obsidian, Medicine Lake, Modoc County. Collected by J. S. Diller, who reports it to be a banded obsidian, containing a few minute feldspar crystals. The banding is due to a multitude of trichites. Analysis by L. G. Eakins, record No. 1072.

B. Tuff from Hyampom, south fork of Trinity River, Trinity County. Almost wholly made up of particles of clear glass. Collected and described by Diller. Analysis by George Steiger, record No. 1427.

C. Diabase-porphyrte, one-half mile west of Browns Valley, Yuba County. Collected and described by W. Lindgren. Contains augite, plagioclase, magnetite, chlorite, and epidote. Analysis by W. F. Hillebrand, record No. 1419.

D. Amphibolitic schist, 1 mile northeast of Browns Valley, Yuba County. Collected and described by W. Lindgren. From metamorphosis of C. Contains chiefly green hornblende and feldspar. Analysis by Hillebrand, No. 1419.

	A.	B.	C.	D.
SiO ₂	73.51	70.40	48.26	54.13
Al ₂ O ₃	14.42	13.50	14.83	14.53
Fe ₂ O ₃46	1.31	3.27	1.50
FeO.....	1.49	1.61	5.97	5.25
MgO.....	.33	.37	8.77	10.93
CaO.....	1.26	.56	11.38	4.91
Na ₂ O.....	4.03	2.11	1.57	3.53
K ₂ O.....	4.29	2.39	1.13	.32
H ₂ O at 100°.....	.40	1.05	.10	.20
H ₂ O above 100°.....		7.41	3.37	4.01
TiO ₂51	.46
P ₂ O ₅04	.08	.25	.09
MnO.....	trace	trace	.15	.15
SrO.....			trace	trace
BaO.....			.05	.02
Li ₂ O.....			trace	none
CO ₂			1.24	-----
	100.23	100.79	100.85	100.03

E. Basalt, base of a lava flow, east of the head of the San Joaquin River, Madera County. Description supplied by Turner. Hitherto unpublished. Contains pyroxene, partly augite, plagioclase, olivine, and iron ores. Analysis by W. F. Hillebrand, record No. 1767.

F. Olivine-basalt, west peak of the Dardanelles, Alpine County. Described by Ransome in Bull. 89. Contains olivine, largely altered to iddingsite, plagioclase, serpentine, and augite. Analysis by George Steiger, record No. 1697.

G. Mica separated from quartz-monzonite, near Bloods Station, Alpine County. Described by Turner in Amer. Journ. Sci., 4th series, vol. 7, p. 294. Analysis by William Valentine, record No. 1736.

	E.	F.	G.
SiO ₂	51.89	48.76	35.62
Al ₂ O ₃	15.28	16.60	15.24
Fe ₂ O ₃	3.10	5.60	4.69
FeO.....	3.60	5.01	13.67
MgO.....	8.68	6.93	12.70
CaO.....	7.38	8.79	.95
Na ₂ O.....	3.27	2.47	.50
K ₂ O.....	2.57	.66	7.72
H ₂ O, at 105°.....	1.17	1.49	.94
H ₂ O, above 105°.....	1.37	2.19	4.36
TiO ₂91	1.26	2.61
P ₂ O ₅61	.19
ZrO ₂	trace
MnO.....	.12	none
NiO.....	.02
SrO.....	.09	trace
BaO.....	.15	trace	.26
Li ₂ O.....	trace	trace
CO ₂	none	.42
	100.21	100.37	100.00

H. Scoriaceous rhyolite, Mono Craters, south of Mono Lake. Described by Russell in 8th Ann., Part I, p. 380. Analysis by T. M. Chatard, record No. 36.

I. Obsidian, Mono Lake. Analysis by W. H. Melville, record No. 1346. Described by Lindgren in Bull. 150, p. 149. P. R. C. 60.

J. Punice, Mono Lake. Analysis by Melville, No. 1346. Described by Lindgren in Bull. 150, p. 148. P. R. C. 59.

K. Volcanic ash, east shore of Owens Lake. Collected and analyzed by Chatard, record No. 783.

	H.	I.	J.	K.
SiO ₂	74.05	75.78	67.39	55.81
Al ₂ O ₃	13.85	12.39	15.99	10.07
Fe ₂ O ₃	trace	.22	.56	3.43
FeO.....		1.25	1.99	.67
MgO.....	.07	.31	.77	2.22
CaO.....	.90	.81	1.63	1.05
Na ₂ O.....	4.60	4.00	4.74	.75
K ₂ O.....	4.31	4.64	4.80	2.98
H ₂ O at 110°.....	} 2.20	} .41	} 2.06	.65
H ₂ O above 110°.....				2.50
TiO ₂80
P ₂ O ₅27
MnO.....				.23
CaCO ₃				14.44
NaCl, soluble.....				1.45
Na ₂ SO ₄ , soluble.....				.80
Na ₂ CO ₃ , soluble.....				2.09
	99.98	99.81	99.93	100.21

OREGON.

1. BASALT, MOUNT THIELSEN.

Partly described by Diller in Am. Journ. Sci., 3d series, vol. 28, p. 257. A hypersthene-basalt containing hypersthene, olivine, feldspar, and magnetite. In the printed paper only the analyses of the groundmass and the fulgurite formed in it are given.

A. Hypersthene-basalt.

B. Pyroxene.

C, D. Feldspars.

E. Groundmass.

F. A fulgurite, or lightning tube.

Analyses A and F by F. W. Clarke, record Nos. 108, 105; B, C, D, and E by T. M. Chatard, record Nos. 135, 133, 134, 128. These analyses were made early in the history of the laboratory and are by no means complete.

	A.	B.	C.	D.	E.	F.
SiO ₂	55.68	53.31	55.48	51.95	55.85	55.04
Al ₂ O ₃	18.93	5.99	26.91	28.84	22.95	
Fe ₂ O ₃	8.73	13.43	2.32	2.24	4.59	28.99
FeO						
MgO	4.86	21.69	2.27	1.34	3.08	5.85
CaO	7.99	3.69	8.11	11.42	8.41	7.86
Na ₂ O	2.12	-----	3.14	3.22	2.16	-----
K ₂ O48	-----	.72	.59	2.67	-----
H ₂ O60	-----	.66	.40	.52	1.11
TiO ₂	-----	-----	.39	trace	-----	-----
P ₂ O ₅	-----	-----	-----	-----	trace	-----
	99.39	98.11	100.00	100.00	100.23	98.85

Iron oxides not separated. Analyses B, C, D, and F made on very small quantities of material. C and D were analyzed by the hydrofluoric acid method, and the silica was determined by difference.

2. PERIDOTITE, DOUGLAS COUNTY.

The matrix of the nickel silicate at Riddle. Described by Diller and Clarke in Bull. 60, p. 21.

The rock, which may be classed as saxonite, consists essentially of olivine and enstatite, with a little chromite and magnetite. Olivine predominates, and the enstatite forms less than one-third of the mass. Quartz, serpentine, and genthite are present as alteration products. Analyses by F. W. Clarke, record Nos. 811, 814, 792.

- A. The fresh rock. P. R. C. 114.
- B. Separated olivine.
- C. The genthite found in the altered rock. Probably derived from olivine.

	A.	B.	C.
SiO ₂	41.43	42.81	44.73
Al ₂ O ₃04	-----	} 1.18
Fe ₂ O ₃	2.52	2.61	
FeO.....	6.25	7.20	-----
MgO.....	43.74	45.12	10.56
CaO.....	.55	none	-----
H ₂ O at 110°.....	-----	-----	8.87
H ₂ O ign.....	4.41	.57	6.99
Cr ₂ O ₃76	.79	-----
MnO.....	none	none	-----
NiO.....	.10	.26	27.57
	99.80	99.36	99.90

3. CRATER LAKE.

Rocks collected by J. S. Diller. Petrographic data, hitherto unpublished, supplied by Horace B. Patton for all except the last rock in the series. Analyses A to J, inclusive, by H. N. Stokes, record No. 1671.

- A. Vitrophyric rhyolite, south edge of Llao Rock flow. Contains plagioclase, hypersthene, hornblende, and apatite in a glassy ground-mass crowded with augite microlites.
- B. Streaked rhyolite, near "Wine Glass" Grotto Cove. Contains plagioclase, hypersthene, hornblende, and magnetite, with black glass. A few small inclusions of basalt and hypersthene-andesite.
- C. Rhyolite, small dike immediately below Llao Rock. Contains plagioclase, hornblende, hypersthene, and magnetite, in a glassy groundmass crowded with microlites of feldspar and augite.

D. Rhyolite, water's edge, head of Cleetwood Cove. Contains plagioclase, hypersthene, brown hornblende, and magnetite, in a feldspathic groundmass of trachytic type.

E. Hypersthene-augite-andesite, large dike transsecting the north-western portion of the crater rim. Contains plagioclase, hypersthene, augite, and magnetite in a groundmass having a moderate amount of glass.

F. Hypersthene-augite-andesite, west edge of Wizard Island. Same minerals as E.

	A.	B.	C.	D.	E.	F.
SiO ₂	70.77	68.17	71.87	70.10	60.09	59.39
Al ₂ O ₃	14.83	15.60	14.53	15.18	17.85	18.45
Fe ₂ O ₃	1.35	2.31	1.28	1.78	2.03	1.79
FeO.....	1.25	.94	1.02	1.09	3.45	3.90
MgO.....	.64	1.02	.48	.74	3.50	3.13
CaO.....	2.12	2.76	1.59	2.27	6.28	6.29
Na ₂ O.....	5.07	5.15	5.08	5.15	4.17	4.29
K ₂ O.....	2.68	2.46	2.84	2.58	1.31	1.29
H ₂ O at 110°...	.07	.09	.06	.10	.12	.10
H ₂ O above 110°	.33	.45	.22	.19	.26	.42
TiO ₂38	.54	.41	.48	.54	.41
P ₂ O ₅13	.13	.10	.13	.23	.22
ZrO ₂05	none	.04	.04	none	none
NiO.....	none	none	none	none	.05	none
SrO.....	.02	.03	.03	.03	.05	.04
BaO.....	.08	.06	.08	.08	.05	.05
Li ₂ O.....	trace	trace	trace	trace	trace	trace
Cl.....	.11	trace	trace	.03	trace	trace
	99.88	99.71	99.63	99.97	99.98	99.77

Traces of manganese in all. Fluorine not sought for. No CO₂, S, SO₃, or Cr₂O₃ in any.

G. Hypersthene-augite-andesite, crater rim, just south of "The Watchman." Same minerals as F.

H. Hypersthene-augite-andesite, Palisades, under Round Top, north-east portion of the rim. Contains plagioclase, hypersthene, augite, and magnetite.

I. Hypersthene-augite-andesite, lake level, under Liao Rock. Same minerals as H.

J. Basalt, base of Red Cone. Contains plagioclase, augite, olivine, and magnetite, with some glass base.

K. Basalt, one mile east of the summit of the Cascade Range, on the road from Fort Klamath to Crater Lake. Described by J. S. Diller as a typical basalt, carrying a considerable amount of hypersthene. Analysis by W. F. Hillebrand, record No. 408.

	G.	H.	I.	J.	K.
SiO ₂	60.98	62.09	58.41	52.99	57.47
Al ₂ O ₃	17.82	17.03	17.85	16.71	18.86
Fe ₂ O ₃	1.83	2.38	2.67	3.80	2.21
FeO	3.33	2.69	3.29	3.55	4.08
MgO	2.76	3.08	3.61	6.95	4.27
CaO	5.73	5.65	6.81	8.49	7.42
Na ₂ O	4.26	4.10	3.77	3.56	3.85
K ₂ O	1.43	1.67	1.23	1.29	.73
H ₂ O at 110°13	.04	.34	.18	} .22
H ₂ O above 110°45	.13	.86	.59	
TiO ₂71	.65	.69	1.18	.75
P ₂ O ₅17	.19	.24	.42	.24
NiO	none	none	none	.02	-----
MnO	trace	trace	trace	trace	.10
SrO05	.07	.05	.12	.11
BaO06	.07	.05	.07	.03
Li ₂ O	none	none	trace	none	-----
Cl	trace	trace?	trace	trace	-----
	99.71	99.84	99.87	99.92	100.34

4. ROCK FROM WILBUR, DOUGLAS COUNTY.

A tuff partly of igneous, partly of organic, origin. The igneous matter contains a few grains of feldspar and augite, with particles of a rock like diabase. The organic remains are partly calcareous and partly siliceous. Description supplied by J. S. Diller. Analysis by H. N. Stokes, record No. 1737.

SiO ₂	55.15
Al ₂ O ₃ <i>a</i>	9.75
Fe ₂ O ₃	7.76
MgO	2.22
CaO	10.48
Na ₂ O	1.00
K ₂ O50
H ₂ O at 110°	2.70
H ₂ O above 110°	6.59
CO ₂	3.64
	99.79

a Includes TiO₂ and P₂O₅ if present.

WASHINGTON.

Rocks from the Mount Stuart quadrangle, Kittitas County. Collected by George Otis Smith, who furnishes the petrographic data. Hitherto unpublished. Analyses A, B, C, F, and G by H. N. Stokes, record No. 1836; D, E, H, and I by W. F. Hillebrand, record No. 1831.

A. Granodiorite, south slope of Mount Stuart. Contains plagioclase, orthoclase, hornblende, biotite, quartz, and magnetite.

B. Granodiorite, ridge between Hardscrabble and Cascade creeks. Contains plagioclase, orthoclase, biotite, hornblende, quartz, magnetite, and apatite.

C. Granodiorite-porphry, dike 2 miles west of Mount Stuart. Contains plagioclase, biotite, hornblende, orthoclase, and quartz.

D. Serpentine, Three Brothers. Derived from saxonite. Contains serpentine, bastite, magnetite, and pyrite.

E. Metamorphic rock, head of Beverly Creek. Believed to be derived from an inclusion of limestone in the peridotite.

	A.	B.	C.	D.	E.
SiO ₂	64.04	63.37	63.78	39.00	32.12
Al ₂ O ₃	15.58	15.90	16.39	1.75	.82
Fe ₂ O ₃	1.26	1.41	1.12	5.16	2.05
FeO.....	3.22	3.18	2.76	1.71	3.50
MgO.....	3.23	3.33	3.27	38.00	26.73
CaO.....	4.51	4.63	4.07	trace	1.81
Na ₂ O.....	4.01	4.05	3.84	} .10	.06
K ₂ O.....	2.22	2.10	2.03		
H ₂ O at 110°.....	.19	.18	.22	1.31	.43
H ₂ O above 110°.....	1.17	1.16	1.82	12.43	.98
TiO ₂69	.69	.44	trace	trace
P ₂ O ₅16	.17	.11	trace	trace
Cr ₂ O ₃	none	none	none	.47	.27
NiO.....	none	none	none	.10	.08
MnO.....	trace	trace	.05	.15	.14
SrO.....	trace	none	trace	none	none
BaO.....	.11	.06	.08	none	none
Li ₂ O.....	trace	trace	trace	none	-----
CO ₂	none	none	none	none	31.04
S.....	trace	trace	trace	-----	-----
FeS ₂ <i>a</i>	-----	-----	-----	.03	none
	100.39	100.23	99.98	100.21	100.03

a Actual condition of sulphur not known.

F. Gabbro, east of Beverly Creek. Contains diallage and "basic" plagioclase, with pyrite and serpentine as alteration products.

G. Olivine-diabase, Camas Land. Contains augite, olivine, plagioclase, magnetite, and apatite.

H. Diabase, dike on ridge west of Turnpike Creek. Contains plagioclase, augite, olivine, magnetite, and apatite.

I. Basalt, middle fork of Teanaway River. Contains augite, plagioclase, magnetite, and apatite, with a glassy base.

	F.	G.	H.	I.
SiO ₂	48.58	51.98	57.21	53.35
Al ₂ O ₃	20.23	15.99	12.99	12.90
Fe ₂ O ₃	1.26	3.10	3.28	2.64
FeO.....	3.02	5.88	10.18	11.28
MgO.....	7.59	5.09	1.59	2.68
CaO.....	14.01	9.68	5.97	6.96
Na ₂ O.....	2.25	2.71	3.07	2.83
K ₂ O.....	.19	.81	1.61	1.40
H ₂ O at 110°.....	.28	.48	.68	.91
H ₂ O above 110°.....	2.68	2.08	1.03	1.76
TiO ₂09	1.71	1.72	2.44
P ₂ O ₅	trace	.31	.44	.45
Cr ₂ O ₃	trace	none	none	none
V ₂ O ₃			none	.04
MnO.....	trace	.10	.24	.25
NiO.....	none	none	trace	trace
SrO.....	none	none	trace	trace
BaO.....	none	.03	.06	.05
Li ₂ O.....	none	trace	trace	trace
FeS ₂13	.13
S.....	.10	.01		
	100.25	99.96	100.20	100.07

ALASKA.

Rocks A to H, inclusive, were collected by G. F. Becker, who furnishes the petrographic data. A, B, C, D, F, and H are described by Becker in the 18th Ann., Part 3, p. 7. Analyses by Hillebrand, record No. 1585.

A. Augite-bronzite-andesite, Delarof Harbor, Unga Island. Greenish black. Contains plagioclase, near labradorite, with much smaller proportion of augite and bronzite, in a groundmass of plagioclase, with a little glass and much light-green indeterminate material. Contains 0.04 V_2O_3 .

B. Augite-bronzite-andesite, St. Augustine Volcano, Cook Inlet. Purplish gray. Contains labradorite, augite, and bronzite, in a groundmass of plagioclase and magnetite.

C. Quartz-porphyry, bed of Bear Creek, 4 miles from its mouth, Turnagain Arm, Cook Inlet. Resembles D, with more feldspar and less quartz.

D. Quartz-porphyry, east of mouth of Indian River, Sitka, Baranof Island. Contains plagioclase, quartz, a little pyroxene, and some carbonaceous matter, with secondary quartz, calcite, and muscovite.

E. Diorite, head of Captains Bay, Unalaska Island. Contains plagioclase, biotite, hornblende, chlorite, magnetite, and sometimes tourmaline.

	A.	B.	C.	D.	E.
SiO ₂	56.63	60.40	62.92	65.94	58.63
Al ₂ O ₃	16.85	16.89	14.29	13.74	16.23
Fe ₂ O ₃	3.62	1.88	.84	.49	1.91
FeO	3.44	3.72	4.66	5.21	4.20
MgO	4.23	3.82	3.14	2.33	4.28
CaO	7.53	7.25	2.72	2.87	6.59
Na ₂ O	3.08	3.80	4.30	2.80	3.51
K ₂ O	2.24	.77	1.39	1.63	2.09
H ₂ O at 110°80	.09	.22	.21	.15
H ₂ O above 110°51	.20	2.84	2.59	1.17
TiO ₂67	.61	.84	.80	.74
P ₂ O ₅16	.16	.13	.21	.20
NiO	trace?	.02	trace	trace?	.02
MnO23	.12	.15	.11	.11
SrO	trace	trace	trace	trace?	trace
BaO09	.06	.10	.12	.06
Li ₂ O	trace	trace	trace	trace	trace
CO ₂	none	none	1.24	.59	none
C20	
FeS ₂06	.08	.32	.41	.04
	100.14	99.87	100.10	100.25	99.93

F. Diorite, Karluk Cliffs, Kadiak Island. Contains labradorite and hornblende, with subordinate quartz, biotite, and magnetite.

G. Diorite, Lane and Hayward mine, Silver Bow Basin. Contains plagioclase, biotite, hornblende, and scattering grains of quartz, with secondary chlorite, epidote, and muscovite.

H. Diorite, Treadwell mine, Douglas Island. Contains plagioclase, mostly albite, with secondary quartz, calcite, and pyrite, the latter apparently replacing ferromagnesian silicates. Contains $0.01 \text{ V}_2\text{O}_5$.

I. Hornblende-andesite, Bogoslof Island. Described by Merrill in Proc. U. S. National Museum, vol. 8, p. 31. Contains hornblende, augite, plagioclase, tridymite, grains of iron ore, a little apatite, and probably sanidine. P. R. C. 315.

J. Like I, but darker colored. Contains more hornblende, no glass, and little or no tridymite. Same locality, also described by Merrill. Analyses I and J, by T. M. Chatard, record Nos. 209, 210. P. R. C. 316.

	F.	G.	H.	I.	J.
SiO_2	61.58	54.20	63.01	56.07	51.54
Al_2O_3	15.89	15.86	18.48	19.06	20.31
Fe_2O_3	2.19	3.32	.06	5.39	4.64
FeO	5.50	4.14	.32	.92	3.56
MgO	2.69	3.51	.06	2.12	3.16
CaO	6.49	5.32	2.66	7.70	9.55
Na_2O	3.04	3.28	10.01	4.52	4.29
K_2O51	3.30	.39	1.24	2.47
H_2O at 110°16	.55	.05	} .99	} .34
H_2O above 110°	1.26	2.40	.27		
TiO_263	1.35	.13	1.24	.32
P_2O_512	.68	.06	.16	.57
NiO	trace ?	.02	none	-----	-----
MnO20	.19	.06	.23	.32
SrO	trace ?	.04	trace	-----	-----
BaO06	.41	.02	-----	-----
Li_2O	trace	trace	none	-----	-----
CO_2	none	1.45	2.01	-----	-----
FeS_206	.26	2.10	-----	-----
	100.38	100.28	99.69	99.64	101.07

Rocks K to S. Collected by J. E. Spurr, who furnishes the petrographic notes. All but S are to appear in the 21st annual. Analyses by Stokes, record Nos. 1809 and 1822.

K. Alaskite, Chilkoot Pass. Dike in granite. Contains quartz, orthoclase, and some twinned feldspars, accessory zircon, actinolite, magnetite, and siderite.

L. Alaskite-porphry, Fortymile Creek, near Canyon Creek. Dike. Contains quartz, orthoclase, and some plagioclase, with accessory biotite and epidote.

M. Alaskite, Skwentna River, 12 miles above its mouth. Dike. Contains quartz, orthoclase, and microcline, with no dark minerals.

N. Alaskite, Tordrillo Mountains. Dike. Consists of quartz, orthoclase, and microcline, with no dark minerals.

	K.	L.	M.	N.
SiO ₂	76.30	67.01	75.01	77.33
Al ₂ O ₃	12.50	17.91	13.88	12.55
Fe ₂ O ₃	1.47	1.30 *	.74	.91
FeO.....				
MgO.....	none	.42	.09	.10
CaO.....	.17	1.86	1.00	.17
Na ₂ O.....	3.86	5.33	3.52	3.19
K ₂ O.....	4.67	4.56	4.89	4.80
H ₂ O at 110°.....	.18	.16	.11	.15
H ₂ O above 110°.....	.32	.48	.26	.53
TiO ₂05	.10	.06	.09
BaO.....	.07	.60	.10	trace
SrO.....	none	.13	trace	trace
Li ₂ O.....	none	none	trace	trace
	99.59	99.86	99.66	99.82

In K to N there are traces of P₂O₅ but no CO₂. Manganese is also present in traces.

O. Andesite-oligoclase-scapolite-biotite rock, Skwentna River, 15 miles above its mouth. Dike. Essential constituents andesine-oligoclase, scapolite, and biotite, with accessory apatite and zircon.

P. Augite-belugite, Skwentna River, near Hayes River. Dike. Contains essential feldspar, intermediate between andesine and labradorite, augite, and some hornblende. Also a considerable amount of pyrite.

Q. Tordrillite, Tordrillo Mountains. Dike. Contains phenocrysts of quartz, orthoclase, anorthoclase, and anorthoclase-albite. Ground-mass consists of quartz and orthoclase. No dark minerals except very small quantities of secondary hornblende.

R. Augite-aleutite, near Kalina Pass, Aleutian Peninsula. Lava. Chiefly labradorite, tending toward andesine, with a considerable amount of pale-green augite.

S. Tonalite-aplite, Yukon River, above Fort Hamlin. Dike. Consists essentially of quartz and soda-lime feldspars.

	O.	P.	Q.	R.	S.
SiO ₂	62.78	50.23	75.84	56.03	74.79
Al ₂ O ₃	17.16	19.46	13.38	18.31	12.59
Fe ₂ O ₃	1.96	4.21	1.45	3.47	1.19
FeO.....	2.31	4.20		4.42	
MgO.....	2.32	3.59	.10	3.64	.31
CaO.....	4.84	10.39	.07	7.43	3.58
Na ₂ O.....	4.11	3.08	3.33	3.60	5.10
K ₂ O.....	2.15	1.32	4.73	1.18	.21
H ₂ O at 110°.....	.24	.16	.18	.12	.09
H ₂ O above 110°.....	.88	1.01	.71	.31	1.03
TiO ₂56	1.30	.09	1.24	.17
P ₂ O ₅15	.41	trace	.13	trace
MnO.....	.06	.07	trace	.11	trace
BaO.....	.04	.04	trace	trace	none
SrO.....	trace	trace	trace	trace	none
Li ₂ O.....	trace	trace	trace	trace	none
CO ₂	none	.25	none	none	.58
S (a).....	.02	.02	-----	trace	-----
Cl.....	trace	-----	-----	-----	-----
	99.58	99.74	99.88	99.99	99.64

a Whether S or SO₃ was not determined.

BRAZIL.

A kyanitic schist from Serra do Gigante, near Diamantina, was analyzed at the request of Prof. O. A. Derby, who describes the rock in Amer. Journ. Sci., 4th ser., vol. 7, p. 343. Analysis by Hillebrand, record No. 1783. Contains kyanite, chlorite, sericite, quartz, and rutile.

A. Bulk analysis of the schist.

B. Portion soluble in hydrochloric acid.

C. Portion soluble in strong sulphuric acid.

D. Residue insoluble in sulphuric acid.

The bracketed figures are deduced from other columns than the one in which they appear.

	A.	B.	C.	D.
SiO ₂	38.32	10.78	14.76	[23.56]
Al ₂ O ₃	28.16	10.42	14.77	[13.39]
Fe ₂ O ₃	2.24	[1.78]	[2.24]	none
FeO.....	4.02	[3.21]	[4.02]	none
MgO.....	12.04	9.34	[12.04]	none
CaO.....	.32	.34	.32	-----
Na ₂ O.....	.16	[.03]	[.03]	.13
K ₂ O.....	1.11	.26	[.26]	.85
H ₂ O at 105°.....	.55	[.55]	[.55]	-----
H ₂ O above 105°.....	7.46	5.36	6.80	.66
TiO ₂	4.93	.10	.20	[4.73]
P ₂ O ₅47	.47	.47	-----
ZrO ₂09	trace	(?)	[.09]
CoO, NiO.....	.04	(?)	[.04]	-----
MnO.....	.16	(?)	[.16]	-----
SrO.....	trace	(?)	(?)	-----
Li ₂ O.....	trace	trace	trace	trace
S.....	trace	-----	-----	-----
F.....	trace?	-----	-----	-----
	100.07	42.64	56.66	43.41

STONY METEORITES.

1. ROCKWOOD, TENNESSEE.

From the Crab Orchard Mountains, Cumberland County, Tennessee, about $8\frac{1}{2}$ miles west of Rockwood. Analyzed by J. E. Whitfield, record No. 735, and described by him in Bull. 60, p. 103.

A. Analysis of the material as a whole.

B. Nickel-iron separated. Forms not over 16 per cent of the mass.

C. Nodule from meteorite.

D. Portion of nodule, 94 per cent, insoluble in hydrochloric acid.

May be enstatite.

The stony part of the meteorite appears to be mainly pyroxene and anorthite. Fe_2O_3 was not determined, and FeO represents the total iron oxide.

	A.	B.	C.	D.
SiO_3	41.92	49.96	51.85
Al_2O_3	9.27	4.75	4.52
FeO	22.94	15.97	13.26
CaO	9.09	1.15	1.09
MgO	8.76	28.15	29.28
Fe.....	3.75	87.59
Ni.....	1.74	12.09
Co.....	trace	trace
Cu.....	trace	trace
P.....	.65
S.....	1.58
Cl.....	.18
	99.88	99.68	99.98	100.00

2. HAMBLLEN COUNTY, TENNESSEE.

Mass of about half and half stone and iron found about 6 miles WSW. of Morristown. Analyzed by L. G. Eakins, and described in Bull. 113, p. 61.

A. The nickel-iron.

B. The part of the stony portion soluble in hydrochloric acid; 37.63 per cent, recalculated to 100, with sulphur deducted.

C. Insoluble part of the stony portion; 62.10 per cent, recalculated to 100.

A petrographic description of this meteorite is given by Merrill in Amer. Journ. Sci., 4th series, vol. 2, p. 149, together with an analysis of the feldspar. He finds it to contain, in addition to the nickel-iron, enstatite, diallage, anorthite, olivine or monticellite, oldhamite or secondary gypsum derived from oldhamite, lawrenceite, troilite, and schreibersite.

	A.		B.	C.
Fe	90.92	SiO ₂	45.61	50.67
Ni	7.71	Al ₂ O ₃	22.62	14.89
Co80	Cr ₂ O ₃	1.32
Cu	trace	FeO.....	11.73	10.55
P19	NiO.....	1.06
S04	MnO.....76
	99.66	CaO.....	14.09	3.61
		MgO.....	3.64	17.98
		K ₂ O.....03
		Na ₂ O.....19
		P ₂ O ₅	1.25
		S
			100.00	100.00

3. WINNEBAGO COUNTY, IOWA.

Fell May 2, 1890. Sp. gr. 3.804, $28^{\circ}.5$. Analyzed by L. G. Eakins, record No. 1190, and described in Bull. 78, p. 95.

Composition of the mass.

Nickel-iron.....	19.40
Troilite.....	6.19
Soluble silicates.....	36.04
Insoluble silicates.....	38.37
	<hr/>
	100.00

Separate analyses:

A. The nickel-iron.

B. Silicate soluble in hydrochloric acid, calculated to 100 per cent.

C. Insoluble silicate, recalculated to 100 per cent. The Cr_2O_3 probably represents chromite.

	A.		B.	C.
Fe	92.65	SiO_2	39.74	55.51
Ni	6.11	Al_2O_3		5.43
Co65	Cr_2O_325
P	trace	FeO.....	18.42	9.45
S	trace	NiO.....	.38	
	99.41	MnO.....	trace	
		CaO.....	.69	3.00
		MgO.....	40.77	24.09
		K_2O	trace	.15
		Na_2O	trace	2.12
		P_2O_5	trace	
			100.00	100.00

4. TANEY COUNTY, MISSOURI.

Analysis by J. E. Whitfield, record No. 736. See Bull. 60, p. 106.
Sp. gr., 4.484.

A. The separated nickel-iron.

B. The stony portion as a whole.

C. Silicates soluble in hydrochloric acid, recalculated to 100 per cent.

D. Insoluble silicates, recalculated to 100 per cent.

	A.		B.	C.	D.
Fe	89.41	SiO ₂	45.88	26.95	52.39
Ni	10.41	Al ₂ O ₃	7.89	17.69	7.11
Co29	FeO	19.73	35.98	14.68
P16	CaO	6.02	15.98	4.49
	100.27	MgO	17.96	3.40	21.33
		NiS	1.67		
		FeS54		
			99.69	100.00	100.00

5. WASHINGTON COUNTY, KANSAS.

Fell July 25, 1890. Analyzed by L. G. Eakins, record No. 1227, and described in Bull. 90, p. 45. Sp. gr. 3.49, 21.6°.

Composition of the mass.

Nickel-iron	7.7
Troilite	5.0
Soluble silicates	46.0
Insoluble silicates	41.5
	100.2

Separate analyses.

A. The nickel-iron.

B. Silicates soluble in hydrochloric acid, calculated to 100 per cent.

C. Insoluble silicates, calculated to 100 per cent.

	A.		B.	C.
Fe	86.76	SiO ₂	38.50	53.80
Ni	12.18	Al ₂ O ₃		4.32
Co83	Cr ₂ O ₃		1.41
	99.77	FeO	23.54	* 11.98
		NiO69	
		CoO	trace	
		MnO34	trace
		CaO12	4.08
		MgO	36.81	22.37
		K ₂ O27
		Na ₂ O		1.77
			100.00	100.00

6. KIOWA COUNTY, KANSAS.

A pallasite found in Brenham Township. Analyzed by L. G. Eakins, record No. 1188, and described in Bull. 78, p. 94.

A. The nickel-iron. Sp. gr. 7.93, 23.4°.

B. The pure olivine. Sp. gr. 3.376, 23.2°.

C. Dark outer zone of olivine, containing troilite.

	A.		B.	C.
Fe	88.49	SiO ₂	40.70	34.14
Ni	10.35	Al ₂ O ₃	trace?
Co57	Fe ₂ O ₃18
Cu03	FeO	10.79	23.20
P14	NiO02	trace
S08	CoO03
Si	trace?	MnO14	.09
C	trace	MgO	48.02	40.19
	99.66	S		5.42
			99.85	103.07
		Less O=S		2.71
				100.36

7. TRAVIS COUNTY, TEXAS.

Analyzed by L. G. Eakins, record No. 1097, and described in Bull. 78, p. 91. Sp. gr. 3.543, 20°. According to Cross, the stony portion contains olivine and enstatite, with a small amount of a colorless mineral, which is probably feldspar. Chromite is also present.

Approximate composition of the mass.

Nickel-iron	2.23
Troilite	5.03
Soluble silicates	39.84
Insoluble silicates	52.42
	<hr/> 99.52

A. Total analysis.

B. Nickel-iron.

C. Silicates soluble in hydrochloric acid, calculated to 100 per cent.

D. Insoluble silicates, calculated to 100 per cent.

	A.	B.	C.	D.
SiO ₂	44.75		38.13	56.14
Al ₂ O ₃	2.72		2.58	3.73
Cr ₂ O ₃52			1.00
CuO	trace			
FeO	16.04		19.76	9.15
Fe	1.83	88.74		
NiO52		1.19	
Ni22	10.68		
Co01	.58		
MnO	trace			
CaO	2.23		1.02	3.59
MgO	27.93		37.32	24.44
K ₂ O13		undet.	.19
Na ₂ O	1.13		undet.	1.76
P ₂ O ₅41			
S	1.83			
H ₂ O84			
	101.11	100.00	100.00	100.00
Less O=S92			
	100.19			

8. BLUFF, FAYETTE COUNTY, TEXAS.

Analyzed by J. E. Whitfield, record No. 824, and described in Bull. 60, p. 107. Sp. gr. 3.510. Examined microscopically by G. P. Merrill, who reports, in addition to nickel-iron and pyrrhotite, olivine, enstatite, and what appears to be augite or an allied pyroxene. See Amer. Journ. Sci., 3d series, vol. 36, August, 1888.

A. Total analysis.

B. Nickel-iron (5.67 per cent of total).

C. Part soluble in hydrochloric acid, calculated to 100 per cent (60.62 per cent of total).

D. Insoluble part, calculated to 100 per cent (33.3 per cent of total).

	A.	B.	C.	D.
SiO ₂	37.70	-----	33.59	49.64
Al ₂ O ₃	2.17	-----	1.34	4.12
FeO	23.82	-----	31.12	15.56
Fe	4.41	82.42	-----	-----
NiO	1.59	-----	2.66	trace
Ni88	15.44	-----	-----
CoO16	-----	.27	trace
Co37	2.14	-----	-----
MnO45	-----	.43	.54
CaO	2.20	-----	1.00	4.93
MgO	25.94	-----	28.08	25.21
P ₂ O ₅25	-----	.42	-----
S	1.30	-----	2.18	-----
	101.24	100.00	101.09	100.00
Less O=S65	-----	1.09	-----
	100.59	-----	100.00	-----

This meteorite also contained a dark vein of sp. gr. 3.585, which carried 2.30 per cent of metallic iron. Analyses, made on less than 0.4 gramme of material, gave as follows, recalculated to 100 per cent:

E. Soluble in hydrochloric acid, metal deducted.

F. Insoluble.

	E.	F.
SiO ₂	27.63	56.52
Al ₂ O ₃	2.41	1.51
FeO	34.31	12.35
NiO+CoO	3.27	4.09
CaO	trace	trace
MgO	32.12	25.53
S52
	100.26	100.00
Less O=S26	
	100.00	

A represents 51 per cent and F 44 per cent of the vein.

9. SAN BERNARDINO COUNTY, CALIFORNIA.

Found in the San Emigdio Mountains. Analyzed by J. E. Whitfield, record Nos. 804 and 936, and described in Bull. 60, p. 114. In fragments, badly altered.

Approximate composition of the mass.

Nickel-iron	6.21
Soluble silicates, etc.	51.26
Insoluble silicates.....	42.23
	<hr/>
	99.70

The soluble part was probably olivine and pyrrhotite, with secondary iron oxide. The insoluble part was enstatite, essentially.

A. The nickel-iron.

B. The enstatite.

	A.		B.
Fe	88.25	SiO ₂	54.42
Ni	11.27	FeO	14.03
Co48	CaO	2.46
	100.00	MgO	29.11
			<hr/>
			100.02

10. BEAVER CREEK, BRITISH COLUMBIA.

Fell May 26, 1893, near Beaver Creek, West Kootenai district. Described by Howell, Hillebrand, and Merrill in Amer. Journ. Sci., 3d series, vol. 47, p. 430.

Composition of the mass.

Nickel-iron	17.13
Magnetite16
Troilite	5.05
Soluble silicates and phosphate	37.23
Insoluble silicates and chromite	40.43
	<hr/> 100.00

According to Merrill, the silicates visible are olivine, enstatite, probably a little plagioclase, and some glassy base.

Analyses by W. F. Hillebrand, record No. 1444.

A. Nickel-iron.

B. Nonmagnetic, stony portion.

C. Portion of B soluble in hydrochloric acid, calculated to 100 per cent.

D. Insoluble portion, calculated to 100 per cent. From C and D troilite and chromite are excluded. The chromite forms about 0.75 per cent of the stony matter.

	A.		B.	C.	D.
Fe	90.68	SiO ₂	45.87	38.26	57.75
Ni	8.80	TiO ₂09		.18
Co49	Al ₂ O ₃	2.30	.56	4.89
Cu03	Cr ₂ O ₃51		
	100.00	FeO	12.68	19.52	8.02
		Fe	3.87		
		NiO07	.09	trace
		MnO26	.27	.35
		CaO	1.96	1.03	3.44
		MgO	28.24	38.74	23.19
		K ₂ O15	.02	.25
		Na ₂ O98	.13	1.87
		H ₂ O34	.70	.06
		P ₂ O ₅30	.68	
		S	2.21		
		Cl	trace	trace	
			99.83	100.00	100.00
		Troilite	6.08		
		Chromite75		

11. LLANO DEL INCA, CHILE.

Analyzed by L. G. Eakins, record No. 1201, and described in Bull. 78, p. 97.

Approximate composition of the mass.

Nickel-iron	25.8
Troilite	10.6
Soluble silicates	30.9
Insoluble silicates	32.6
	<hr/>
	99.9

Separate analyses.

A. The nickel-iron.

B. Silicates soluble in hydrochloric acid, calculated to 100 per cent.

C. Insoluble silicates, calculated to 100 per cent.

Is the P_2O_5 in B derived from schreibersite?

	A.		B.	C.
Fe	89.77	SiO_2	28.08	53.11
Ni	9.17	Al_2O_3	12.74	2.32
Co61	Cr_2O_390
	<hr/>	FeO	42.52	18.82
	99.55	NiO	2.90	-----
		MnO20	-----
		CaO	9.33	1.75
		MgO	1.98	23.10
		P_2O_5	2.25	-----
			<hr/>	<hr/>
			100.00	100.00

METEORIC IRON.

A. The Mount Joy meteorite, found near Two Taverns post-office, near Gettysburg, Pennsylvania. Analysis by L. G. Eakins, record No. 1318.

B. From Pulaski County, Virginia. Sp. gr. 7.95, 23°. Analysis by Eakins, No. 1228. Described by Eakins in Bull. 90, p. 45.

C. From Ellenboro, Rutherford County, North Carolina. Described and analyzed by Eakins, Bull. 78, p. 93. Record No. 1160.

D. From Linnville Mountain, North Carolina. Analyzed by J. E. Whitfield, record No. 822, and described in Bull. 60, p. 107. Sp. gr. 7.778.

E. From Cherokee County, Georgia. Analysis by H. N. Stokes, record No. 1527.

	A.	B.	C.	D.	E.
Fe	93.80	93.59	88.05	84.56	91.96
Ni	4.81	5.56	10.37	14.95	6.70
Co51	.53	.68	.33	.50
Cu005	trace	.04	-----	.03
P19	.27	.21	trace	.11
S01	.01	.08	.12	.01
Si		trace	.02	none	trace
C			-----	trace	trace?
	90.325	99.96	99.45	99.96	99.31

F. From near Holland's store, Chattooga County, Georgia. Analysis by J. E. Whitfield, record No. 765. See Bull. 60, p. 106. Sp. gr. 7.801.

G. From Hamilton County, Texas. Sp. gr. 7.95, 27°. Analysis by L. G. Eakins, record No. 1189. See Bull. 78, p. 95.

H. From near Scottsville, Allen County, Kentucky. Analysis by Whitfield, record No. 509. See Bull. 55, p. 64.

I. Fell 6 miles east of Cabin Creek, Johnson County, Arkansas, March 27, 1886. Analysis by Whitfield, record No. 505. See Bull. 55, p. 63.

J. From near Grand Rapids, Michigan. Sp. gr. 7.87. Analysis by R. B. Riggs, record No. 296. See Bull. 42, p. 94.

	F.	G.	H.	I.	J.
Fe	94.60	86.54	94.32	91.87	88.71
Ni	4.97	12.77	5.01	6.60	10.69
Co21	.63	trace	trace
Cu0207
Mg02
Mn	trace
P21	.16	.16	.41	.26
S	trace	.03	.34	.05	.03
Si	none
C	trace	.11	.12	.15	.13
Insoluble34
	99.99	100.26	99.95	99.42	99.91

- K. The El Capitan iron, from near Bonito, New Mexico. Analysis by H. N. Stokes, record No. 1527.
- L. From La Bella Roca, Sierra de San Francisco, Mexico, State of Durango. Analysis by J. E. Whitfield, record No. 1037. Bull. 64, p. 28.
- M. Troilite nodule from L, outer part, somewhat altered.
- N. Troilite nodule M, inner part. Analyses L and M also by Whitfield, record No. 1037.
- O. From Puquios, Chile. Sp. gr. 7.93, 25.2°. Analysis by L. G. Eakins, record No. 1181. See Bull. 78, p. 95.
- P. The Abert iron, of unknown origin. Analysis by R. B. Riggs, record No. 356. Sp. gr. 7.89. See Bull. 42, p. 94.

	K.	L.	M.	N.	O.	P.
Fe	90.51	91.48	9.37	88.67	92.07
Ni	8.40	7.92	9.83	7.01
Co60	.2271	.66
Cu0504
P24	.2117	.08
S	trace	.2109	.01
Si	trace?
C0604	.05
NiS	2.07	2.13
FeS	37.51	85.27
Fe ₂ O ₃	37.80
Moisture	19.85
	99.80	100.10	97.23	96.77	99.55	99.88

The two following meteoric irons were analyzed by L. G. Eakins in the Denver laboratory.

A. Found near Albuquerque, New Mexico. Described by Eakins in Proc. Colorado Sci. Soc., vol. 2, p. 14.

B. From Wyoming. Partial analysis.

	A.	B.
Fe	88.76	89.26
Ni	9.86	5.94
Co51	.78
Cu034
Zn03
Mn	trace
P182	.24
S012
Si044
C	undet.
	99.432	96.22

SANDSTONES, CHERTS AND SINTERS.

SANDSTONES FROM OHIO.

A. Blue sandstone from near Cleveland. Analysis by T. M. Chatard, record No. 214.

B. Sandstone from Berea. Analysis by L. G. Eakins, record No. 914.

C, D, E. Three samples of the "Peebles-Henley sandstone," from Portsmouth. Analyses by H. M. Stokes, record No. 1239. Alkalies, etc., undetermined.

	A.	B.	C.	D.	E.
SiO ₂	91.67	92.91	90.40	89.32	87.12
Al ₂ O ₃	6.92	3.78	5.15	5.52	5.96
Fe ₂ O ₃	trace	trace	.65	.87	.85
FeO91	.27	.35	.85
MgO34	trace	.28	.51	.73
CaO28	.31
Na ₂ O34
K ₂ O61
H ₂ O	1.17	1.19	.99	1.49	2.00
	100.38	100.05	97.74	98.06	97.51
Insol. in HCl	97.50	98.00	96.90	95.52

SANDSTONES FROM CALIFORNIA.

1. SANDSTONES FROM MOUNT DIABLO.

Described by Turner and Melville in Bull. Geol. Soc. Amer., vol. 2, pp. 383-414. Analyses made by W. H. Melville in the San Francisco laboratory.

A. Upper Cretaceous, Chico sandstone. Light brown, finely granular, carrying grains of mica and feldspar.

B. Lower Cretaceous, Neocomian sandstone. Hard, granular, greenish. From head waters of Bagley Creek.

C. Miocene sandstone. Granular, particles light brown and black, friable. From near Wall Point.

D. Chico sandstone. Greenish gray, compact, crystalline.

	A.	B.	C.	D.
SiO ₂	73.71	56.84	44.54	36.93
Al ₂ O ₃	10.40	11.37	12.63	7.22
Fe ₂ O ₃	3.89	1.46	2.50	1.59
FeO	1.88	4.95	3.08	2.95
MgO	1.62	3.10	5.55	2.34
CaO96	7.62	14.65	29.34
Na ₂ O	3.48	3.26	3.35	2.94
K ₂ O99	.86	1.37	.64
H ₂ O at 100°	1.06	1.45	1.43	.57
H ₂ O above 100°	2.60	3.34	2.25	3.45
P ₂ O ₅	none	.10	.29	.16
MnO17	.22	.44	.57
CO ₂	none	5.10	7.76	<i>a</i> 11.30
	100.76	99.67	99.84	100.00

a CO₂ determined by difference.

Traces of organic matter are found in all four sandstones, but were not determined.

2. SANDSTONE DIKES IN SHASTA COUNTY.

Described by J. S. Diller in Bull. Geol. Soc. Amer., vol. 1, p. 411. The rock is made up of quartz, feldspar, and biotite, with a calcite cement. Serpentine, sphene, magnetite, and zircon also occur in it, but are less common.

A. From Salt Creek, one half mile above McNett's.

B. One and one-fourth miles below Ono Bridge, north fork of Cottonwood.

C, D. Three-fourths of a mile below John Allen's, Dry Creek.

E. From John Allen's, Dry Creek.

Analyses A and B by T. M. Chatard, record No. 1106; C, D, and E by J. E. Whitfield, record Nos. 972, 973.

	A.	B.	C.	D.	E.
SiO ₂	48.13	48.10	59.10	61.60	54.55
Al ₂ O ₃	11.19	12.16	14.02	12.15	10.64
Fe ₂ O ₃	1.25	1.02	3.16	2.09	1.59
FeO.....	1.47	2.14	1.42	3.30	1.16
MgO	2.22	1.65	1.72	2.33	1.29
CaO	16.39	15.88	9.35	6.92	14.30
Na ₂ O.....	2.29	2.46	2.21	2.16	2.60
K ₂ O.....	1.17	1.56	1.49	1.41	1.68
H ₂ O at 110°78	.46	-----	-----	-----
H ₂ O at red heat.....	1.78	3.27	2.63	3.10	1.60
TiO ₂24	.47	.70	trace	trace
P ₂ O ₅14	.13	trace	.08	.10
MnO29	.26	trace	trace	1.53
BaO04	undet.	-----	-----	-----
CO ₂	12.73	10.36	4.65	5.05	9.05
SO ₃	-----	-----	trace	.27	.10
Cl	-----	-----	trace	trace	.72
	100.11	99.92	100.45	100.46	100.31

The following bedded sandstones of the same region are also described:

F. From middle fork of Cottonwood, 1 mile above Miller's.

G. Top of cascade, $1\frac{1}{2}$ miles up Byron Creek from north fork of Cottonwood.

H. Two and one-half miles above John Allen's, Dry Creek.

Analyses F and G by Chatard, record No. 1106; H by Whitfield, record No. 974.

	F.	G.	H.
SiO ₂	55.85	67.62	60.74
Al ₂ O ₃	13.20	13.63	10.25
Fe ₂ O ₃	2.56	1.25	4.31
FeO	4.77	3.27	6.21
MgO	1.90	2.34	3.69
CaO	6.93	2.80	4.97
Na ₂ O	2.60	2.78	1.83
K ₂ O	1.89	1.11	.52
H ₂ O at 110°	1.13	.64	-----
H ₂ O at redness	2.99	2.83	4.36
TiO ₂76	.48	.86
P ₂ O ₅18	.08	trace
MnO24	.15	trace
BaO	undet.	.03	-----
CO ₂	4.97	.72	2.29
SO ₃40
Cl			trace
	99.97	99.73	100.43

3. SANDSTONE FROM SULPHUR BANK.

Described by Becker in Mon. XIII, p. 92. Analysis made by Melville in the San Francisco laboratory. An altered sandstone, showing grains of quartz, plagioclase, and orthoclase.

SiO ₂	68.50	H ₂ O at 100°28
Al ₂ O ₃	12.82	H ₂ O above 100°	2.11
Fe ₂ O ₃	1.29	TiO ₂60
FeO	3.37	P ₂ O ₅16
MgO	2.21	MnO02
CaO	1.82		100.47
Na ₂ O	6.03		
K ₂ O	1.26		

MISCELLANEOUS SANDSTONES.

A. Triassic sandstone from the Jaittelle quarry, near Hancock, Maryland. Hard, compact, brown. Analysis by F. W. Clarke, record No. 613.

B. Brown sandstone, Hummelstown, Pennsylvania. Analysis by E. A. Schneider, record No. 1280. Described by Diller in Bull. 150, p. 77. P. R. C. 14.

C. Yellow sandstone from Stony Point, Michigan. Analysis by F. W. Clarke, record No. 213.

D. Carboniferous sandstone adjoining the peridotite dike of Elliott County, Kentucky. See Diller, Amer. Journ. Sci., 3d series, vol. 32, p. 125. Analysis by T. M. Chatard, record No. 354.

E. Yellow sandstone, Armejo quarry, Colorado. Analysis, partial, by T. M. Chatard, record No. 289.

	A.	B.	C.	D.	E.
SiO ₂	76.43	88.13	84.57	60.78	81.27
Al ₂ O ₃	17.78	5.81	5.90	10.54	9.81
Fe ₂ O ₃		1.77	6.48	3.27	1.44
FeO.....		.31			
MgO.....	.92	.53	.68	1.59	.42
CaO.....	.84	.20		10.15	.44
Na ₂ O.....	undet.	.06	undet.	1.41	undet.
K ₂ O.....	undet.	2.63	undet.	2.36	undet.
H ₂ O at 100°.....	2.79	.23	1.92	.85	1.19
H ₂ O above 100°.....		.26		2.32	
TiO ₂03	
P ₂ O ₅09	
MnO.....	trace			.10	
CO ₂				6.29	
	98.76	99.93	99.55	99.78	94.57
Insol. in HCl.....	88.68		91.87		95.54

F. Potsdam sandstone, Ablemans, Sauk County, Wisconsin. Described by Diller in Bull. 150, p. 80. Analysis by Schneider, record No. 1280. P. R. C. 15.

G. Banded sandstone, Peoa, Utah. Described by Diller in Bull. 150, p. 80. Analysis by Schneider, record No. 1280. P. R. C. 16.

H. Brown sandstone, Flagstaff, Arizona. Analysis by T. M. Chatard, record No. 1144.

I. Sandstone, Robinson mine, Summit County, Colorado. Analysis by L. G. Eakins, made in the Denver laboratory, partial.

	F.	G.	H.	I.
SiO ₂	99.42	96.60	<i>a</i> 79.19	56.33
Al ₂ O ₃	} .31	} 2.02	1.30	.77
Fe ₂ O ₃			2.45	.97
MgO08	.23	7.30
CaO04	7.76	14.01
Na ₂ O				undet.
K ₂ O				undet.
H ₂ O at 100°	} .18	.11	.32	undet.
H ₂ O above 100°29	2.94	undet.
CO ₂			5.77	<i>b</i> 19.04
	99.91	99.14	99.96	98.42

a Silica and insoluble matter.

b Calculated for bases.

CHERT.

From the lead-zinc region of southwestern Missouri and its extension into Kansas.

Collected by W. P. Jenney.

A. Unaltered chert, East Hollow, Belleville, Jasper County, Missouri.

B. Partly altered, same locality.

C. Altered to "cotton rock," same locality.

D. From the Surprise mine, Joplin, Missouri.

Analyses by E. A. Schneider, record No. 1205.

	A.	B.	C.	D.
SiO ₂	98.17	98.92	98.71	99.46
Al ₂ O ₃ , Fe ₂ O ₃83	.48	.43	.29
MgO.....	.01	.02	.02	trace
CaO.....	.05	.03	.03	.04
Ignition.....	.78	.42	.50	.34
	99.84	99.87	99.69	100.13

E. Blue chert, unaltered, Bonanza shaft, Galena, Kansas.

F. Same locality.

G. Altered, same locality.

H. Jasperite, Joplin, Missouri.

I. Jasperite, Galena, Kansas.

Analyses E, F, and G by E. A. Schneider, record No. 1205; H and I by L. G. Eakins, record No. 1208.

	E.	F.	G.	H.	I.
SiO ₂	99.23	98.60	99.13	95.77	97.33
Al ₂ O ₃ , Fe ₂ O ₃22	.52	.16	1.84	1.89
MgO.....	trace	trace	.01	.24	.09
CaO.....	.02	.10	trace	.54	.11
Ignition.....	.50	.40	.20	1.17	.77
	99.97	99.62	99.50	99.56	100.19

SILICEOUS SINTERS.

From the Yellowstone National Park and similar localities. Described by W. H. Weed in 9th Ann., p. 619. Analyses A to H, inclusive, by J. E. Whitfield, record Nos. 97, 100, 707, 708, 812, and 998; analysis I by E. A. Schneider, record No. 1254. A to E from Yellowstone National Park.

A. Dried siliceous jelly from Emerald Spring, Upper Basin.

B. Sinter from Solitary Spring, Upper Basin.

C. Grayish sinter from margin of Splendid Geyser.

D. Compact sinter from Old Faithful Geyser.

E. From Asta Spring, Hillside Group.

F, G, H. Three samples of sinter from Rotorua, New Zealand. For comparison with Yellowstone sinters.

I. Sinter from Mount Morgan gold mine, Queensland, Australia. Described by Weed in Am. Journ. Sci., 3d series, vol. 42, p. 165.

	A.	B.	C.	D.	E.
SiO ₂	93.37	93.88	81.95	89.54	89.72
Al ₂ O ₃	1.16	1.73	6.49	2.12	} 1.02
Fe ₂ O ₃	trace	.14	trace	trace	
MgO.....	.05	.07	.15	trace	trace
CaO.....	.29	.25	.56	1.71	2.01
Na ₂ O.....	.11	.28	2.56	1.12	-----
K ₂ O.....	.02	.23	.65	.30	-----
H ₂ O (ign.).....	4.17	3.37	7.50	5.13	7.34
NaCl.....	.08	.18	trace	trace	-----
SO ₃31	.20	.16	trace	trace
C, organic.....	.78	-----	-----	-----	-----
H, organic.....	.07	-----	-----	-----	-----
	100.41	100.33	100.02	99.92	100.09

	F.	G.	H.	I.
SiO ₂	92.47	90.28	74.63	94.02
Al ₂ O ₃	2.54	3.00	15.59	} 2.27
Fe ₂ O ₃	-----	-----	-----	
MgO.....	.15	trace	trace	trace
CaO.....	.79	.44	1.00	.07
Na ₂ O.....	-----	-----	.30	-----
K ₂ O.....	-----	-----	1.02	-----
H ₂ O at 105°.....	-----	-----	-----	1.07
Ignition.....	3.99	6.24	7.43	2.29
	99.94	99.96	99.97	99.72

THE CARBONATE ROCKS: LIMESTONE, DOLOMITE, SIDERITE, ETC.

VERMONT AND MASSACHUSETTS.

A. White marble, Rutland, Vermont. Analysis by L. G. Eakins, record No. 1213.

B. The portion of A insoluble in dilute hydrochloric acid. Same analyst and number.

C. White marble, Lee, Massachusetts. Analysis by E. A. Schneider, record No. 1279. Described by Diller in Bull. 150, p. 299. P. R. C. 116.

D. Limestone, Lee, Massachusetts. Collected by B. K. Emerson. From cut on west side of railroad. Analysis by George Steiger, record No. 1654.

E. Dolomite, Charlemont, Massachusetts. Collected by Emerson. Analysis by Eakins, record No. 1343.

F. Dolomite, Webster, Massachusetts. Collected by Emerson. Analysis by H. N. Stokes, record No. 1634.

	A.	B.	C.	D.	E.	F.
Insoluble	8.00	-----	.19	-----	-----	-----
SiO ₂ -----	-----	56.69	-----	.95	.67	1.01
Al ₂ O ₃ -----	.39	31.16	} .24	.09	trace	.17
Fe ₂ O ₃ -----	-----	-----		none	.08	none
FeO-----	.14	2.13	-----	.10	7.60	.37
MnO-----	-----	-----	-----	-----	1.61	.08
NiO-----	-----	-----	-----	-----	.03	-----
CaO-----	50.79	2.68	30.88	54.75	28.63	30.82
MgO-----	trace	3.27	21.42	.56	16.17	21.35
K ₂ O-----	-----	undet.	-----	.15	-----	.10
Na ₂ O-----	-----	undet.	-----	.02	-----	.01
H ₂ O-----	1.01	-----	-----	.08	-----	.09
P ₂ O ₅ -----	-----	-----	-----	.03	-----	.06
CO ₂ -----	39.80	-----	46.72	43.38	45.35	45.84
SO ₃ -----	-----	-----	-----	.05	-----	-----
	100.13	95.93	99.45	100.16	100.14	99.90

NEW YORK, PENNSYLVANIA, MARYLAND.

- A. Dolomite-marble, New York Quarry Company, Tuckahoe, Westchester County, New York. Analysis by W. F. Hillebrand, record No. 746.
- B. Hydraulic-cement rock, Akron, New York. Analysis by George Steiger, record No. 1655.
- C. Compact gray limestone, Greason, Pennsylvania. Analysis by E. A. Schneider, record No. 1279.
- D. Dolomite-marble, Cockeysville, Maryland. Analysis by Schneider, No. 1279. P. R. C. 117.
- E. An earlier sample of D. Analysis by J. E. Whitfield, record No. 827. P. R. C. 117.
- Rocks B, C, D, E are described by Diller in Bull. 150, pp. 127, 133, 300.

	A.	B.	C.	D.	E.
Insoluble.....	1.33	-----	11.07	5.57	-----
SiO ₂		9.03	-----	-----	.44
TiO ₂16	-----	-----	-----
Al ₂ O ₃		2.25	-----	} .40	1.22
Fe ₂ O ₃21	.85	-----		-----
FeO.....		.52	-----		trace
CaO.....	30.68	26.84	39.26	29.08	30.73
MgO.....	20.71	18.37	9.00	20.30	20.87
K ₂ O.....		.85	-----	-----	-----
Na ₂ O.....		none	-----	-----	-----
H ₂ O.....	.16	.98	.18	-----	1.22
P ₂ O ₅03	-----	-----	-----
CO ₂	46.66	40.33	38.82	44.26	45.85
Organic matter.....		-----	.75	-----	-----
	99.75	100.21	99.08	99.61	100.33

VIRGINIA AND WEST VIRGINIA.

A. Limestone, upper ledge, Moundsville Narrows, West Virginia.

B. Same as A. lower ledge. Analysis A and B by T. M. Chatard, record No. 127.

C. Trenton limestone, Lexington, Virginia. Analysis by R. B. Riggs, record No. 365.

D. Limestone, Staunton, Virginia. Analysis by George Steiger, record No. 1630.

E. Part of D insoluble in one-tenth hydrochloric acid.

F. Soluble part of D. Analyses E and F also by Steiger, same number. These three analyses are accompanied by analyses of the residual clay, formed by the weathering of the limestone. See section on clays.

	A.	B.	C.	D.	E.	F.
Insoluble	10.33	1.53				
SiO ₂44	7.37	6.98	.39
TiO ₂09	.09	none
Al ₂ O ₃				1.92	1.39	.53
Fe ₂ O ₃90	.96	.42	.29	.25	.04
FeO63	none	.63
MnO	trace	trace		none	none	none
CaO	48.02	53.26	54.77	28.39	.04	28.35
MgO	1.08	.93	trace	18.30	.15	18.15
K ₂ O				1.09	.91	.18
Na ₂ O09	.04	.05
H ₂ O at 100°05	.10	1.08	.09	undet.	undet.
H ₂ O above 100° }				.49	.15	.34
P ₂ O ₅	trace	trace		.03	none	.03
CO ₂	39.18	43.16	42.72	41.85	none	41.85
	99.56	99.94	99.43	100.63	10.00	90.54

GEORGIA.

A. Marble, Happy Valley.

B. "Creole" marble, Happy Valley.

C. Portion of B insoluble in dilute hydrochloric acid.

D. "Cherokee" marble, Happy Valley.

E. Portion of D insoluble in dilute hydrochloric acid.

Analyses by L. G. Eakins, record Nos. 464, 485.

	A.	B.	C.	D.	E.
Insoluble		1.84		2.01	
SiO ₂	2.23		58.21		55.48
Al ₂ O ₃91	.17	7.37	.15	15.58
Fe ₂ O ₃22				
FeO05	.31	.06	trace
CaO	52.16	53.91	12.53	53.69	14.52
MgO	2.09	.83	20.42	.83	12.88
H ₂ O45	.13		.17	
CO ₂	42.22	43.16		43.13	
	100.28	99.69	98.84	100.04	98.46

FLORIDA.

1. CORAL AND SHELL ROCKS, COLLECTED BY N. S. SHALER.

Partial analyses only. Chlorides and sulphates present; alkalies and phosphates not looked for.

A, B, C, D. Coquina gravel, Tortugas.

E. The same, Key West.

Analyses by F. W. Clarke, record No. 878.

	A.	B.	C.	D.	E.
SiO ₂19	.22	.32	.21	.25
Al ₂ O ₃ , Fe ₂ O ₃19	.47	.56	.76	.56
CaO	52.24	51.24	49.38	51.95	51.52
MgO	1.53	2.09	1.93	1.44	2.08
CO ₂	41.46	41.07	40.39	41.53	41.58
H ₂ O	3.27	3.57	5.12	3.07	3.19
	98.88	98.66	97.70	98.96	99.18

F. Near Fort Worth.

G. East side of St. Johns River, near Seville.

H. Corroded surface, Miami Reef.

I. Near Oak Hill.

J. Near Melbourne.

Analyses by F. W. Clarke, record No. 885.

	F.	G.	H.	I.	J.
SiO ₂	2.94	8.50	2.99	5.87	17.83
Al ₂ O ₃ , Fe ₂ O ₃23	.73	.65	.95	1.18
CaO.....	51.51	47.29	51.22	50.34	43.85
MgO.....	.71	1.51	.06	.37	.26
CO ₂	41.59	39.00	41.22	39.62	34.31
H ₂ O.....	2.64	3.37	2.23	3.21	2.53
	99.62	100.40	98.37	100.36	99.96

K. Coarse shell mass, Senote.

L. Coral rock, Salt Key Bank.

M. Loggerhead Key.

Analyses by L. G. Eakins, record No. 882. The following analyses, also by Eakins, No. 883, are included for comparison:

N. Coralline bottom, Barbados.

O. Recent coral (*Siderostria*), Bermuda.

	K.	L.	M.	N.	O.
SiO ₂22	.11	.20	1.17	.23
Al ₂ O ₃18	.04	.22	.31	trace
Fe ₂ O ₃					
CaO.....	54.87	53.54	53.54	46.45	55.16
MgO.....	.64	.71	.78	5.15	.20
CO ₂	43.89	43.87	43.71	43.40	43.74
H ₂ O.....	.11	1.13	.81	2.73	.54
	99.91	99.40	99.26	99.21	99.87

2. MISCELLANEOUS ROCKS.

A to M inclusive. Thirteen borings from the artesian well at Key West. Partial analyses by Steiger, record No. 1553. The figures at tops of columns give depths in feet from which samples were taken.

	A.	B.	C.	D.	E.	F.
	25	100	150	350	600	775
SiO ₂17	.25	.12	3.52	5.10	.13
Al ₂ O ₃20	.17	.08	.40	.35	.14
Fe ₂ O ₃07	.07				
CaO.....	54.03	54.01	54.38	51.46	48.87	46.53
MgO.....	.29	.77	.86	1.67	2.50	6.70
CO ₂	42.52	42.84	43.36	41.77	40.72	43.60
	97.28	98.11	98.80	98.82	97.54	97.10

	G.	H.	I.	J.	K.	L.	M.
	1125	1325	1400	1475	1625	1850	2000
SiO ₂05	.07	.19	.06	.05	.03	.07
Al ₂ O ₃ , Fe ₂ O ₃21	.11	.16	.14	.17	.17	.16
CaO.....	53.84	54.49	55.12	54.48	53.90	54.28	54.02
MgO.....	.86	.62	.30	.73	1.14	1.12	1.06
CO ₂	42.87	43.29	43.28	43.38	43.37	43.13	43.20
	97.83	98.58	99.05	98.79	98.63	98.73	98.51

P₂O₅ is present. Is included with Al₂O₃ and Fe₂O₃.

N. Supposed cement rock, River Junction. Received from D. T. Day. Analysis by George Steiger; record No. 1844.

SiO ₂	12.31
Al ₂ O ₃	12.19
Fe ₂ O ₃66
CaO.....	26.28
MgO.....	16.72
K ₂ O.....	none
Na ₂ O.....	.50
H ₂ O at 100°.....	.94
H ₂ O above 100°.....	2.05
P ₂ O ₅05
CO ₂	38.12
	<hr/> 99.82

TENNESSEE, ALABAMA, LOUISIANA.

A. Limestone, Knoxville, Tennessee. Analysis by L. G. Eakins, record No. 1159.

B. Knox dolomite, Morrisville, Alabama. Described by Russell in Bull. 52, together with a residual clay derived from it. See section on clays. Analysis by W. F. Hillebrand, record No. 797.

C. Limestone from Rayborn's salt-lick, Bienville Parish, Louisiana. Analysis by R. B. Riggs, record No. 323.

D. White marble, streaked with black. From 5 miles west of Winnfield, Louisiana. Analysis by W. F. Hillebrand, record No. 760. In addition to the constituents named in the table, this marble contains traces of barium, strontium, chlorine, and organic matter.

	A.	B.	C.	D.
Insoluble65
SiO ₂17	3.24	.55	
Al ₂ O ₃04	.17	} 1.61	trace
Fe ₂ O ₃23	.17		
FeO06		trace
MnO			trace	.10
CaO	55.47	29.58	54.09	55.01
MgO30	20.84	.06	.60
H ₂ O21	.30		.13
P ₂ O ₅05	
CO ₂	43.63	45.54	44.12	43.43
SO ₃05	.27
	100.05	99.90	100.53	100.19

TEXAS.

Supposed cement rock, Uvalde quadrangle. Collected by T. Wayland Vaughan. Analysis by W. F. Hillebrand, record No. 1759.

A, the rock; B, the part insoluble in dilute nitric acid.

	A.		B.
SiO ₂ 10	SiO ₂	20. 80
Fe ₂ O ₃ , Al ₂ O ₃ 09	Al ₂ O ₃	} 3. 44
CaO.....	40. 82	Fe ₂ O ₃	
MgO.....	. 18	P ₂ O ₅	
CO ₂	32. 41	H ₂ O.....	1. 14
Insoluble.....	25. 57	Alkalies.....	. 19
NaCl, Mn, H ₂ O <i>a</i> 83		25. 57
	100. 00		

a By difference.

OHIO.

A, B, C, D. Trenton limestone from New Vienna.

Partial analyses by F. W. Clarke and R. B. Riggs, record Nos. 729, 730, 732. Iron and alumina are present in soluble form.

	A.	B.	C.	D.
Insoluble.....	8. 47	9. 93	2. 12	28. 43
CaO.....	47. 16	49. 04	51. 18	23. 00
MgO.....	1. 52	. 58	3. 08	12. 90
CO ₂	36. 20	37. 64	42. 04	30. 82
	92. 35	97. 19	98. 42	95. 15

E. Trenton limestone, Arcadia, Hancock County.

F. Air-line Junction, Toledo. This sample contains a great deal of ferrous carbonate.

G. Gas rock, St. Henry's well, Mercer County.

H. Oil rock, Lima.

All Trenton. Analyses, partial, by Clarke and Riggs, record Nos. 729, 730.

	E.	F.	G.	H.
Insoluble.....	8. 56	3. 52	2. 27	1. 64
CaO.....	47. 17	30. 64	50. 34	32. 24
MgO.....	2. 59	18. 05	2. 86	17. 36
CO ₂	38. 54	42. 82	40. 96	43. 92
	96. 86	95. 03	96. 43	95. 16

The following partial analyses by F. W. Clarke, record No. 738, all relate to Trenton limestones:

- A. Well No. 3, Bryan. Gas rock.
 B. McElree well, Kenton. Depth, 1,315 feet.
 C. Huntsville. Depth, 1,405 feet.
 D. Prospect. Depth, 1,650 feet.
 E. Findlay street well, Dayton. Depth, 975 feet.
 F. Xenia. Depth, 1,075 feet.
 G. New Madison. Depth, 1,150 feet.

	A.	B.	C.	D.
Insoluble	9.22	5.26	4.41	26.12
Fe ₂ O ₃ , Al ₂ O ₃	1.51	1.10	3.15	2.57
CaCO ₃	49.00	84.32	57.23	66.02
MgCO ₃	38.59	8.43	33.16	3.77
	98.32	99.11	98.95	98.58
	E.		F.	G.
Insoluble	12.34		9.23	11.11
Fe ₂ O ₃ , Al ₂ O ₃58		.18	3.60
CaCO ₃	82.36		86.54	64.91
MgCO ₃	1.67		2.99	17.98
	96.95		98.94	97.60

In C and G there is ferrous carbonate.

Partial analyses by Charles Catlett, record Nos. 754, 755, 756, 757; all of Trenton limestones from the natural-gas belt.

A. London. Depth, 1,594 feet.

B. Air-line Junction, Toledo. Depth, 1,415 feet.

C. Celina. Depth, 1,112 feet.

D. City well No. 2, Upper Sandusky.

E. Sandusky. Depth, 2,260 feet.

F. Gas rock. Pauck well, St. Marys Township, Auglaize County.

G. Gas rock. Bennett well, St. Marys Township. Depth, 1,121 feet.

H. First city well, Carey. Depth, about 1,350 feet.

I. Well No. 2, Fort Recovery. Depth, 1,065 feet.

J. Waggoner well, 6 miles west of Fremont. Gas rock.

K. Loomis and Nyman well, Tiffin. Depth, 1,470 to 1,481 feet.

L. Loomis and Nyman well, Tiffin. Depth, 1,488 to 1,494 feet.

M. Port Clinton. Depth, 1,660 to 1,700 feet.

N. Wauseon. Depth, 2,135 feet.

O. Napoleon. Depth, 1,830 feet.

P. Kossuth, Allen County. Oil rock.

Q. Doenze's well, Franklin Township, Mercer County. Depth, 1,107 feet.

	A.	B.	C.	D.	E.	F.
Insoluble	15.90	2.88	2.95	8.18	3.65	3.18
Al ₂ O ₃ , Fe ₂ O ₃	1.84	8.68	2.95	4.31	4.58	3.12
CaCO ₃	77.69	54.68	68.41	64.25	54.62	52.18
MgCO ₃	1.89	25.73	24.18	15.93	33.67	38.42
	97.32	91.97	98.49	92.67	96.52	96.90
	G.	H.	I.	J.	K.	L.
Insoluble	1.66	5.72	1.89	5.22	5.66	9.88
Al ₂ O ₃ , Fe ₂ O ₃	2.48	3.08	1.57	6.32	4.86	1.46
CaCO ₃	56.94	80.11	87.88	52.93	52.89	79.39
MgCO ₃	35.55	8.09	7.43	32.75	33.46	6.20
	96.63	97.00	98.77	97.22	96.87	96.93
	M.	N.	O.	P.	Q.	
Insoluble	7.46	18.24	2.66	1.08	3.68	
Al ₂ O ₃ , Fe ₂ O ₃	4.16	7.28	2.14	.66	8.38	
CaCO ₃	71.96	42.82	53.85	90.72	69.53	
MgCO ₃	14.34	28.11	37.33	6.69	10.98	
	97.92	96.45	95.98	99.15	92.27	

INDIANA,

A. Buff limestone, Hoosier Stone Company, Bedford.
B. Blue limestone, same locality.
Analyses by F. W. Clarke, record Nos. 306, 307.

	A.	B.
SiO ₂ 63	1. 69
Fe ₂ O ₃ 39	. 49
CaO.....	54. 19	54. 18
MgO.....	. 39	. 37
P ₂ O ₅	trace	trace
CO ₂	44. 01	43. 08
SO ₃	none	none
	99. 61	99. 81

The following Trenton limestones are all from the natural-gas belt:
C. Union City. Depth, 1,160 feet.
D. Bluffton. Depth, 1,062 to 1,067 feet.
E. Muncie. Depth, 920 feet.
F. Greensburg. Depth, 867 feet.
G. Vernon. Depth, 905 feet.
H. Wabash. Depth, about 900 feet.
Analysis C by F. W. Clarke, record No. 738. The remainder by Charles Catlett, Nos. 753, 758.

	C.	D.	E.	F.	G.	H.
Insoluble.....	2. 14	2. 37	3. 30	. 87	8. 00	3. 52
Al ₂ O ₃ , Fe ₂ O ₃ ...	1. 23	4. 48	3. 72	. 55	. 60	7. 58
CaCO ₃	83. 21	53. 43	51. 96	94. 60	85. 56	53. 18
MgCO ₃	12. 48	37. 47	38. 11	. 36	trace	30. 53
	99. 06	97. 75	97. 09	96. 38	94. 16	94. 81

MISSOURI.

Limestones and dolomites collected by W. P. Jenney. Analyses by L. G. Eakins, record Nos. 1184, 1207. CO₂ calculated.

A, B. Cherokee limestone, quarry near Seneca, Newton County.

C, D. The same, near Grand Falls, Newton County.

E, F. Dolomite, Oswego land, Joplin.

	A.	B.	C.	D.	E.	F.
Insoluble .	.66	1.21	1.01	1.01	29.77	11.66
Al ₂ O ₃11	.13	.08	.13	1.32	1.03
FeO.....	.05	.07	.05	trace		
MnO.....	trace	trace	.03	trace		
CaO.....	55.29	54.92	54.98	55.11	21.46	28.72
MgO.....	.23	.20	.31	.32	14.79	17.26
CO ₂	43.69	43.31	43.54	43.65	33.13	41.55
	100.03	99.84	100.00	100.22	100.47	100.22

KANSAS.

A. Limestone, Silverdale. Analysis by Charles Catlett, record No. 967.

B. Cherokee limestone, Short Creek, near Spring River, Cherokee County. Analysis by L. G. Eakins, record No. 1184.

C. Supposed marl, large surface deposit near Wakeeney, Trego County. Analysis by F. W. Clarke, record No. 212.

	A.	B.	C.
Insoluble.....		.32	
SiO ₂	5.27		14.06
Al ₂ O ₃	1.07	.17	
Fe ₂ O ₃71		5.10
FeO.....	.32	.20	
MnO.....		.02	
CaO.....	50.36	55.25	43.05
MgO.....	.56	.35	.50
K ₂ O.....	.10		
Na ₂ O.....	.20		
H ₂ O.....	.78		1.77
P ₂ O ₅06		
CO ₂	40.54	43.79	35.03
SO ₂07		
	99.84	100.10	99.51

MICHIGAN, WISCONSIN, MINNESOTA, CANADA.

Most of the rocks considered under this heading were described by Irving and Van Hise in Mon. XIX, pages 131 and 191. A few other analyses, also representing Van Hise's collections, are taken from the laboratory records. The Canadian rocks are from near the boundary line, and relate directly to others gathered upon the Minnesota side.

A. Dolomite, near Sunday Lake, Gogebic district, Michigan. Analysis by W. F. Hillebrand, record No. 767.

B. Dolomite, Penokee region, Wisconsin, NW. $\frac{1}{4}$ sec. 22, T. 44 N., R. 5 W. Analysis by Hillebrand, record No. 768.

C. Limestone, bed of Slate Creek, Huron Bay slate quarries, Michigan. Analysis by T. M. Chatard, record No. 894. From laboratory records; not in the monograph cited.

D. Limestone, east end of Ogiskemannissi Lake, Minnesota. Analysis by Chatard, record No. 899. Not in monograph.

	A.	B.	C.	D.
SiO ₂	3.07	.63	7.05	41.99
Al ₂ O ₃48	1.24
Fe ₂ O ₃09	.03	1.33	.42
FeO86	.75	undet.	4.77
MnO15	.08	.19	.26
CaO	29.72	30.94	50.08	16.85
MgO	19.95	20.68	.57	8.41
H ₂ O at 105°30	.27		.05
H ₂ O ign25	1.02
P ₂ O ₅27	.05
CO ₂	45.31	46.27	39.68	24.70
SO ₃21	.32
Cl	trace	trace		
	99.45	99.65	100.11	100.08

- E. Iron carbonate, SE. $\frac{1}{4}$ sec. 20, T. 47 N., R. 43 W., Michigan.
F. Iron carbonate, south side of Sunday Lake, Michigan.
G. Iron carbonate, Palms mine, Gogebic district, Michigan.
Analyses E, F, and G by W. F. Hillebrand, record Nos. 769, 770, 771.
H. Iron carbonate, Miner and Wells Option, sec. 13, T. 47 N., R. 46 W., Michigan. Analysis by T. M. Chatard, record No. 893.
I. Iron carbonate, NW. $\frac{1}{4}$ sec. 18, T. 47 N., R. 45 W., Michigan. Analysis by Chatard, No. 895.

	E.	F.	G.	H.	I.
SiO ₂	3. 16	28. 86	46. 47	46. 01	36. 73
TiO ₂ 20	. 10	. 12	. 19
Al ₂ O ₃ 08	1. 29	. 70	. 83	. 38
Fe ₂ O ₃ 93	1. 01	. 86	1. 35	. 98
FeO.....	15. 18	37. 37	28. 57	26. 07	34. 81
MnO	1. 15	. 97	. 40	2. 09	. 52
CaO.....	26. 65	. 74	. 49	. 63	. 48
MgO	11. 01	3. 64	2. 30	2. 86	2. 74
H ₂ O at 105°.....	} . 54	} . 68	} . 60	} 1. 71	. 12
H ₂ O ign					1. 40
P ₂ O ₅ 06	trace	trace	. 07	. 01
CO ₂	41. 10	25. 21	19. 24	17. 72	22. 44
SO ₃ 15	. 16
Cl	trace		?		
FeS ₂ 34				
	100. 20	99. 97	99. 73	99. 61	100. 96

J. Iron carbonate, Penokee iron range, NE. $\frac{1}{4}$ sec. 6, T. 45, R. 2 E., Wisconsin. P. R. C. 999.

K, L. Black, slaty, carbonaceous iron carbonates, Animikie formation, Kakabikka Falls, Kaministiquia River, Canada.

M. Iron carbonate, west end of Gunflint Lake, Minnesota.

N. Iron carbonate, north side of Gunflint Lake, Minnesota.

O. Iron carbonate, north side of Gunflint Lake, Canada.

Analyses J, K, and L by R. B. Riggs, record Nos. 376, 377, 378; M, N, and O by T. M. Chatard, record Nos. 897, 898, 900.

	J.	K.	L.	M.	N.	O.
SiO ₂	15.62	37.73	54.26	58.23	46.46	23.90
TiO ₂				trace ?	trace ?	none
Al ₂ O ₃	4.27	3.41	2.57	.06	.24	.07
Fe ₂ O ₃	8.14	6.42	3.62	5.01	.64	.44
FeO.....	32.85	22.92	19.63	18.48	26.34	10.72
MnO.....	5.06	.40	.19	.25	.21	.28
CaO.....	.81	1.26	1.07	.38	1.87	22.25
MgO.....	2.66	3.98	2.93	9.59	3.10	8.52
H ₂ O at 110°.....	.68	2.74	1.20	.07	.07	none
H ₂ O at redness.....				2.01	1.15	.99
P ₂ O ₅03	.13	trace
CO ₂	30.32	18.01	14.93	5.22	19.96	32.42
SO ₃19	.14	.17
C.....		3.54	.45			
	100.41	100.41	100.85	99.52	100.31	99.76

P. Ferrodolomite, Marquette district, Michigan. Analysis by George Steiger, record No. 1473. P. R. C. 994.

Q. Ferrodolomite, Marquette district, Michigan.

R. Portion of Q insoluble in hydrochloric acid.

S. Soluble portion of Q.

Analyses Q, R, and S by George Steiger, record No. 1442. These rocks are not in the monograph cited.

	P.	Q.	R.	S.
SiO ₂	42.37	26.97	26.67	.30
Al ₂ O ₃		1.30	.12	1.18
Fe ₂ O ₃	1.09	2.31	.16	2.15
FeO.....	31.41	39.77	-----	39.77
MnO.....		.29	-----	.29
CaO.....	.50	.66	-----	.66
MgO.....	2.48	1.94	.10	1.84
Alkalies.....		.09	-----	.09
H ₂ O at 100°.....		.10	-----	
H ₂ O above 100°.....		.51	-----	
P ₂ O ₅03	-----	.03
CO ₂	21.80	26.20	-----	26.20
	99.65	100.17	27.05	72.51

YELLOWSTONE NATIONAL PARK.

Travertines described by W. H. Weed in 9th Ann., p. 619.

A. Terrace below the hotel.

B. Cupids Cave.

C. Near Sulphur Spring No. 246, Mammoth Hot Springs.

D. Extinct spring, main terrace, Mammoth Hot Springs.

E. Ridge behind main terrace.

Analysis C by F. A. Gooch, record No. 243; A, B, D, and E by J. E. Whitfield, record Nos. 240, 242, 244, 245.

	A.	B.	C.	D.	E.
SiO ₂08	.15	.01	.06	.26
Al ₂ O ₃15	.49	} .05	.14	.11
Fe ₂ O ₃					
CaO.....	53.83	53.41	55.02	55.02	54.06
MgO.....	.90	.42	.07	.06	.66
K ₂ O.....		.01	.04		
Na ₂ O.....		.03			
H ₂ O.....	1.43	2.44	1.61	1.06	1.19
NaCl.....	.02	.13	.12	.20	.26
KCl.....				.08	
SO ₃	1.72	.55	.49	.70	1.34
CO ₂	41.79	41.96	42.25	42.25	42.14
C (organic).....	.21	.37	.11	.24	none
	100.13	99.96	99.77	99.81	100.02

MONTANA.

Rocks collected by A. C. Peale. Analyses by Charles Catlett, record Nos. 890, 905.

A. North of East Gallatin River.

B. West of North Boulder River.

C, D. North of East Gallatin River.

E. Base of Carboniferous, west side of Bridger Range.

F. Middle Carboniferous, north of Gallatin River.

G, H. Upper Carboniferous, north of Gallatin River.

Analyses all partial. A, B, C, D, and H were published by Peale in Bull. 110, pp. 16, 28, and 40.

	A.	B.	C.	D.
Insoluble34	1.78	23.50	35.26
Fe ₂ O ₃ Al ₂ O ₃22	.40	2.50	1.92
CaCO ₃	54.54	54.54	67.85	59.11
MgCO ₃	43.63	42.62	6.18	1.96
	99.73	99.34	100.03	98.25

	E.	F.	G.	H.
Insoluble	9.98	5.99	50.74	25.24
Fe ₂ O ₃ , Al ₂ O ₃38	.58	.30	5.30
CaCO ₃	88.50	91.96	32.28	40.21
MgCO ₃95	1.35	13.91	25.25
	99.81	99.88	97.23	96.00

COLORADO.

1. DENVER BASIN.

Rocks described by Emmons in Mon. XXVII. Analyses made by L. G. Eakins, in the Denver laboratory.

A. Upper Wyoming limestone, Morrison.

B. Niobrara dolomite.

	A.	B.
Insoluble.....	5.32	12.01
Al ₂ O ₃53	.54
Fe ₂ O ₃38	.11
MnO.....	.49	.20
CaO.....	48.73	27.49
MgO.....	2.95	18.03
H ₂ O.....	.11	.61
P ₂ O ₅03	.03
CO ₂	41.71	41.40
	100.25	100.42

2. LEADVILLE DISTRICT.

Rocks described by Emmons in Mon. XII. Analyses A and E by W. F. Hillebrand; B, C, and D by A. Guyard; all made in the Denver laboratory.

A. Upper blue limestone, Silver Wave mine.

B. Upper blue limestone, Dugan quarry.

C. Upper blue limestone, Glass-Pendery mine.

D. Montgomery quarry, near base of blue limestone.

E. White limestone, upper part, Carbonate Hill quarry.

	A.	B.	C.	D.	E.
SiO ₂21	.70	.27	7.76	11.84
Al ₂ O ₃27	.17	.04	.11	1.66
Fe ₂ O ₃21	.11	.22	.10	1.51
FeO.....	.24	.38	.13	.57	.83
MnO.....	trace	.05	.20	.06	-----
CaO.....	30.79	30.43	29.97	27.26	26.60
MgO.....	21.14	20.78	21.52	20.05	17.41
K ₂ O.....	.03	.05	.01	.02	.02
Na ₂ O.....	.06	.09	.02	.04	.03
H ₂ O.....	.22	.04	.07	.05	.48
P ₂ O ₅	trace	.12	.03	.07	trace
CO ₂	46.84	46.93	47.39	43.79	40.01
SO ₃	trace	-----	-----	trace	-----
Cl.....	.10	.14	.04	.06	.05
FeS ₂	trace	trace	-----	trace	-----
Organic matter.....	.03	.03	.02	.07	-----
	100.14	100.02	99.93	100.01	100.44

3. FAIRPLAY, PARK COUNTY.

Analyses made by W. F. Hillebrand, in the Denver laboratory.

A. Serpentinous limestone, Buckskin Gulch.

B. Limestone, Fairplay.

C. Limestone, Mount Silverheels.

D. Dolomite-limestone, Mount Silverheels.

Analyses B, C, and D partial, with CO₂ calculated to satisfy bases.

	A.	B.	C.	D.
Insoluble		2. 37	. 51	1. 98
SiO ₂	17. 64			
Al ₂ O ₃ 99			
Fe ₂ O ₃ 62			
FeO 18	} . 19	} . 10	} . 46
MnO	trace			
CaO	32. 23	53. 64	55. 50	30. 19
MgO	19. 01	. 73	. 17	20. 47
Alkalies 07			
H ₂ O	3. 72	. 51		
P ₂ O ₅ 05			
CO ₂	25. 33	42. 93	43. 82	46. 52
Cl 08			
	99. 92	100. 37	100. 10	99. 62

4. PITKIN COUNTY.

Limestones and dolomites collected under the direction of S. F. Emmons. Analyses by George Steiger, record No. 1559. CO₂ calculated to satisfy bases. Analyses partial only.

	A.	B.	C.	D.	E.	F.	G.
Insoluble 16	. 80	1. 02	13. 63	1. 42	31. 12	7. 78
Fe ₂ O ₃ 22	1. 63	2. 10	1. 88	3. 34	. 36	. 88
FeO 09	. 23	. 06	. 64	. 42	. 19	. 22
CaO	30. 66	31. 19	33. 74	35. 98	31. 61	37. 28	38. 85
MgO	20. 94	19. 69	16. 76	8. 25	18. 06	. 54	9. 97
CO ₂	47. 13	46. 16	44. 94	37. 35	44. 70	29. 88	41. 47
	99. 20	99. 70	98. 62	97. 73	99. 55	99. 37	99. 17

The following samples from Aspen were analyzed by L. G. Eakins in the Denver laboratory. Partial analyses, CO₂ calculated as before.

H. Blue limestone.

I. Limestone.

J. Dolomite.

	H.	I.	J.
Insoluble52	.33	.84
Fe ₂ O ₃88	trace	1.31
CaO	31.16	55.81	30.46
MgO	20.64	.16	20.90
CO ₂	47.19	44.03	46.92
	100.39	100.33	100.43

5. GLENWOOD SPRINGS, GARFIELD COUNTY.

Limestones and dolomites collected under the direction of S. F. Emmons. Analyses, partial only, by George Steiger, record No. 1559. CO₂ calculated to satisfy bases.

	A.	B.	C.	D.	E.	F.	G.
Insoluble	21.45	47.74	6.47	3.71	9.44	17.82	1.96
Fe ₂ O ₃97	.18	.42	none	.26	.74	.03
FeO23	.71	.35	.55	.32	.57	.35
CaO	40.64	15.87	46.65	47.40	39.56	26.50	32.14
MgO73	10.60	2.64	4.49	8.56	14.86	18.72
CO ₂	32.73	24.13	39.55	42.15	40.52	37.18	45.85
	96.75	99.23	96.08	98.30	98.66	97.67	99.05

	H.	I.	J.	K.	L.	M.
Insoluble	2.27	.22	.23	.06	.22	.11
Fe ₂ O ₃14	trace	.09	none	.10	.03
FeO				none	.10	.07
CaO	53.79	55.17	55.49	55.81	55.45	55.68
MgO46	.21	.24	trace	.24	trace
CO ₂	42.76	43.58	43.87	43.85	43.84	43.75
	99.42	99.18	99.92	99.72	99.95	99.64

6. TENMILE DISTRICT, SUMMIT COUNTY.

Partial analyses, made in the Denver laboratory, by W. F. Hillebrand. CO_2 calculated to satisfy bases. Manganese and iron present as carbonates, but Fe_2O_3 and Al_2O_3 were not separated from them.

A. A. V. Fletcher shaft, Copper Mountain.

B. Pittston tunnel.

C. Middle Carboniferous, Pearl Hill.

D. Summit quarry.

E. Dolomite, Sheep Mountain.

F. Oolitic limestone, northwest corner of area.

G. Pittston tunnel.

	A.	B.	C.	D.	E.	F.	G.
Insoluble . . .	2.69	.62	10.09	1.75	.78	1.37	7.91
FeO , MnO . .	.21	.25	1.19	.32	1.50	.20	.32
CaO	54.23	55.24	28.01	53.60	30.55	55.17	50.83
MgO21	.24	18.33	1.23	20.15	.28	.70
CO_2	42.97	43.81	42.63	43.65	47.04	43.76	40.90
	100.31	100.16	100.25	100.55	100.02	100.78	100.66

H. Dolomite, Blackbird tunnel, Tucker Mountain.

I. Summit King shaft, Summit City.

J. Middle Carboniferous dolomite, Ptarmigan Hill.

K. Hill north of Sugar Loaf.

L. Quarry on southeast side of Searls Gulch.

M. Open cut below Sabbath Rest tunnel, Elk Mountain.

N. Triassic, Jacque Mountain.

	H.	I.	J.	K.	L.	M.	N.
Insoluble . . .	2.68	6.75	.65	4.42	.36	.82	2.04
FeO , MnO . .	1.52	3.08	1.67	.10	.17	.07	.15
CaO	31.60	28.05	30.90	52.97	55.58	55.47	54.62
MgO	18.27	18.15	19.75	.40	.37	.22	.25
CO_2	45.75	43.88	47.02	42.12	44.17	43.86	43.28
	99.82	99.91	99.99	100.01	100.65	100.44	100.34

UTAH.

A. Marble from the Ontario mine, east end of the 1,000-foot level. Analysis made by L. G. Eakins, in the Denver laboratory.

B. Oolitic sand from shore of Great Salt Lake. Analysis by T. M. Chatard, record No. 156.

	A.	B.
Insoluble in HCl.....	9.61	4.03
Al ₂ O ₃	trace	} .20
Fe ₂ O ₃		
CaO.....	50.63	51.33
MgO.....	.09	.72
Alkalies.....		.63
CO ₂	<i>a</i> 39.89	41.07
P ₂ O ₅		trace
SO ₃89
H ₂ O.....		.83
Organic matter.....		.27
	100.22	99.97

a Calculated to satisfy bases.

NEVADA AND CALIFORNIA.

A. Crystalline limestone, Eureka, Nevada. Analysis by E. A. Schneider, record No. 1279.

B. From base of the Hamburg limestone, Eureka district, Nevada.

C. From summit of Hamburg limestone, Eureka.

D. Pogonip limestone (Silurian), Eureka district. Analyses B, C, and D made by W. F. Hillebrand in the Denver laboratory. Described by Hague in Mon. XX, pp. 40, 49. In D the CO_2 was taken by difference.

E. Cretaceous limestone from Mount Diablo, California. Analysis made by W. H. Melville in the San Francisco laboratory and published in Bull. Geol. Soc. Amer., vol. 2, p. 409. FeO and alkalies undetermined.

F. White deposit, White Terrace, west shore of Pyramid Lake, Nevada. Analysis by T. M. Chatard, record No. 34.

	A.	B.	C.	D.	E.	F.
Insoluble53					
SiO_2		24.00	3.94	9.34	21.19	22.00
Al_2O_312	.64	.31	.39	5.14
Fe_2O_312	.43	.29	1.52	2.04
FeO20			
MnO61		3.61	
CaO	30.60	41.97	51.96	50.01	35.61	37.22
MgO	21.69	.80	.52	.54	1.39	1.89
Alkalies		traces	traces	traces		
H_2O at 100°		} .16	} .37	} .13	.76	} 3.32
H_2O above 100°					2.33	
P_2O_507	.50	.24	2.55	
CO_2	47.13	32.62	40.71	39.11	26.84	28.53
Organic matter		traces	.03	traces		
Cl01	.01	.03		
	99.95	99.87	99.92	100.00	96.19	100.14

HAWAIIAN ISLANDS.

Coral and shell rocks, analyzed for N. S. Shaler by L. G. Eakins, record Nos. 886, 887, 889. Analyses only partial.

- A. Laie.
- B, C. Kohuku Bluff.
- D. Kohuku coral flat.
- E. Point near coral flat.
- F. "Modern chalk," Oahu.
- G. Diamond Head.
- H. Under lava, Honolulu.
- I. Old reef, Waialua.
- J. Campbell's ranch, Waianea, Oahu.
- K. Wailuku Bay.
- L. Reef No. 3, Honolulu.
- M. Prison Knoll, Honolulu.

	A.	B.	C.	D.	E.	F.	G.
SiO ₂ 35	. 19	. 67	. 25	. 26	33. 25	2. 97
Al ₂ O ₃ 19	. 52	. 73	. 49	. 21	19. 53	2. 88
Fe ₂ O ₃						10. 71	
CaO.....	49. 38	49. 34	51. 09	53. 34	52. 17	11. 37	44. 82
MgO.....	1. 74	4. 60	2. 50	. 67	1. 51	3. 06	5. 32
CO ₂	41. 89	44. 33	43. 64	43. 89	43. 95	11. 09	40. 81
H ₂ O.....	4. 74	. 40	. 79	. 93	. 70	9. 84	1. 86
	98. 29	99. 38	99. 42	99. 57	98. 80	98. 85	98. 66

	H.	I.	J.	K.	L.	M.
SiO ₂	5. 34	1. 05	. 53	. 45	3. 53	. 81
Al ₂ O ₃	5. 11	1. 26	. 62	1. 82	2. 26	1. 19
Fe ₂ O ₃						
CaO.....	42. 24	51. 07	50. 69	50. 54	46. 52	52. 67
MgO.....	5. 95	. 11	2. 98	1. 83	2. 45	. 42
CO ₂	38. 71	42. 68	43. 96	42. 80	40. 59	42. 81
H ₂ O.....	1. 61	1. 33	. 46	1. 93	2. 75	1. 24
	98. 96	97. 50	99. 24	99. 37	98. 10	99. 14

SLATES AND SHALES.

VERMONT.

Samples A to I, inclusive, are described by T. Nelson Dale in 19th Ann., Part III. Analyses by W. F. Hillebrand, record Nos. 1567, 1656. Roofing slates of Cambrian age.

A. McCarty quarry, South Poultney.

B. Unfading green, Eureka quarry, Poultney.

C. Sea green, Griffith and Nathaniel quarry, South Poultney.

D. Sea green, Rising and Nelson's quarry, Pawlet.

E. Sea green, Brownell quarry, Pawlet.

F. Black, American Black Slate Company, Benson.

	A.	B.	C.	D.	E.	F.
SiO ₂	61.63	59.27	62.37	67.76	59.84	59.70
Al ₂ O ₃	16.33	18.81	15.43	14.12	15.02	16.98
Fe ₂ O ₃	4.10	1.12	1.34	.81	1.23	.52
FeO.....	2.71	6.58	5.34	4.71	4.73	4.88
MgO.....	2.92	2.21	3.14	2.38	3.41	3.23
CaO.....	.50	.42	.77	.63	2.20	1.27
Na ₂ O.....	1.26	1.88	1.14	1.39	1.12	1.35
K ₂ O.....	5.54	3.75	4.20	3.52	4.48	3.77
H ₂ O at 100°...	.31	.32	.34	.23	.41	.30
H ₂ O above 100°	3.24	3.98	3.71	2.98	3.44	3.82
TiO ₂68	.99	.74	.71	.74	.79
P ₂ O ₅16	.11	.06	.07	.09	.16
MnO.....	.09	.13	.22	.10	.34	.16
BaO.....	.06	.05	.07	.04	.09	.08
CO ₂41	.21	.87	.40	2.98	1.40
FeS ₂04	.15	.06	.22	.05	1.18
C.....	none	none	trace	none	trace?	.46
	99.98	99.98	99.80	100.07	100.17	100.05

All six contain traces of lithia, of sulphates, and of nickel or cobalt.

- G. Unfading green, Valley Slate Company's quarry, Poultney.
H. Mottled, purple and green, Eureka quarry, Poultney.
I. Purple, 1 mile south of Hydeville, in Castleton.

	G.	H.	I.
SiO ₂	59.48	60.24	60.96
Al ₂ O ₃	18.22	18.46	16.15
Fe ₂ O ₃	1.24	2.56	5.16
FeO	6.81	5.18	2.54
MgO	2.50	2.33	3.06
CaO56	.33	.71
Na ₂ O	1.55	1.57	1.50
K ₂ O	3.81	4.09	5.01
H ₂ O at 110°17	.18	.17
H ₂ O above 110°	4.05	3.81	3.08
TiO ₂	1.02	.92	.86
P ₂ O ₅10	.11	.23
MnO07	.07	.07
BaO05	.03	.04
CO ₂39	.08	.68
FeS ₂13	.16	none
F08	undet.	undet.
N, as NH ₃03	.03	.01
	100.26	100.15	100.23

All three contain traces of lithia, nickel, and chlorine, and possibly of zirconia; in G and H traces of strontia, but none in I.

J. Slate from Guilford, collected for the Educational Series of Rock Specimens. Analysis by L. G. Eakins, record No. 1316.

K. Slate from the Lakeshore quarry, Hydeville. Analysis by Eakins, record No. 1159.

	J.	K.
SiO ₂	60.72	58.15
Al ₂ O ₃	22.59	18.93
Fe ₂ O ₃		2.91
FeO	6.03	5.64
MgO	2.05	2.70
CaO41	.60
Na ₂ O86	1.17
K ₂ O	3.69	3.92
H ₂ O	3.01	4.56
TiO ₂93
P ₂ O ₅13	.12
MnO	trace	.07
SO ₃16
C57	
	100.06	99.86

NEW YORK.

Roofing slates from Washington County, collected by T. Nelson Dale. Of Cambrian age. Described by Dale in 19th Ann., Part III.

A. Red, three-fourths mile south of Hampton Village.

B. Empire Red Slate Company, near Granville.

C. National Red Slate Company, Granville.

D. Green, three-fourths mile northwest of Janesville.

Analyses by W. F. Hillebrand, record No. 1567.

	A.	B.	C.	D.
SiO ₂	67.61	67.55	56.49	67.89
Al ₂ O ₃	13.20	12.59	11.59	11.03
Fe ₂ O ₃	5.36	5.61	3.48	1.47
FeO.....	1.20	1.24	1.42	3.81
MgO.....	3.20	3.27	6.43	4.57
CaO.....	.11	.26	5.11	1.43
Na ₂ O.....	.67	.61	.52	.77
K ₂ O.....	4.45	4.13	3.77	2.82
H ₂ O at 110°.....	.45	.40	.37	.36
H ₂ O above 110°.....	2.97	3.03	2.82	3.21
TiO ₂56	.58	.48	.49
P ₂ O ₅05	.10	.09	.10
MnO.....	.10	.19	.30	.16
BaO.....	.04	.31	.06	.04
CO ₂	none	.11	1.42	1.89
FeS ₂03	.04	.03	.04
	100.00	100.02	100.38	100.08

All contain traces of lithia, of sulphates, and of nickel or cobalt. No carbonaceous matter is present.

E. Red slate, from quarry 3 miles north of Raceville.

F. Green spot in E.

G. Purple rim of green spot F.

Analyses by W. F. Hillebrand, record No. 1656.

	E.	F.	G.
SiO ₂	63.88	65.44	64.59
Al ₂ O ₃	9.77	9.38	10.23
Fe ₂ O ₃	3.86	1.09	1.79
FeO.....	1.44	1.06	1.19
MgO.....	5.37	4.92	5.12
CaO.....	3.53	4.53	4.07
Na ₂ O.....	.20	.22	.23
K ₂ O.....	3.45	3.57	3.70
H ₂ O at 110°.....	.27	.25	.28
H ₂ O above 110°.....	2.48	2.10	2.29
TiO ₂47	.52	.51
P ₂ O ₅08	.08	.08
MnO.....	.21	.32	.26
BaO.....	.05	.06	.05
CO ₂	5.08	6.55	5.84
FeS ₂	trace	.04	trace
	100.14	100.13	100.23

Contain traces of lithia and nickel. Fluorine not determined.

KENTUCKY, GEORGIA, ALABAMA.

A. Indurated Carboniferous shale, in contact with the peridotite dike of Elliott County, Kentucky. Described by Diller in Bull. 38. Analysis by T. M. Chatard, record No. 351.

B. Fragment of shale included in the Elliott County dike. Analysis by Chatard, record No. 353.

C. Bituminous shale, Dry Gap, Georgia. Analysis by L. G. Eakins, record No. 1316. P. R. C. 22. Described by Diller in Bull. 150, p. 90.

D. Middle Cambrian shale, Coosa Valley, near Blaine, Cherokee County, Alabama. Analysis by H. N. Stokes, record No. 1549.

	A.	B.	C.	D.
SiO ₂	41.32	35.53	51.03	55.02
Al ₂ O ₃	20.71	18.23	13.47	21.02
Fe ₂ O ₃	2.59	2.46	8.06	5.00
FeO.....	5.46	4.81	-----	1.54
MgO.....	1.91	2.01	1.15	2.32
CaO.....	9.91	21.17	.78	1.60
Na ₂ O.....	7.19	2.53	.41	.81
K ₂ O.....	.88	1.08	3.16	3.19
H ₂ O at 110°.....	-----	1.40	} .81	2.44
H ₂ O above 110°.....	8.78	9.00		5.65
TiO ₂48	.95	-----	.65
P ₂ O ₅08	.08	.31	.06
MnO.....	.17	.13	-----	trace
BaO.....	-----	-----	-----	.04
SrO.....	-----	-----	-----	trace
Li ₂ O.....	-----	-----	-----	.03
SO ₃	-----	-----	-----	.02
S.....	-----	-----	7.29	-----
Cl.....	-----	-----	-----	trace
CO ₂55	.88	-----	.83
Carbonaceous matter.....	-----	-----	-----	.32
Fixed carbon.....	-----	-----	13.11	-----
Volatile hydrocarbons.....	-----	-----	3.32	-----
	100.03	100.26	102.90	100.54
Less O = S.....	-----	-----	2.74	-----
			100.16	

OHIO.

Three samples of Utica shale from New Vienna. Collected by Edward Orton. Partial analyses by F. W. Clarke and R. B. Riggs, record No. 731.

	A.	B.	C.
Insoluble.....	60. 17	29. 51	25. 80
CaO.....	17. 11	33. 43	35. 27
MgO.....	1. 25	2. 16	1. 32
CO ₂	15. 24	27. 16	27. 40
	93. 77	92. 26	89. 79

Iron and alumina are present in the soluble portions of these shales, the solvent being dilute hydrochloric acid.

MICHIGAN.

Clay slate, sec. 17, T. 43 N., R. 31 W., near Mansfield. Contains principally quartz, white mica, actinolite, rutile, hematite, and carbonaceous matter. Described by J. M. Clements in Mon. XXXVI, pp. 59, 61, 210. Analysis by George Steiger, record No. 1709.

SiO ₂	60. 28
Al ₂ O ₃	22. 61
Fe ₂ O ₃	2. 53
FeO.....	. 45
MgO.....	1. 35
CaO.....	. 13
Na ₂ O.....	. 54
K ₂ O.....	5. 73
H ₂ O at 100°.....	. 60
H ₂ O above 100°.....	3. 62
TiO ₂ 69
P ₂ O ₅ 03
MnO.....	trace
BaO.....	. 04
C.....	. 97
	<hr/> 99. 57

WISCONSIN.

Slates of the Penokee-Gogebic series, collected by C. R. Van Hise. A and C are described in Mon. XIX, p. 306, as magnetitic clay slates. Analyses by L. G. Eakins, record No. 392.

- A. Sec. 6, T. 45 N., R. 2 E.
- B. Sec. 1, T. 45 N., R. 1 E.
- C. Sec. 4, T. 44 N., R. 2 W.

	A.	B.	C.
SiO ₂	53.44	59.73	52.58
Al ₂ O ₃	19.62	22.78	20.76
Fe ₂ O ₃	11.38	.11	12.17
FeO.....	5.35	5.98	4.08
MgO.....	1.58	2.94	1.33
CaO.....	.42	.53	.30
Na ₂ O.....	2.61	1.41	.37
K ₂ O.....	1.73	3.48	4.87
H ₂ O.....	4.07	3.28	3.43
P ₂ O ₅	trace	-----	-----
MnO.....	trace	.09	.21
Li ₂ O.....	trace	-----	trace
	100.20	100.33	100.10

COLORADO.

Shales from the Pueblo quadrangle, collected by G. K. Gilbert.

A, B. Near Nushbaum Spring.

C. Salt Creek.

D. Head of Rock Creek.

E. Near Rush Creek.

Analyses by George Steiger, record No. 1466.

	A.	B.	C.	D.	E.
SiO ₂	60.80	51.69	60.60	63.60	45.89
Al ₂ O ₃	15.63	16.50	16.42	16.74	13.24
Fe ₂ O ₃	4.62	7.90	4.95	4.63	3.88
MgO.....	2.73	2.10	1.43	1.19	2.12
CaO.....	1.63	4.41	1.61	.68	12.09
Na ₂ O.....	1.45	2.07	.92	.29	.47
K ₂ O.....	2.55	2.68	2.98	2.92	2.31
H ₂ O at 100°.....	3.19	3.02	3.91	2.88	1.38
H ₂ O above 100°.....	4.16	6.00	5.72	5.99	4.16
TiO ₂47	.66	.35	.66	.52
P ₂ O ₅10	.22	.31	.16	.17
CO ₂		3.19			10.38
Organic matter.....	2.87	.53	.84	.46	3.47
	100.20	100.97	100.04	100.20	100.08

Calcareous shales from Fairplay, Park County. Partial analyses, by W. F. Hillebrand, made in the Denver laboratory.

	F.	G.
Insoluble.....	68.72	35.14
Fe ₂ O ₃ , Al ₂ O ₃ , etc.....	2.10	
FeO, MnO.....		2.10
MgO.....	5.72	12.55
CaO.....	9.06	19.34
H ₂ O.....	1.01	.73
CO ₂ , calculated.....	13.41	30.28
	100.02	100.14

CALIFORNIA.

Cretaceous shales from Mount Diablo. Described by Turner and Melville, Bull. Geol. Soc. Amer., vol. 3, pp. 383-414. Analyses by W. H. Melville, made in the laboratory at San Francisco, except F (record No. 1166), which was done in the Washington laboratory.

A. Brownish black, resinous. From Bagley Canyon.

B. Slate colored, soft, friable, little altered. From near Bagley Creek.

C. Same locality as B, less friable, but considerably altered.

D. Slate colored, friable. From Arroyo del Cerro.

E. Very friable. Same locality as D.

	A.	B.	C.	D.	E.
SiO ₂	56.66	53.65	49.14	25.05	40.17
Al ₂ O ₃	17.64	17.64	16.91	8.28	12.76
Fe ₂ O ₃49	4.06	4.39	.27	2.10
FeO.....	5.22	3.72	3.82	2.41	3.56
MgO.....	3.50	5.15	5.43	2.61	15.42
CaO.....	1.67	2.27	3.28	27.87	4.24
Na ₂ O.....	2.17	2.53	4.67	undet.	.57
K ₂ O.....	2.27	2.22	1.53	undet.	1.36
H ₂ O at 100°.....	3.01	3.95	3.39	1.44	9.19
H ₂ O above 100°.....	5.92	4.57	6.97	2.86	6.73
P ₂ O ₅15	.23	.24	.08	.08
NiO.....		trace	trace	trace	trace
MnO.....	.19	.01	.22	4.11	.16
CO ₂				24.20	3.48
SO ₃93				
	99.82	100.00	99.99	99.18	99.82

F. Neocomian shale, altered, light brown, friable. From near Arroyo del Cerro.

G. Calcareous shale, near Arroyo del Cerro. Hard, compact, dark colored. Very much altered.

H. Red shale, metamorphic area at head of Bagley Creek.

I. Silicified shale or phthanite, same locality as H.

J. Clay slate, near the head of Yaqui Gulch, in Mariposa County. Described by Turner in Bull. 150, p. 342. Contains grains of quartz and feldspar, abundant carbonaceous particles, a chloritic substance (?), and a fibrous alteration of sillimanite (?). Analysis by George Steiger, record No. 1643.

	F.	G.	H.	I.	J.
SiO ₂	45.64	44.56	69.98	93.54	60.35
Al ₂ O ₃	15.42	3.12	11.69	2.26	17.62
Fe ₂ O ₃	3.40	1.27	6.23	.48	5.64
FeO.....	3.73	5.21	1.08	.79	2.20
MgO.....	4.62	3.39	1.29	.66	1.04
CaO.....	8.11	12.70	.38	.09	.45
Na ₂ O.....	3.13	3.09	.73	.37	1.00
K ₂ O.....	1.86	.88	3.72	.51	3.16
H ₂ O at 100°.....	8.74	1.41	1.03	.21	1.02
H ₂ O above 100°.....		6.24	2.92	.72	4.36
TiO ₂75
P ₂ O ₅27	.16	.05		.17
Cr ₂ O ₃12				
MnO.....	.33	trace	.49	.23	none
BaO.....					.12
CO ₂	4.59	17.62			none
SO ₃05
Cl.....					.01
F.....					trace
C.....					1.72
	99.96	99.65	99.59	99.86	99.76

CLAYS, SOILS, ETC.

MASSACHUSETTS.

Clays and soils from Marthas Vineyard, collected by N. S. Shaler. See 7th Ann., p. 303. Analyses by F. W. Clarke, record Nos. 439, 440, 441, 442, 443, 444, 445, 446, 454, and 455. Partial analyses only.

- A. Average sample of white clay, east end of Chilmark Cliffs.
 B. Average sample of clays, Weyquosque series, Chilmark Cliffs.
 C. Average sample of fine clay and soil, east end of Weyquosque Cliffs.

D. Sandy white clay, south end of Gay Head Cliffs.

E. Average sample of fine white clay, south end of Gay Head Cliffs.

	A.	B.	C.	D.	E.
SiO ₂	82.95	61.76	70.81	56.19	73.46
Al ₂ O ₃ , Fe ₂ O ₃	13.45	25.35	20.67	30.65	19.06
MgO.....	trace	1.95	1.99	trace	trace
CaO.....	none	.51	trace	none	none
Na ₂ O.....		1.83	1.23	-----	.70
K ₂ O.....		3.01	1.67	-----	.73
Ignition.....	3.47	5.76	3.39	10.79	6.36
P ₂ O ₅	none	trace	none	none	none
SO ₃			-----	2.45	none
	99.87	100.17	99.76	100.08	100.31

F. Average sample of clay, north end of Gay Head Cliffs.

G. Average sample of southernmost red clays, Gay Head.

H. Brown clay, south of light-house, Gay Head Cliffs.

I. Average sample of red clay from the greensand, north end of Gay Head.

J. Pyritiferous clay, central part of Gay Head section.

	F.	G.	H.	I.	J.
SiO ₂	49.19	57.50	56.62	55.93	72.74
Al ₂ O ₃ , Fe ₂ O ₃	39.77	31.21	31.24	33.51	21.46
MgO.....	trace	.20	1.97	.19	trace
CaO.....	none	.19	trace	none	none
Na ₂ O.....		.40	.40	undet.	-----
K ₂ O.....			2.76	undet.	-----
Ignition.....	11.47	9.83	7.57	9.98	5.69
P ₂ O ₅	none	none	none	none	none
	100.43	99.33	100.56	99.61	99.89

NEW YORK, PENNSYLVANIA, DELAWARE.

A. Clay, near Richfield Springs, New York. Partial analysis by Charles Catlett, record No. 946.

B, C. Clays, Northumberland County, Pennsylvania. Analyses by Charles Catlett, record No. 952.

D. Kaolin, Hockessin, Delaware. P. R. C. 149.

E. Portion of D insoluble in sulphuric acid.

F. Portion of D soluble in sulphuric acid.

Analyses D, E, and F by George Steiger, record No. 1626.

	A.	B.	C.	D.	E.	F.
SiO ₂	49.65	65.97	59.16	48.73	29.55	19.18
Al ₂ O ₃	23.82	20.37	18.68	37.02	18.44	18.58
Fe ₂ O ₃		2.75	10.32	.79	.27	.52
MgO.....	trace	.52	.67	.11	trace	.11
CaO.....	6.48	.64	.52	.16	.02	.14
Na ₂ O.....	undet.	.05	.11	.04	.02	.02
K ₂ O.....	undet.	3.32	3.35	.41	.41	none
H ₂ O at 100°...	16.18	6.28	6.87	.52	6.84	5.99
H ₂ O above 100°				12.83		
TiO ₂17	.11	.06
P ₂ O ₅03	undet.	undet.
	96.13	99.90	99.68	100.81	55.66	44.60

MARYLAND.

Clays from the Matawan formation, received from W. B. Clark.
Analyses by George Steiger, record No. 1684.

A. Below Barnard's wharf, near Betterton, Kent County.

B. Severn River, below Round Bay, Anne Arundel County.

C. Magothy River near Wilson's wharf, Anne Arundel County.

D. Fort Washington Bluff.

	A.	B.	C.	D.
SiO ₂	73.47	87.15	82.86	73.02
Al ₂ O ₃ <i>a</i>	12.69	6.46	6.49	10.00
Fe ₂ O ₃	4.62	2.15	3.54	4.78
MgO59	.27	.52	.90
CaO15	.10	.29	.57
Na ₂ O09	.14	.16	.59
K ₂ O	1.55	.90	1.16	1.92
H ₂ O at 100°89	.32	.69	1.09
H ₂ O above 100°	3.85	1.90	2.24	3.07
SO ₃19	none	.21	1.04
	98.09	99.39	98.16	96.98

a Titanic and phosphoric oxides not separated.

No carbonates present. Sulphides undetermined.

VIRGINIA, NORTH CAROLINA.

A. Residual clay from decay of Trenton limestone, Lexington, Virginia. Described by Russell in Bull. 52. Analysis by R. B. Riggs, record No. 373. See also analysis of the limestone.

B. Residual clay from limestone, Staunton, Virginia.

C. Portion of B soluble in weak hydrochloric acid.

D. Insoluble portion of B. Analyses B, C, and D by George Steiger, record No. 1630. See also analysis of limestone.

E. Decomposed dolerite, near Wadesboro, North Carolina. Described by Russell in Bull. 52. Analysis by T. M. Chatard, record No. 327.

F. Residual clay from decay of chloritic schist, Cary, 8 miles west of Raleigh, North Carolina. Analysis by R. B. Riggs, record No. 364. Described by Russell in Bull. 52.

	A.	B.	C.	D.	E.	F.
SiO ₂	43.07	55.90	3.09	52.81	39.55	54.54
Al ₂ O ₃	25.07	19.92	3.96	15.96	28.76	26.43
Fe ₂ O ₃	15.16	7.30	6.25	1.05	16.80	9.04
FeO.....		.39	.30	.09		
MgO.....	.03	1.18	.43	.75	.59	
CaO.....	.63	.50	.30	.20	.37	
Na ₂ O.....	1.20	.23	.20	.03	undet.	
K ₂ O.....	2.50	4.79	.28	4.51	undet.	
H ₂ O at 110°....	12.98	2.54			13.26	9.87
H ₂ O above 110°		6.52	2.10	4.42		
TiO ₂20	.04	.16	.64	
P ₂ O ₅10	.04	.06	.10	
Cr ₂ O ₃					trace	
MnO.....		none			trace	
CO ₂38	.38	none		
	100.64	99.95	17.37	80.04	100.07	99.88

SOUTH CAROLINA, GEORGIA.

A, B, C. Clays, near Augusta, Georgia. Partial analyses by George Steiger, record No. 1395.

D. Kaolin, Aiken, South Carolina. Analysis by Steiger, No. 1472.

	A.	B.	C.	D.
SiO ₂	60.24	61.36	60.70	44.94
Al ₂ O ₃	26.72	29.04	29.24	39.18
Fe ₂ O ₃				
MgO	present	present	present	.52
CaO88	.76	.68	-----
Alkalies	undet.	undet.	undet.	-----
H ₂ O at 100°	6.28	7.46	6.74	.47
H ₂ O at 200°20
H ₂ O at 300°27
H ₂ O, ignition				13.38
TiO ₂	-----	-----	-----	.65
P ₂ O ₅	-----	-----	-----	.12
	94.12	98.62	97.36	99.73

FLORIDA.

A. Hammock clay, Melborne Creek. Collected by N. S. Shaler. Partial analysis by L. G. Eakins, record No. 881.

B. Clay, Tampa.

C. Clay, Lakeland.

B and C collected by W. H. Dall. Analyses by L. G. Eakins, record No. 1255, partial.

	A.	B.	C.
SiO ₂	38.04	70.78	80.39
Al ₂ O ₃ , Fe ₂ O ₃	27.19	11.33	15.03
MgO46	-----	-----
CaO	10.73	2.18	1.22
H ₂ O	^a 23.61	14.55	4.34
	100.03	98.84	100.98

^a Includes some CO₂.

Clays collected by G. H. Eldridge.

D, E. From the Sandlin place, 2 miles southeast of Marion, Hamilton County.

F. From Richmond's, 6 miles south of Leesburg, Lake County.

G, H. From Bartow Junction.

Analyses D, E, and F by H. N. Stokes, record No. 1493; G and H by George Steiger, No. 1545.

	D.	E.	F.	G.	H.
SiO ₂	15.68	78.23	84.41	79.99	79.48
Al ₂ O ₃61	7.30	11.02	10.82	12.14
Fe ₂ O ₃45	1.85	trace	3.25	2.64
FeO.....				.25	.09
MgO.....	17.28	2.11	trace	.07	.07
CaO.....	26.11	1.60	.20	.23	.31
H ₂ O at 100°.....	} 1.97	} a 8.48	} a 4.25	.90	.86
H ₂ O above 100°.....				4.09	4.73
P ₂ O ₅	trace	trace	trace		
CO ₂	37.90			none	none
	100.00	99.57	99.88	99.87	100.32

a Includes a little CO₂.

I. "Filtering clay," Ocala. Received from D. T. Day. Analysis by H. N. Stokes, record No. 1738.

SiO ₂	36.73
Al ₂ O ₃	27.78
Fe ₂ O ₃	3.21
MgO.....	.64
CaO.....	.81
Na ₂ O.....	none
K ₂ O.....	.42
H ₂ O at 110°.....	7.38
H ₂ O above 110°.....	12.14
TiO ₂	1.27
P ₂ O ₅	5.54
CO ₂	none
Organic matter.....	3.61
	99.53

ALABAMA, MISSISSIPPI.

A. Kaolin, Greenville, Alabama. Contains about 40 per cent of kaolin, with fragments of quartz, feldspar, and mica. Analysis by T. M. Chatard, record No. 1148.

B. Residual clay from decay of Knox dolomite, Morrisville, Alabama. Described by Russell in Bull. 52. Analysis by W. F. Hillebrand, record No. 797. See also analysis of the dolomite.

C. Loess from Vicksburg, Mississippi. Described by Chamberlin and Salisbury, 6th Ann., p. 282. Analysis by R. B. Riggs, record No. 294.

	A.	B.	C.
SiO ₂	69.84	55.42	60.69
Al ₂ O ₃	19.91	22.17	7.95
Fe ₂ O ₃90	8.30	2.61
FeO.....		trace	.67
MgO.....	.28	1.45	4.56
CaO.....	.07	.15	8.96
Na ₂ O.....	.21	.17	1.17
K ₂ O.....	2.14	2.32	1.08
H ₂ O at 110°.....	.06	2.10	
H ₂ O, ign.....	6.72	7.76	1.14
TiO ₂52
P ₂ O ₅13
MnO.....	trace		.12
CO ₂			9.63
C, organic.....			.19
SO ₃12
Cl.....			.08
	100.13	99.84	99.62

ILLINOIS, IOWA, MINNESOTA.

A, B. Clays from Henry County, Illinois. Analyses by T. M. Chatard, record No. 144.

C. Loess, a stratum overlying residuary clay, 350 feet above the Mississippi River, near Galena, Illinois. Described by Chamberlin and Salisbury, 6th Ann., p. 282. Analysis by R. B. Riggs, record No. 293. Dried at 100°.

D. Loess, 300 feet above the Mississippi, 3½ miles northwest of Dubuque, Iowa. Described by Chamberlin and Salisbury (*l. c.*), and analyzed by Riggs, No. 292. Dried at 100°.

E. Tallow clay, lead mine at Lansing, Iowa. Collected by W. P. Jenney. Analysis by H. N. Stokes, record No. 1337. Dried at 100°. Partial analysis.

F. Greenish-gray clay, New Ulm, Minnesota. Analysis by T. M. Chatard, record No. 825.

	A.	B.	C.	D.	E.	F.
SiO ₂	46.12	42.58	64.61	72.68	52.08	61.32
Al ₂ O ₃	15.24	12.16	10.64	12.03	23.11	12.27
Fe ₂ O ₃	4.41	3.90	2.61	3.53	9.34	3.62
FeO.....			.51	.96		4.18
MgO.....	3.63	4.32	3.69	1.11	2.12	1.76
CaO.....	8.63	11.33	5.41	1.59	1.04	.99
Na ₂ O.....	1.54	1.96	1.35	1.68	undet.	.42
K ₂ O.....	3.79	3.88	2.06	2.13	undet.	3.59
H ₂ O.....	15.57	18.64	2.05	2.50	9.80	10.73
TiO ₂79	.64	.40	.72		.66
P ₂ O ₅08	.10	.06	.23		.27
MnO.....	.28	.09	.05	.06		.27
ZnO.....					trace	
PbO.....					trace	
BaO.....						.05
CO ₂			6.31	.39		
C, organic.....			.13	.09		
SO ₃11	.51		.19
Cl.....			.07	.01		
	100.08	99.60	100.06	100.22	97.49	100.32

WISCONSIN.

Clays, etc., described by Chamberlin and Salisbury in 6th Ann., pp. 250 and 282. Analyses by R. B. Riggs, record Nos. 259, 260, 261, 262, 290, 295. Dried at 100°.

A. Residuary clay from Dodgeville, $4\frac{1}{2}$ feet below surface.

B. The same, $8\frac{1}{2}$ feet below surface.

C. Residuary clay from near Cobb, $4\frac{1}{2}$ feet below surface.

D. Same as C, $3\frac{1}{2}$ feet below surface.

E. Red, putty-like clay, containing pebbles, Milwaukee.

F. Red pebble clay, Milwaukee.

	A.	B.	C.	D.	E.	F.
SiO ₂	71.13	49.59	49.13	53.09	40.22	48.81
Al ₂ O ₃	12.50	18.64	20.08	21.43	8.47	7.54
Fe ₂ O ₃	5.52	17.19	11.04	8.53	2.83	2.53
FeO.....	.45	.27	.93	.86	.48	.65
MgO.....	.38	.73	1.92	1.43	7.80	7.05
CaO.....	.85	.93	1.22	.95	15.65	11.83
Na ₂ O.....	2.19	.80	1.33	1.45	.84	.92
K ₂ O.....	1.61	.93	1.60	.83	2.36	2.60
H ₂ O.....	4.63	10.46	11.72	10.79	1.95	2.02
TiO ₂45	.28	.13	.16	.35	.45
P ₂ O ₅02	.03	.04	.03	.05	.13
MnO.....	.04	.01	.06	.03	trace	.03
CO ₂43	.30	.39	.29	18.76	15.47
C, organic ..	.19	.34	1.09	.22	.32	.38
SO ₃	-----	-----	-----	-----	.13	.05
Cl.....	-----	-----	-----	-----	.06	.04
	100.39	100.50	100.68	100.09	100.27	100.50

MISSOURI, ARKANSAS.

A. Typical loess, Kansas City, Missouri. Dried at 100°. Described by Chamberlin and Salisbury, 6th Ann., p. 282. Analysis by R. B. Riggs, record No. 291.

B, C, D, E. Tallow clays, Joplin, Missouri. Collected by W. P. Jenney. Analyses by T. M. Chatard, record No. 1210.

F. Tallow clay, Aurora, Missouri. Collected by Jenney. Analysis by Chatard, No. 1210. In analyses B, C, D, E, and F the percentages of bases relate to the portion soluble in hydrochloric acid. Analyses only partial.

	A.	B.	C.	D.	E.	F.
Insoluble		40.64	43.07	39.34	39.62	34.04
SiO ₂	74.46					
Al ₂ O ₃	12.26	5.72	7.60	6.17	6.45	10.01
Fe ₂ O ₃	3.25	1.30	1.12	1.16	1.53	3.62
FeO12					
MgO	1.12	.27	.32	.27	.30	.25
CaO	1.69	1.80	1.70	2.13	1.77	2.09
Na ₂ O	1.43					
K ₂ O	1.83					
H ₂ O	2.70	17.19	16.74	17.63	16.95	16.96
TiO ₂14					
P ₂ O ₅09					
MnO02					
ZnO		32.46	29.43	34.28	33.55	33.49
CO ₂49					
C, organic ..	.12					
SO ₃06					
Cl05					
	99.83	99.38	99.98	100.98	100.17	100.46

The following partial analyses by H. N. Stokes, record No. 1260, all relate to tallow clays collected by W. P. Jenney. The same remarks apply as to B, C, D, E, and F.

G. Cave Springs mine, Jasper County, Missouri.

H. Great Western mine, Granby, Missouri.

I, J, K. Woodcock mine, Granby, Missouri.

L. Coon Hollow, Boone County, Arkansas.

Material dried at 103°.

	G.	H.	I.	J.	K.	L.
Insoluble	34.89	11.25	2.41	16.17	3.85	18.18
Soluble SiO ₂ ..	16.75	32.89	36.71	28.62	37.08	29.02
Al ₂ O ₃	7.38	10.78	8.21	8.93	6.46	6.34
Fe ₂ O ₃	10.34	3.89	2.75	5.98	3.49	4.40
ZnO	14.35	29.54	38.59	26.23	38.90	30.50
CaO	1.55	2.65	2.77	2.01	2.56	1.91
MgO35	.90	.78	.46	.42	.75
Ignition	10.37	8.22	7.99	9.19	7.52	8.36
	95.98	100.12	100.21	97.59	100.28	99.46

COLORADO.

A. Loess, Denver.

B. Loess, Highland.

C. Concretion in loess, Wray.

A, B, and C collected by S. F. Emmons, analyses by L. G. Eakins, record No. 1066.

D. Clay, Davis ranch, Pueblo quadrangle.

E. Clay, head of Rock Creek, Pueblo quadrangle.

D and E collected by G. K. Gilbert, analyses by George Steiger, record No. 1457.

	A.	B.	C.	D.	E.
SiO ₂	69.27	60.97	70.63	63.52	76.56
Al ₂ O ₃	13.51	15.67	10.43	24.72	8.30
Fe ₂ O ₃	3.74	5.22	2.58	.43	.38
FeO	1.02	.35	.48	-----	-----
MgO	1.09	1.60	1.13	.13	.24
CaO	2.29	2.77	4.64	.30	.12
Na ₂ O	1.70	.97	1.29	trace	trace
K ₂ O	3.14	2.28	2.50	trace	trace
H ₂ O at 100°	4.19	9.83	3.77	1.58	1.26
H ₂ O above 100°				8.41	4.40
TiO ₂	-----	-----	-----	.68	.60
P ₂ O ₅45	.19	.20	trace	.06
MnO	trace	trace	-----	-----	-----
CO ₂	trace	.31	2.59	-----	-----
Organic matter	-----	-----	-----	.40	8.31
	100.40	100.16	100.24	100.17	100.23

F. From Red Creek Canyon, south part Colorado Springs quadrangle.

G. From 2 miles southeast of F.

H. From near Canyon.

I. Overlying H.

Collected as probable fire clays by G. K. Gilbert. Analyses by George Steiger, record No. 1578. Fe_2O_3 represents total iron. Al_2O_3 includes TiO_2 . In I, the ignition includes some CO_2 , which is absent from the others.

	F.	G.	H.	I.
SiO_2	85.09	86.79	57.98	69.04
Al_2O_3	6.98	8.29	27.51	14.51
Fe_2O_3	1.10	.75	1.68	3.78
MgO27	.13	.32	.73
CaO21	.34	.42	1.24
Na_2O	none	none	.03	.08
K_2O13	.25	.56	.48
Ignition.....	6.37	3.78	11.80	10.50
P_2O_506	.05	.06	.07
	100.21	100.38	100.36	100.43

J to K. Supposed fire clays collected in the area of the Apishapa sheet, by G. K. Gilbert. Analyses, partial, by H. N. Stokes, record No. 1503. Titanium present, alkalies undetermined. Analyses made on ignited material, reckoned as 100. The loss on ignition is separately stated below each analysis.

	J.	K.	L.	M.	N.
SiO ₂	86.58	78.07	76.96	61.98	93.11
Al ₂ O ₃	12.72	20.22	20.77	37.51	5.56
Fe ₂ O ₃45	.89	1.11	.45	1.15
MgO11	.26	.32	.09	.10
CaO11	-----	.71	.19	.32
	98.97	99.44	99.87	100.22	100.24
Ignition	4.75	7.51	7.98	12.51	4.45

	O.	P.	Q.	R.
SiO ₂	85.98	85.25	54.93	58.56
Al ₂ O ₃	13.67	11.45	43.65	39.17
Fe ₂ O ₃41	2.24	.69	.55
MgO	-----	.21	.05	.45
CaO21	.26	.64	1.08
	100.27	99.41	99.96	99.81
Ignition	5.07	4.81	16.80	19.58

S. Loess-like alluvium, Golden, Jefferson County.

T. Fire clay, Golden, Jefferson County.

Analyses S and T made by W. F. Hillebrand in the Denver laboratory.

	S.	T.
SiO ₂	72.31	50.35
Al ₂ O ₃	12.66	34.44
Fe ₂ O ₃	4.67	.75
MgO94	trace
CaO	1.15	-----
Na ₂ O	2.47	.10
K ₂ O	3.75	.48
H ₂ O+organic matter	1.80	13.88
P ₂ O ₅23	-----
	99.98	100.00

WYOMING, UTAH, NEW MEXICO.

A. Loess, Cheyenne, Wyoming. Analysis by L. G. Eakins, record No. 1066.

B. Adobe soil, Salt Lake City, Utah.

C. Adobe soil, Santa Fe, New Mexico.

D. Adobe soil, Fort Wingate, New Mexico.

Analyses B, C, and D by Eakins, Nos. 981, 990.

	A.	B.	C.	D.
SiO ₂	67.10	19.24	66.69	26.67
Al ₂ O ₃	10.26	3.26	14.16	.91
Fe ₂ O ₃	2.52	1.09	4.38	.64
FeO.....	.31			
MgO.....	1.24	2.75	1.28	.51
CaO.....	5.88	38.94	2.49	36.40
Na ₂ O.....	1.42	trace	.67	trace
K ₂ O.....	2.68	trace	1.21	trace
H ₂ O.....	5.09	1.67	4.94	2.26
P ₂ O ₅11	.23	.29	.75
MnO.....		trace	.09	trace
CO ₂	3.67	29.57	.77	25.84
SO ₃53	.41	.82
Cl.....		.11	.34	.07
Organic matter.....		2.96	2.00	5.10
	100.28	100.35	99.72	99.97

NEVADA.

A. Grayish clay from Upper Lahontan lake beds, Humboldt River bridge, Mill City.

B. Grayish clay, Lower Lahontan beds, same locality. Analyses by T. M. Chatard, record Nos. 32, 33.

C. Adobe soil, Humboldt. Analysis by L. G. Eakins, record No. 981.

D. Halloysite, pale greenish, Lucia mining district, Elko County. Analysis by George Steiger, record No. 1472.

	A.	B.	C.	D.
SiO ₂	56.30	50.70	44.64	42.11
Al ₂ O ₃	16.52	19.01	13.19	33.83
Fe ₂ O ₃	5.08		5.12	.04
FeO.....				.28
MgO.....	2.64	3.19	2.96	.30
CaO.....	5.45	10.26	13.91	.33
Na ₂ O.....	2.60	1.91	.59	-----
K ₂ O.....	2.17	2.16	1.71	-----
H ₂ O at 100°.....	9.78	13.03	3.89	6.54
H ₂ O at 200°.....				1.07
H ₂ O at 300°.....				1.26
H ₂ O, ignition.....				12.04
P ₂ O ₅94	trace
MnO.....			.13	-----
CuO.....				2.83
CO ₂			8.55	-----
SO ₃64	-----
Cl.....			.14	-----
Organic matter.....			3.43	-----
	100.54	100.26	99.84	100.63

CALIFORNIA, WASHINGTON, HAWAIIAN ISLANDS.

A. Sandy clay, Owens Lake, California.

B. Blue clay, Owens Lake, California. Analyses by T. M. Chatard, record No. 551.

C. Clay from foot of Rickey Hill, Kittle Falls, Stevens County, Washington. Analysis by W. F. Hillebrand, record No. 1428.

D. Lava soil, Diamond Head, Hawaiian Islands. Analysis by L. G. Eakins, record No. 888.

	A.	B.	C.	D.
SiO ₂	53.24	54.92	62.74	32.88
Al ₂ O ₃	10.84	11.25	16.45	12.02
Fe ₂ O ₃	2.59	2.77	2.62	11.52
FeO.....	.77	.94	1.91
MgO.....	5.82	4.91	2.41	11.70
CaO.....	9.18	8.76	3.68	12.20
Na ₂ O.....	2.06	2.10	3.05	undet.
K ₂ O.....	2.64	2.77	3.53	undet.
H ₂ O at 110°.....	1.41	2.05	} 2.69	} 5.30
H ₂ O at redness.....	2.73	2.40		
TiO ₂25	.30
P ₂ O ₅24
MnO.....	.10	.08	trace	trace
SrO.....	trace
CO ₂	8.75	7.24	.65	11.41
SO ₃08	trace
Cl.....	.05	trace91
	100.51	100.49	99.73	98.18

INDEX TO ROCKS.

	Page.		Page.
Absarokite.....	97, 99, 103	Augite-microcline-granite	55
Acmite-trachyte.....	123	Augite-oligoclase-scapolite-biotite	
Actinolite-magnetite-schist.....	83	rock	229
Adinole.....	69	Augite-porphyrityte	167
Adobe soil.....	302, 303	Augite-porphyrity	112
Alaskite	228	Augite-syenite.....	95, 204
Albite	21, 23, 34	Augite-syenite-porphyrity	95
Aleutite	229	Augite-vogesite.....	130
Amphibole	116, 190, 206, 208	Banakite	101, 102
Amphibole-biotite-granite.....	208	Barkevikite	133
Amphibole-gabbro	206	Basalt.....	35, 39,
Amphibole-schist	217	61, 62, 63, 68, 92, 109, 110, 111, 113,	
Amphibolite	27, 31, 32, 66, 197	130, 132, 138, 140, 144, 145, 146, 169,	
Amygdaloid	53	170, 171, 173, 175, 176, 184, 185, 186,	
Analcite	34, 146	189, 191, 202, 218, 220, 222, 223, 225	
Analcite-basalt	146	Basanite	61, 171
Andalusite-hornfels.....	210	Belugite	229
Andalusite-schist	210	Biotite.....	116
Andesite.....	19,	Biotite-augite-latite.....	205
51, 90, 91, 97, 98, 108, 109, 114,		Biotite-diorite.....	44
119, 141, 145, 148, 153, 166, 170,		Biotite-gneiss	72
171, 174, 176, 181, 182, 183, 188,		Biotite-granite	30, 46, 47, 208
191, 193, 203, 212, 222, 226, 227		Biotite-porphyrityte	155
Andesite-basalt.....	176	Biotite-quartz-monzonite.....	192
Andesite-perlite	175	Biotite-trachyte	98
Andesite-porphyrity	119	Boltonite	33
Anorthoclase.....	77, 123	Bostonite	19
Anorthosite.....	36, 37	Bronzite.....	43
Aplite	94, 129, 192, 207, 229	Bronzite-norite.....	67
Aporhyolite.....	40	Camptonite	23, 26, 110, 162, 163, 197
Argillite	195, 196	Cancrinite	22
Arkose-gneiss	33	Cement rock	253, 257, 259
Asperite	212	Chert.....	250
Augite.....	63, 140, 146, 170	Chiasmolite-schist	210
Augite-aleutite	229	Chloritic gneiss	48
Augite-andesite	98, 141	Chloritic granite	27
Augite-andesite-porphyrity	89, 93	Clay	289, 290, 291, 292, 293, 294,
Augite-belugite.....	229	295, 296, 297, 298, 300, 301, 303, 304	
Augite-bronzite-andesite	226	Clay slate.....	284, 288
Augite-diorite	147, 161, 164	Coquina	255
Augite-granite.....	198	Coral	256
Augite-latite	205	Coral rock	255, 256, 277
Augite-mica-syenite.....	140	Cordierite-hornfels	26

	Page.		Page.
Cortlandtite	43	Granite-porphyrý ..	25, 120, 125, 158, 207
Dacite	86, 91,	Granite-syenite-porphyrý	126, 134
106, 107, 149, 174, 177, 179, 180, 203		Granitite	24, 120, 142, 141, 199
Dacite-porphyrý	106, 177	Granitoid gneiss	27, 142
Diabase	20, 26,	Granodiorite	139, 187,
30, 37, 50, 52, 73, 76, 96, 121, 141,		190, 194, 195, 196, 197, 198, 199, 224	
193, 194, 209, 211, 213, 217, 225		Granodiorite-porphyrý	224
Diabase-diorite	213	Granulite	192, 207
Diabase-porphyrý	121, 217	Graywacke	75
Diabase-porphyrý	193	Greenstone	64, 71, 72
Diabasic basalt	138	Grünerite-magnetite-schist	64
Diallage	75, 76	Halloysite	303
Diallage-gabbro	81	Hornblende	133
Diopside	43, 86	Hornblende-andesite	90,
Diorite	25,	108, 176, 181, 188, 227	
26, 45, 52, 55, 67, 71, 83, 87, 88, 94,		Hornblende-augite-andesite	97
96, 118, 121, 122, 126, 130, 137, 147,		Hornblende-basalt	186
158, 159, 160, 161, 164, 177, 190,		Hornblende-diorite	44
191, 192, 200, 201, 204, 209, 226, 227		Hornblende-gabbro	67
Diorite-monzonite	163	Hornblende-granite	27
Diorite-porphyrý	121	Hornblende-granitite	120
Diorite-porphyrý	72,	Hornblende-mica-andesite	91,
126, 158, 161, 192, 200		94, 107, 108	
Dolerite	68, 140, 189	Hornblende-mica-porphyrý	160
Dolomite	252,	Hornblende-mica-porphyrý	89
253, 258, 263, 264, 270, 272, 273, 274		Hornblende-mica-syenite	130
Dunite	54	Hornblende-picrite	114, 124
Elæolite	21, 23	Hornblende-porphyrý	155,
Elæolite-syenite	21, 23, 39, 123, 144	164, 167, 173, 194	
Enstatite	28	Hornblende-porphyrý	89
Enstatite-díabase-porphyrý	141	Hornblende-pyroxene-andesite	90,
Epidote-chlorite-schist	53	97, 188, 193	
Feldspar	21, 23,	Hornfels	26, 210
34, 41, 49, 73, 74, 75, 77, 82, 170, 220		Hornstone	122
Feldspar-porphyrý	130	Hydraulic cement rock	253
Felsite	51	Hydronephelite	21
Felsophyre	51	Hypersthene	81, 153
Ferrodolomite	267	Hypersthene-andesite	114,
Fulgurite	220	153, 174, 176, 182, 183, 188, 193	
Gabbro	36, 37,	Hypersthene-augite-andesite	222
41, 44, 55, 59, 67, 70, 81, 82, 83, 92, 93,		Hypersthene-basalt	220
95, 97, 122, 137, 163, 198, 206, 214, 225		Hypersthene-gabbro	44, 81
Gabbro-diorite	44, 71, 83, 93	Ilmenite	56
Gabbro-porphyrý	97, 163	Iron carbonate	265, 266
Garnet	56	Jasperite	250
Genthite	221	Kaolin	290, 295
Glaucophane-schist	211, 213	Keratophyre	33, 77
Gneiss	27, 33,	Kersantite	110
47, 48, 54, 65, 66, 72, 83, 200, 201, 202		Kimberlite	56
Granite	25, 27,	Kyanitic schist	230
30, 46, 47, 55, 56, 59, 66, 77, 80, 94, 96,		Lamprophyre	112, 113, 162, 163, 165
116, 117, 118, 120, 125, 129, 134, 141,		Lapilli	184
142, 158, 164, 187, 192, 198, 199, 207		Latite	205

	Page.		Page.
Lava soil.....	304	Novaculite	65
Lepidomelane.....	21	Obsidian..... 104, 119, 172, 212, 217, 219	
Leucite-absarokite.....	97, 99	Olivine	56, 81, 221
Leucite-banakite	102	Olivine-basalt	35, 189, 218
Leucite-basalt	132	Olivine-diabase.....	76, 225
Leucite-monchiquite.....	132	Olivine-gabbro	44, 81, 122, 206
Leucite-shoshonite	100	Oolitic sand.....	275
Leucite-syenite	133	Orendite	86
Leucitite.....	136	Orthoclase	23, 155, 156
Lherzolite	42	Orthoclase-gabbro-diorite	93
Limburgite	38	Ottrelite	50
Limestone.....	17, 252,	Ottrelite-phyllite	50
253, 254, 255, 257, 258, 259, 260,		Ouachitite	60
261, 262, 263, 264, 268, 269, 270,		Pantellarite	60
271, 272, 273, 274, 275, 276, 277		Pegmatite.....	207
Litchfieldite	21	Peridotite.....	30,
Lithoidite.....	104	38, 54, 56, 58, 64, 114, 147, 191, 221	
Loess..... 295, 296, 298, 299, 301, 302		Perlite.....	108, 175
Madupite	85	Phlogopite	85
Magnetite.....	75, 83	Phonolite	33, 62, 84, 143, 144, 171
Marble	252, 255, 258, 275	Phonolitic andesite	170
Marl	263	Phthanite	288
Melaphyr.....	202	Phyllite	26
Metaandesite.....	191, 203	Pierite	67, 114, 124
Metabasalt	68	Pierite-porphry	67
Metadacite	203	Pitchstone	35, 119, 129, 150, 168
Metadolerite	68	Plagioclase	49
Meta-quartz-diorite	55	Plagioclase-basalt.....	61, 145, 171
Metarhyolite.....	187	Plagioclase-gneiss.....	201
Meteorites	231-244	Porphyrite	112, 120, 121, 155, 157,
Mica..... 21, 50, 85, 202, 208, 218		160, 164, 167, 173, 179, 199, 200, 217	
Mica-andesite	170	Porphyrite-diorite	160
Mica-basalt	173	Porphyry	23, 24, 25, 40, 52, 59,
Mica-dacite.....	107, 149	67, 72, 77, 89, 92, 93, 94, 95, 96, 97,	
Mica-diorite.....	67, 72	106, 107, 112, 119, 124, 125, 126, 129,	
Mica-gabbro.....	92	130, 134, 135, 141, 154, 155, 158, 166,	
Mica-peridotite	58	167, 177, 192, 200, 204, 207, 226, 228	
Mica-schist.....	66, 83, 210	Pseudo-diabase	211
Microcline	49	Pseudo-diorite.....	211
Micropegmatite	207	Pseudo-leucite-sodalite-tinguaite ..	136
Microperthite-hornfels	26	Pseudo-leucite-syenite	133
Minette.....	39, 128	Pumice	219
Missourite	133	Pyroxene	111, 165, 220
Monchiquite	60, 122, 128, 130	Pyroxene-andesite	108,
Monzonite	93, 118, 128, 133,	109, 145, 171, 181	
135, 137, 139, 162, 163, 166, 192, 200		Pyroxene-mica-diorite	87
Nepheline	21, 23	Pyroxene-porphry.....	87
Nepheline-basalt	62, 63, 144	Pyroxenite.....	42, 43, 53, 213
Nepheline-basanite	171	Quartz-alunite rock	151
Nepheline-melilite-basalt	63	Quartz-andesite	193
Nepheline-syenite 21, 33, 39, 123, 144		Quartz-banakite	102
Nevadite.....	156	Quartz-basalt.....	184, 185
Norite	36, 37, 45, 67, 83	Quartz-biotite-hornblende-diorite ..	45

	Page.		Page.
Quartz-diaspore rock.....	169	Siliceous sinter	251
Quartz-diorite..... 94, 121, 190, 192, 200		Slate... 79, 278, 279, 280, 281, 282, 284, 285	
Quartz-diorite-gneiss.....	200	Sodalite.....	22, 134
Quartz-diorite-porphry	94, 192	Sodalite-syenite	133
Quartz-gabbro.....	55	Sölvbergite.....	123
Quartz - hornblende - mica-porphyr- rite.....	157	Spherulites	152
Quartzite	78, 79, 80	Spilosite	69
Quartz-keratophyre	77	Syenite.....	21, 24,
Quartz-mica-diorite... 88, 89, 94, 192, 209		25, 39, 95, 121, 123, 125, 126, 130,	
Quartz-mica-diorite-porphry	94	132, 133, 134, 140, 144, 151, 161, 204	
Quartz-monzonite.....	118,	Syenite-diorite-porphry.....	126
137, 139, 163, 192, 200		Syenite-porphry	24, 95, 125, 204
Quartz-norite-gneiss	83	Tachylite	35
Quartz-pantellarite	60	Talcose schist	72
Quartz-porphryrite..... 157, 160, 194, 200		Tallow clay	296, 298
Quartz-porphryrite-schist	200	Teschenite.....	19
Quartz-porphry	23,	Theralite	123, 124
40, 52, 72, 119, 124, 129, 166, 226		Tinguaite.....	123, 136, 165
Quartz-pyroxene-diorite	204	Tonalite	30, 229
Quartz-pyroxene-mica-diorite	87	Tonalite-aplite	229
Quartz-schist.....	50	Tordrillite	229
Quartz-syenite	95, 135	Tourmaline-biotite-schist	84
Quartz-syenite-porphry.....	126, 135	Trachy-andesitic breccia	131
Quartz-tourmaline-porphry	129	Trachyte.....	19,
Quartz-trachyte	19	98, 106, 123, 131, 145, 147, 172	
Rhyolite	20, 52,	Trachyte-andesite.....	98
53, 104, 105, 106, 108, 119, 129, 130,		Trachytic phonolite.....	144
138, 149, 150, 152, 156, 159, 161, 164,		Trachytic rhyolite.....	105, 106
166, 168, 174, 178, 187, 188, 219, 221		Travertine	268
Sandstone.....	17,	Uralite-diabase	195
19, 57, 244, 245, 246, 247, 248, 249		Uralite-diorite.....	191
Saussurite-gabbro.....	70	Vitrophyre	161, 221
Saxonite	221	Vogesite	130
Schist..... 50, 53, 64, 65, 66, 70, 71, 72,		Volcanic ash	219
83, 84, 200, 210, 211, 213, 217, 230		Volcanic glass	115, 152, 168
Sericite-schist	50, 65	Volcanic sand	115, 184
Serpentine	28,	Volcanic tuff	20
29, 33, 189, 195, 212, 215, 216, 224		Websterite.....	43, 53, 213
Shale..... 17, 57, 283, 284, 286, 287, 288		Wehrlite.....	33, 67, 114
Shell rock	256, 277	Wollastonite-gneiss	202
Shonkinite.....	128, 136	Wyomingite	85
Shoshonite.....	100, 103	Yogoite.....	135

ADVERTISEMENT.

[Bulletin 168.]

The statute approved March 3, 1879, establishing the United States Geological Survey, contains the following provisions:

"The publications of the Geological Survey shall consist of the annual report of operations, geological and economic maps illustrating the resources and classification of the lands, and reports upon general and economic geology and paleontology. The annual report of operations of the Geological Survey shall accompany the annual report of the Secretary of the Interior. All special memoirs and reports of said Survey shall be issued in uniform quarto series if deemed necessary by the Director, but otherwise in ordinary octavos. Three thousand copies of each shall be published for scientific exchanges and for sale at the price of publication; and all literary and cartographic materials received in exchange shall be the property of the United States and form a part of the library of the organization; and the money resulting from the sale of such publications shall be covered into the Treasury of the United States."

Except in those cases in which an extra number of any special memoir or report has been supplied to the Survey by resolution of Congress or has been ordered by the Secretary of the Interior, this office has no copies for gratuitous distribution.

ANNUAL REPORTS.

- I. First Annual Report of the United States Geological Survey, by Clarence King. 1880. 8°. 79 pp. 1 map.—A preliminary report describing plan of organization and publications.
- II. Second Annual Report of the United States Geological Survey, 1880-'81, by J. W. Powell. 1882. 8°. iv, 588 pp. 62 pl. 1 map.
- III. Third Annual Report of the United States Geological Survey, 1881-'82, by J. W. Powell. 1883. 8°. xviii, 564 pp. 67 pl. and maps.
- IV. Fourth Annual Report of the United States Geological Survey, 1882-'83, by J. W. Powell. 1884. 8°. xxxii, 473 pp. 85 pl. and maps.
- V. Fifth Annual Report of the United States Geological Survey, 1883-'84, by J. W. Powell. 1885. 8°. xxxvi, 469 pp. 58 pl. and maps.
- VI. Sixth Annual Report of the United States Geological Survey, 1884-'85, by J. W. Powell. 1885. 8°. xxix, 570 pp. 65 pl. and maps.
- VII. Seventh Annual Report of the United States Geological Survey, 1885-'86, by J. W. Powell. 1888. 8°. xx, 656 pp. 71 pl. and maps.
- VIII. Eighth Annual Report of the United States Geological Survey, 1886-'87, by J. W. Powell. 1889. 8°. 2 pt. xix, 474, xii pp. 53 pl. and maps; 1 p.l., 475-1063 pp. 54-76 pl. and maps.
- IX. Ninth Annual Report of the United States Geological Survey, 1887-'88, by J. W. Powell. 1889. 8°. xiii, 717 pp. 88 pl. and maps.
- X. Tenth Annual Report of the United States Geological Survey, 1888-'89, by J. W. Powell. 1890. 8°. 2 pt. xv, 774 pp., 98 pl. and maps; viii, 123 pp.
- XI. Eleventh Annual Report of the United States Geological Survey, 1889-'90, by J. W. Powell. 1891. 8°. 2 pt. xv, 757 pp., 66 pl. and maps; ix, 351 pp., 30 pl.
- XII. Twelfth Annual Report of the United States Geological Survey, 1890-'91, by J. W. Powell. 1891. 8°. 2 pt. xiii, 675 pp., 53 pl. and maps; xviii, 576 pp., 146 pl. and maps.
- XIII. Thirteenth Annual Report of the United States Geological Survey, 1891-'92, by J. W. Powell. 1893. 8°. 3 pt. vii, 240 pp., 2 maps; x, 372 pp., 105 pl. and maps; xi, 486 pp., 77 pl. and maps.
- XIV. Fourteenth Annual Report of the United States Geological Survey, 1892-'93, by J. W. Powell. 1893. 8°. 2 pt. vi, 321 pp., 1 pl.; xx, 597 pp., 74 pl.
- XV. Fifteenth Annual Report of the United States Geological Survey, 1893-'94, by J. W. Powell. 1895. 8°. xiv, 755 pp. 48 pl.
- XVI. Sixteenth Annual Report of the United States Geological Survey, 1894-'95, by Charles D. Walcott, Director. 1895. (Part I, 1896.) 8°. 4 pt. xxii, 910 pp., 117 pl. and maps; xix, 598 pp., 43 pl. and maps; xv, 646 pp., 23 pl.; xix, 735 pp., 6 pl.

XVII. Seventeenth Annual Report of the United States Geological Survey, 1895-'96, Charles D. Walcott, Director. 1896. 8°. 3 pt. in 4 vol. xxii, 1076 pp., 67 pl. and maps; xxv, 864 pp., 113 pl. and maps; xxiii, 542 pp., 8 pl. and maps; iii, 543-1058 pp., 9-13 pl.

XVIII. Eighteenth Annual Report of the United States Geological Survey, 1896-'97, Charles D. Walcott, Director. 1897. (Parts II and III, 1898.) 8°. 5 pt. in 6 vol. 440 pp., 4 pl. and maps; v, 653 pp., 105 pl. and maps; v, 861 pp., 118 pl. and maps; x, 756 pp., 102 pl. and maps; xii, 642 pp., 1 pl.; 643-1400 pp.

XIX. Nineteenth Annual Report of the United States Geological Survey, 1897-'98, Charles D. Walcott, Director. 1898. (Parts II, III, and V, 1899.) 8°. 6 pt. in 7 vol. 422 pp., 2 maps; v, 958 pp., 172 pl. and maps; v, 785 pp., 99 pl. and maps; viii, 814 pp., 118 pl. and maps; xvii, 400 pp., 110 pl. and maps; viii, 651 pp., 11 pl.; viii, 706 pp.

XX. Twentieth Annual Report of the United States Geological Survey, 1898-'99, Charles D. Walcott, Director. 1899. 8°. 7 pt. in 8 vol. 551 pp., 2 pl.; v, 953 pp., 193 pl. and maps; v, 595 pp., 78 pl. and maps; vii, 660 pp., 75 pl. and maps; xix, 498 pp., 159 pl. and maps; viii, 616 pp.; xi, 804 pp., 1 pl.; v, 509 pp., 38 pl. and maps.

MONOGRAPHS.

I. Lake Bonneville, by Grove Karl Gilbert. 1890. 4°. xx, 438 pp. 51 pl. 1 map. Price \$1.50.

II. Tertiary History of the Grand Cañon District, with Atlas, by Clarence E. Dutton, Capt., U. S. A. 1882. 4°. xiv, 264 pp. 42 pl. and atlas of 24 sheets folio. Price \$10.00.

III. Geology of the Comstock Lode and the Washoe District, with Atlas, by George F. Becker. 1882. 4°. xv, 422 pp. 7 pl. and atlas of 21 sheets folio. Price \$11.00.

IV. Comstock Mining and Miners, by Eliot Lord. 1883. 4°. xiv, 451 pp. 3 pl. Price \$1.50.

V. The Copper-Bearing Rocks of Lake Superior, by Roland Duer Irving. 1883. 4°. xvi, 464 pp. 15 l. 29 pl. and maps. Price \$1.85.

VI. Contributions to the Knowledge of the Older Mesozoic Flora of Virginia, by William Morris Fontaine. 1883. 4°. xi, 144 pp. 54 l. 54 pl. Price \$1.05.

VII. Silver-Lead Deposits of Eureka, Nevada, by Joseph Story Curtis. 1884. 4°. xiii, 200 pp. 16 pl. Price \$1.20.

VIII. Paleontology of the Eureka District, by Charles Doolittle Walcott. 1884. 4°. xiii, 298 pp. 24 l. 24 pl. Price \$1.10.

IX. Brachiopoda and Lamellibranchiata of the Raritan Clays and Greensand Marls of New Jersey, by Robert P. Whitfield. 1885. 4°. xx, 338 pp. 35 pl. 1 map. Price \$1.15.

X. Dinocerata. A Monograph of an Extinct Order of Gigantic Mammals, by Othniel Charles Marsh. 1886. 4°. xviii, 243 pp. 56 l. 56 pl. Price \$2.70.

XI. Geological History of Lake Lahontan, a Quaternary Lake of Northwestern Nevada, by Israel Cook Russell. 1885. 4°. xiv, 288 pp. 46 pl. and maps. Price \$1.75.

XII. Geology and Mining Industry of Leadville, Colorado, with Atlas, by Samuel Franklin Emmons. 1886. 4°. xxix, 770 pp. 45 pl. and atlas of 35 sheets folio. Price \$8.40.

XIII. Geology of the Quicksilver Deposits of the Pacific Slope, with Atlas, by George F. Becker. 1888. 4°. xix, 486 pp. 7 pl. and atlas of 14 sheets folio. Price \$2.00.

XIV. Fossil Fishes and Fossil Plants of the Triassic Rocks of New Jersey and the Connecticut Valley, by John S. Newberry. 1888. 4°. xiv, 152 pp. 26 pl. Price \$1.00.

XV. The Potomac or Younger Mesozoic Flora, by William Morris Fontaine. 1889. 4°. xiv, 377 pp. 180 pl. Text and plates bound separately. Price \$2.50.

XVI. The Paleozoic Fishes of North America, by John Strong Newberry. 1889. 4°. 340 pp. 53 pl. Price \$1.00.

XVII. The Flora of the Dakota Group, a Posthumous Work, by Leo Lesquereux. Edited by F. H. Knowlton. 1891. 4°. 400 pp. 66 pl. Price \$1.10.

XVIII. Gasteropoda and Cephalopoda of the Raritan Clays and Greensand Marls of New Jersey, by Robert P. Whitfield. 1891. 4°. 402 pp. 50 pl. Price \$1.00.

XIX. The Penokee Iron-Bearing Series of Northern Wisconsin and Michigan, by Roland D. Irving and C. R. Van Hise. 1892. 4°. xix, 534 pp. 37 pl. Price \$1.70.

XX. Geology of the Eureka District, Nevada, with Atlas, by Arnold Hague. 1892. 4°. xvii, 419 pp. 8 pl. Price \$5.25.

XXI. The Tertiary Rhynchophorous Coleoptera of North America, by Samuel Hubbard Scudder. 1893. 4°. xi, 206 pp. 18 pl. Price 90 cents.

XXII. A Manual of Topographic Methods, by Henry Gannett, Chief Topographer. 1893. 4°. xiv, 300 pp. 18 pl. Price \$1.00.

XXIII. Geology of the Green Mountains in Massachusetts, by Raphael Pumpelly, J. E. Wolff, and T. Nelson Dale. 1894. 4°. xiv, 206 pp. 23 pl. Price \$1.30.

XXIV. Mollusca and Crustacea of the Miocene Formations of New Jersey, by Robert Parr Whitfield. 1894. 4°. 195 pp. 24 pl. Price 90 cents.

XXV. The Glacial Lake Agassiz, by Warren Upham. 1895. 4°. xxiv, 658 pp. 38 pl. Price \$1.70.

XXVI. Flora of the Amboy Clays, by John Strong Newberry; a Posthumous Work, edited by Arthur Hollick. 1895. 4°. 260 pp. 58 pl. Price \$1.00.

XXVII. Geology of the Denver Basin, Colorado, by S. F. Emmons, Whitman Cross, and George H. Eldridge. 1896. 4°. 556 pp. 31 pl. Price \$1.50.

- XXVIII. The Marquette Iron-Bearing District of Michigan, with Atlas, by C. R. Van Hise and W. S. Bayley, including a Chapter on the Republic Trough, by H. L. Smyth. 1897. 4°. 608 pp. 35 pl. and atlas of 39 sheets folio. Price \$5.75.
- XXIX. Geology of Old Hampshire County, Massachusetts, comprising Franklin, Hampshire, and Hampden Counties, by Benjamin Kendall Emerson. 1898. 4°. xxi, 790 pp. 35 pl. Price \$1.90.
- XXX. Fossil Medusæ, by Charles Doolittle Walcott. 1898. 4°. ix, 201 pp. 47 pl. Price \$1.50.
- XXXI. Geology of the Aspen Mining District, Colorado, with Atlas, by Josiah Edward Spurr. 1898. 4°. xxxv, 260 pp. 43 pl. and atlas of 30 sheets folio. Price \$3.60.
- XXXII. Geology of the Yellowstone National Park, Part II, Descriptive Geology, Petrography, and Paleontology, by Arnold Hague, J. P. Iddings, W. Harvey Weed, Charles D. Walcott, G. H. Girty, T. W. Stanton, and F. H. Knowlton. 1899. 4°. xvii, 693 pp. 121 pl. Price \$2.45.
- XXXIII. Geology of the Narragansett Basin, by N. S. Shaler, J. B. Woodworth, and August F. Foerste. 1899. 4°. xx, 402 pp. 31 pl. Price \$1.
- XXXIV. The Glacial Gravels of Maine and their Associated Deposits, by George H. Stone. 1899. 4°. xiii, 499 pp. 52 pl. Price \$1.30.
- XXXV. The Later Extinct Floras of North America, by John Strong Newberry; edited by Arthur Hollick. 1898. 4°. xviii, 295 pp. 68 pl. Price \$1.25.
- XXXVI. The Crystal Falls Iron-Bearing District of Michigan, by J. Morgan Clements and Henry Lloyd Smyth; with a Chapter on the Sturgeon River Tongue, by William Shirley Bayley, and an Introduction by Charles Richard Van Hise. 1899. 4°. xxxvi, 512 pp. 53 pl. Price \$2.
- XXXVII. Fossil Flora of the Lower Coal Measures of Missouri, by David White. 1899. 4°. xi, 467 pp. 73 pl. Price \$1.25.
- XXXVIII. The Illinois Glacial Lobe, by Frank Leverett. 1899. 4°. xxi, 817 pp. 24 pl. Price \$1.60.
- In preparation:*
- XXXIX. The Eocene and Lower Oligocene Coral Faunas of the United States, with Descriptions of a Few Doubtfully Cretaceous Species, by Thomas Wayland Vaughan.
- Adephagous and Clavicorn Coleoptera from the Tertiary Deposits at Florissant, Colorado, with Descriptions of a Few Other Forms and including a Systematic List of the Non-Rhynchophorous Tertiary Coleoptera of North America, by Samuel Hubbard Scudder.
- Flora of the Laramie and Allied Formations, by Frank Hall Knowlton.

BULLETINS.

1. On Hypersthene-Andesite and on Triclinic Pyroxene in Augitic Rocks, by Whitman Cross, with a Geological Sketch of Buffalo Peaks, Colorado, by S. F. Emmons. 1883. 8°. 42 pp. 2 pl. Price 10 cents.
2. Gold and Silver Conversion Tables, giving the Coining Value of Troy Onnces of Fine Metal, etc., computed by Albert Williams, jr. 1883. 8°. 8 pp. Price 5 cents.
3. On the Fossil Faunas of the Upper Devonian, along the Meridian of 76° 30', from Tompkins County, New York, to Bradford County, Pennsylvania, by Henry S. Williams. 1884. 8°. 36 pp. Price 5 cents.
4. On Mesozoic Fossils, by Charles A. White. 1884. 8°. 36 pp. 9 pl. Price 5 cents.
5. A Dictionary of Altitudes in the United States, compiled by Henry Gannett. 1884. 8°. 325 pp. Price 20 cents.
6. Elevations in the Dominion of Canada, by J. W. Spencer. 1884. 8°. 43 pp. Price 5 cents.
7. *Mapoteca Geologica Americana*. A Catalogue of Geological Maps of America (North and South), 1752-1881, in Geographic and Chronologic Order, by Jules Marcou and John Belknap Marcou. 1884. 8°. 184 pp. Price 10 cents.
8. On Secondary Enlargements of Mineral Fragments in Certain Rocks, by R. D. Irving and C. R. Van Hise. 1884. 8°. 56 pp. 6 pl. Price 10 cents.
9. A Report of Work done in the Washington Laboratory during the Fiscal Year 1883-'84. F. W. Clarke, Chief Chemist. T. M. Chatard, Assistant Chemist. 1884. 8°. 40 pp. Price 5 cents.
10. On the Cambrian Faunas of North America. Preliminary Studies, by Charles Doolittle Walcott. 1884. 8°. 74 pp. 10 pl. Price 5 cents.
11. On the Quaternary and Recent Mollusca of the Great Basin; with Descriptions of New Forms, by R. Ellsworth Call. Introduced by a Sketch of the Quaternary Lakes of the Great Basin, by G. K. Gilbert. 1884. 8°. 66 pp. 6 pl. Price 5 cents.
12. A Crystallographic Study of the Thinolite of Lake Lahontan, by Edward S. Dana. 1884. 8°. 24 pp. 3 pl. Price 5 cents.
13. Boundaries of the United States and of the Several States and Territories, with a Historical Sketch of the Territorial Changes, by Henry Gannett. 1885. 8°. 135 pp. Price 10 cents. (Exhausted.)
14. The Electrical and Magnetic Properties of the Iron-Carburets, by Carl Barns and Vincent Strouhal. 1885. 8°. 238 pp. Price 15 cents.
15. On the Mesozoic and Cenozoic Paleontology of California, by Charles A. White. 1885. 8°. 33 pp. Price 5 cents.
16. On the Higher Devonian Faunas of Ontario County, New York, by John M. Clarke. 1885. 8°. 86 pp. 3 pl. Price 5 cents.

17. On the Development of Crystallization in the Igneous Rocks of Washoe, Nevada, with Notes on the Geology of the District, by Arnold Hague and Joseph P. Iddings. 1885. 8°. 44 pp. Price 5 cents.
18. On Marine Eocene, Fresh-water Miocene, and Other Fossil Mollusca of Western North America, by Charles A. White. 1885. 8°. 26 pp. 3 pl. Price 5 cents.
19. Notes on the Stratigraphy of California, by George F. Becker. 1885. 8°. 28 pp. Price 5 cents. (Exhausted.)
20. Contributions to the Mineralogy of the Rocky Mountains, by Whitman Cross and W. F. Hillebrand. 1885. 8°. 114 pp. 1 pl. Price 10 cents.
21. The Lignites of the Great Sioux Reservation. A Report on the Region between the Grand and Moreau Rivers, Dakota, by Bailey Willis. 1885. 8°. 16 pp. 5 pl. Price 5 cents.
22. On New Cretaceous Fossils from California, by Charles A. White. 1885. 8°. 25 pp. 5 pl. Price 5 cents.
23. Observations on the Junction between the Eastern Sandstone and the Keweenaw Series on Keweenaw Point, Lake Superior, by R. D. Irving and T. C. Chamberlin. 1885. 8°. 124 pp. 17 pl. Price 15 cents.
24. List of Marine Mollusca, comprising the Quaternary Fossils and Recent Forms from American Localities between Cape Hatteras and Cape Roque, including the Bermudas, by William Healy Dall. 1885. 8°. 336 pp. Price 25 cents.
25. The Present Technical Condition of the Steel Industry of the United States, by Phineas Barnes. 1885. 8°. 85 pp. Price 10 cents.
26. Copper Smelting, by Henry M. Howe. 1885. 8°. 107 pp. Price 10 cents.
27. Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1884-'85. 1886. 8°. 80 pp. Price 10 cents.
28. The Gabbros and Associated Hornblende Rocks occurring in the Neighborhood of Baltimore, Maryland, by George Huntington Williams. 1886. 8°. 78 pp. 4 pl. Price 10 cents.
29. On the Fresh-water Invertebrates of the North American Jurassic, by Charles A. White. 1886. 8°. 41 pp. 4 pl. Price 5 cents.
30. Second Contribution to the Studies on the Cambrian Faunas of North America, by Charles Doolittle Walcott. 1886. 8°. 369 pp. 33 pl. Price 25 cents.
31. Systematic Review of our Present Knowledge of Fossil Insects, including Myriapods and Arachnids, by Samuel Hubbard Scudder. 1886. 8°. 128 pp. Price 15 cents.
32. Lists and Analyses of the Mineral Springs of the United States; (a Preliminary Study), by Albert C. Peale. 1886. 8°. 235 pp. Price 20 cents.
33. Notes on the Geology of Northern California, by J. S. Diller. 1886. 8°. 23 pp. Price 5 cents.
34. On the Relation of the Laramie Molluscan Fauna to that of the Succeeding Fresh-water Eocene and Other Groups, by Charles A. White. 1886. 8°. 54 pp. 5 pl. Price 10 cents.
35. Physical Properties of the Iron-Carburets, by Carl Barns and Vincent Strouhal. 1886. 8°. 62 pp. Price 10 cents.
36. Subsidence of Fine Solid Particles in Liquids, by Carl Barns. 1886. 8°. 58 pp. Price 10 cents.
37. Types of the Laramie Flora, by Lester F. Ward. 1887. 8°. 354 pp. 57 pl. Price 25 cents.
38. Peridotite of Elliott County, Kentucky, by J. S. Diller. 1887. 8°. 31 pp. 1 pl. Price 5 cents.
39. The Upper Beaches and Deltas of the Glacial Lake Agassiz, by Warren Upham. 1887. 8°. 84 pp. 1 pl. Price 10 cents.
40. Changes in River Courses in Washington Territory due to Glaciation, by Bailey Willis. 1887. 8°. 10 pp. 4 pl. Price 5 cents.
41. On the Fossil Faunas of the Upper Devonian—the Genesee Section, New York, by Henry S. Williams. 1887. 8°. 121 pp. 4 pl. Price 15 cents.
42. Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1885-'86. F. W. Clarke, Chief Chemist. 1887. 8°. 152 pp. 1 pl. Price 15 cents.
43. Tertiary and Cretaceous Strata of the Tuscaloosa, Tombigbee, and Alabama Rivers, by Eugene A. Smith and Lawrence C. Johnson. 1887. 8°. 189 pp. 21 pl. Price 15 cents.
44. Bibliography of North American Geology for 1886, by Nelson H. Darton. 1887. 8°. 35 pp. Price 5 cents.
45. The Present Condition of Knowledge of the Geology of Texas, by Robert T. Hill. 1887. 8°. 94 pp. Price 10 cents.
46. Nature and Origin of Deposits of Phosphate of Lime, by R. A. F. Penrose, jr., with an Introduction by N. S. Shaler. 1888. 8°. 143 pp. Price 15 cents.
47. Analyses of Waters of the Yellowstone National Park, with an Account of the Methods of Analysis employed, by Frank Austin Gooch and James Edward Whitfield. 1888. 8°. 84 pp. Price 10 cents.
48. On the Form and Position of the Sea Level, by Robert Simpson Woodward. 1888. 8°. 88 pp. Price 10 cents.
49. Latitudes and Longitudes of Certain Points in Missouri, Kansas, and New Mexico, by Robert Simpson Woodward. 1889. 8°. 133 pp. Price 15 cents.
50. Formulas and Tables to facilitate the Construction and Use of Maps, by Robert Simpson Woodward. 1889. 8°. 124 pp. Price 15 cents.

51. On Invertebrate Fossils from the Pacific Coast, by Charles Abiathar White. 1889. 8°. 102 pp. 14 pl. Price 15 cents.
52. Subaërial Decay of Rocks and Origin of the Red Color of Certain Formations, by Israel Cook Russell. 1889. 8°. 65 pp. 5 pl. Price 10 cents.
53. The Geology of Nantucket, by Nathaniel Southgate Shaler. 1889. 8°. 55 pp. 10 pl. Price 10 cents.
54. On the Thermo-Electric Measurement of High Temperatures, by Carl Barus. 1889. 8°. 313 pp. incl. 1 pl. 11 pl. Price 25 cents.
55. Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1886-'87. Frank Wigglesworth Clarke, Chief Chemist. 1889. 8°. 96 pp. Price 10 cents.
56. Fossil Wood and Lignite of the Potomac Formation, by Frank Hall Knowlton. 1889. 8°. 72 pp. 7 pl. Price 10 cents.
57. A Geological Reconnaissance in Southwestern Kansas, by Robert Hay. 1890. 8°. 49 pp. 2 pl. Price 5 cents.
58. The Glacial Boundary in Western Pennsylvania, Ohio, Kentucky, Indiana, and Illinois, by George Frederick Wright, with an Introduction by Thomas Chrowder Chamberlin. 1890. 8°. 112 pp. 8 pl. Price 15 cents.
59. The Gabbros and Associated Rocks in Delaware, by Frederick D. Chester. 1890. 8°. 45 pp. 1 pl. Price 10 cents.
60. Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1887-'88. F. W. Clarke, Chief Chemist. 1890. 8°. 174 pp. Price 15 cents.
61. Contributions to the Mineralogy of the Pacific Coast, by William Harlow Melville and Waldemar Lindgren. 1890. 8°. 40 pp. 3 pl. Price 5 cents.
62. The Greenstone Schist Areas of the Menominee and Marquette Regions of Michigan; a Contribution to the Subject of Dynamic Metamorphism in Eruptive Rocks, by George Huntington Williams; with an Introduction by Roland Duer Irving. 1890. 8°. 241 pp. 16 pl. Price 30 cents.
63. A Bibliography of Paleozoic Crustacea from 1698 to 1889, including a List of North American Species and a Systematic Arrangement of Genera, by Anthony W. Vogdes. 1890. 8°. 177 pp. Price 15 cents.
64. A Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1888-'89. F. W. Clarke, Chief Chemist. 1890. 8°. 60 pp. Price 10 cents.
65. Stratigraphy of the Bituminous Coal Field of Pennsylvania, Ohio, and West Virginia, by Israel C. White. 1891. 8°. 212 pp. 11 pl. Price 20 cents.
66. On a Group of Volcanic Rocks from the Tewan Mountains, New Mexico, and on the Occurrence of Primary Quartz in Certain Basalts, by Joseph Paxson Iddings. 1890. 8°. 34 pp. Price 5 cents.
67. The Relations of the Traps of the Newark System in the New Jersey Region, by Nelson Horatio Darton. 1890. 8°. 82 pp. Price 10 cents.
68. Earthquakes in California in 1889, by James Edward Keeler. 1890. 8°. 25 pp. Price 5 cents.
69. A Classified and Annotated Bibliography of Fossil Insects, by Samuel Hubbard Scudder. 1890. 8°. 101 pp. Price 15 cents.
70. Report on Astronomical Work of 1889 and 1890, by Robert Simpson Woodward. 1890. 8°. 79 pp. Price 10 cents.
71. Index to the Known Fossil Insects of the World, including Myriapods and Arachnids, by Samuel Hubbard Scudder. 1891. 8°. 744 pp. Price 50 cents.
72. Altitudes between Lake Superior and the Rocky Mountains, by Warren Upham. 1891. 8°. 229 pp. Price 20 cents.
73. The Viscosity of Solids, by Carl Barus. 1891. 8°. xii, 139 pp. 6 pl. Price 15 cents.
74. The Minerals of North Carolina, by Frederick Augustus Genth. 1891. 8°. 119 pp. Price 15 cents.
75. Record of North American Geology for 1887 to 1889, inclusive, by Nelson Horatio Darton. 1891. 8°. 173 pp. Price 15 cents.
76. A Dictionary of Altitudes in the United States (Second Edition), compiled by Henry Gannett, Chief Topographer. 1891. 8°. 393 pp. Price 25 cents.
77. The Texan Permian and its Mesozoic Types of Fossils, by Charles A. White. 1891. 8°. 51 pp. 4 pl. Price 10 cents.
78. A Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1889-'90. F. W. Clarke, Chief Chemist. 1891. 8°. 131 pp. Price 15 cents.
79. A Late Volcanic Eruption in Northern California and its Peculiar Lava, by J. S. Diller. 1891. 8°. 33 pp. 17 pl. Price 10 cents.
80. Correlation Papers—Devonian and Carboniferous, by Henry Shaler Williams. 1891. 8°. 279 pp. Price 20 cents.
81. Correlation Papers—Cambrian, by Charles Doolittle Walcott. 1891. 8°. 447 pp. 3 pl. Price 25 cents.
82. Correlation Papers—Cretaceous, by Charles A. White. 1891. 8°. 273 pp. 3 pl. Price 20 cents.
83. Correlation Papers—Eocene, by William Bullock Clark. 1891. 8°. 173 pp. 2 pl. Price 15 cents.

84. Correlation Papers—Neocene, by W. H. Dall and G. D. Harris. 1892. 8°. 349 pp. 3 pl. Price 25 cents.
85. Correlation Papers—The Newark System, by Israel Cook Russell. 1892. 8°. 344 pp. 13 pl. Price 25 cents.
86. Correlation Papers—Archean and Algonkian, by C. R. Van Hise. 1892. 8°. 549 pp. 12 pl. Price 25 cents.
87. A Synopsis of American Fossil Brachiopoda, including Bibliography and Synonymy, by Charles Schuchert. 1897. 8°. 464 pp. Price 25 cents.
88. The Cretaceous Foraminifera of New Jersey, by Rufus Mather Bagg, jr. 1898. 8°. 89 pp. 6 pl. Price 10 cents.
89. Some Lava Flows of the Western Slope of the Sierra Nevada, California, by F. Leslie Ransome. 1898. 8°. 74 pp. 11 pl. Price 15 cents.
90. A Report of Work done in the Division of Chemistry and Physics, mainly during the Fiscal Year 1890-'91. F. W. Clarke, Chief Chemist. 1892. 8°. 77 pp. Price 10 cents.
91. Record of North American Geology for 1890, by Nelson Horatio Darton. 1891. 8°. 88 pp. Price 10 cents.
92. The Compressibility of Liquids, by Carl Barus. 1892. 8°. 96 pp. 29 pl. Price 10 cents.
93. Some Insects of Special Interest from Florissant, Colorado, and Other Points in the Tertiaries of Colorado and Utah, by Samuel Hubbard Scudder. 1892. 8°. 35 pp. 3 pl. Price 5 cents.
94. The Mechanism of Solid Viscosity, by Carl Barus. 1892. 8°. 138 pp. Price 15 cents.
95. Earthquakes in California in 1890 and 1891, by Edward Singleton Holden. 1892. 8°. 31 pp. Price 5 cents.
96. The Volume Thermodynamics of Liquids, by Carl Barus. 1892. 8°. 100 pp. Price 10 cents.
97. The Mesozoic Echinodermata of the United States, by William Bullock Clark. 1893. 8°. 207 pp. 50 pl. Price 20 cents.
98. Flora of the Outlying Carboniferous Basins of Southwestern Missouri, by David White. 1893. 8°. 139 pp. 5 pl. Price 15 cents.
99. Record of North American Geology for 1891, by Nelson Horatio Darton. 1892. 8°. 73 pp. Price 10 cents.
100. Bibliography and Index of the Publications of the U. S. Geological Survey, 1879-1892, by Philip Creveling Warman. 1893. 8°. 495 pp. Price 25 cents.
101. Insect Fauna of the Rhode Island Coal Field, by Samuel Hubbard Scudder. 1893. 8°. 27 pp. 2 pl. Price 5 cents.
102. A Catalogue and Bibliography of North American Mesozoic Invertebrata, by Cornelius Breckinridge Boyle. 1893. 8°. 315 pp. Price 25 cents.
103. High Temperature Work in Igneous Fusion and Ebullition, chiefly in Relation to Pressure, by Carl Barus. 1893. 8°. 57 pp. 9 pl. Price 10 cents.
104. Glaciation of the Yellowstone Valley north of the Park, by Walter Harvey Weed. 1893. 8°. 41 pp. 4 pl. Price 5 cents.
105. The Laramie and the Overlying Livingston Formation in Montana, by Walter Harvey Weed, with Report on Flora, by Frank Hall Knowlton. 1893. 8°. 68 pp. 6 pl. Price 10 cents.
106. The Colorado Formation and its Invertebrate Fauna, by T. W. Stanton. 1893. 8°. 288 pp. 45 pl. Price 20 cents.
107. The Trap Dikes of the Lake Champlain Region, by James Furman Kemp and Vernon Freeman Marsters. 1893. 8°. 62 pp. 4 pl. Price 10 cents.
108. A Geological Reconnaissance in Central Washington, by Israel Cook Russell. 1893. 8°. 108 pp. 12 pl. Price 15 cents.
109. The Eruptive and Sedimentary Rocks on Pigeon Point, Minnesota, and their Contact Phenomena, by William Shirley Bayley. 1893. 8°. 121 pp. 16 pl. Price 15 cents.
110. The Paleozoic Section in the Vicinity of Three Forks, Montana, by Albert Charles Peale. 1893. 8°. 56 pp. 6 pl. Price 10 cents.
111. Geology of the Big Stone Gap Coal Field of Virginia and Kentucky, by Marius R. Campbell. 1893. 8°. 106 pp. 6 pl. Price 15 cents.
112. Earthquakes in California in 1892, by Charles D. Perrine. 1893. 8°. 57 pp. Price 10 cents.
113. A Report of Work done in the Division of Chemistry during the Fiscal Years 1891-'92 and 1892-'93. F. W. Clarke, Chief Chemist. 1893. 8°. 115 pp. Price 15 cents.
114. Earthquakes in California in 1893, by Charles D. Perrine. 1894. 8°. 23 pp. Price 5 cents.
115. A Geographic Dictionary of Rhode Island, by Henry Gannett. 1894. 8°. 31 pp. Price 5 cents.
116. A Geographic Dictionary of Massachusetts, by Henry Gannett. 1894. 8°. 126 pp. Price 15 cents.
117. A Geographic Dictionary of Connecticut, by Henry Gannett. 1894. 8°. 67 pp. Price 10 cents.
118. A Geographic Dictionary of New Jersey, by Henry Gannett. 1894. 8°. 131 pp. Price 15 cents.
119. A Geological Reconnaissance in Northwest Wyoming, by George Homans Eldridge. 1894. 8°. 72 pp. 4 pl. Price 10 cents.
120. The Devonian System of Eastern Pennsylvania and New York, by Charles S. Prosser. 1895. 8°. 81 pp. 2 pl. Price 10 cents.

121. A Bibliography of North American Paleontology, by Charles Rollin Keyes. 1894. 8°. 251 pp. Price 20 cents.
122. Results of Primary Triangulation, by Henry Gannett. 1894. 8°. 412 pp. 17 pl. Price 25 cents.
123. A Dictionary of Geographic Positions, by Henry Gannett. 1895. 8°. 183 pp. 1 pl. Price 15 cents.
124. Revision of North American Fossil Cockroaches, by Samuel Hubbard Seudder. 1895. 8°. 176 pp. 12 pl. Price 15 cents.
125. The Constitution of the Silicates, by Frank Wigglesworth Clarke. 1895. 8°. 109 pp. Price 15 cents.
126. A Mineralogical Lexicon of Franklin, Hampshire, and Hampden Counties, Massachusetts, by Benjamin Kendall Emerson. 1895. 8°. 180 pp. 1 pl. Price 15 cents.
127. Catalogue and Index of Contributions to North American Geology, 1732-1891, by Nelson Horatio Darton. 1896. 8°. 1045 pp. Price 60 cents.
128. The Bear River Formation and its Characteristic Fauna, by Charles A. White. 1895. 8°. 108 pp. 11 pl. Price 15 cents.
129. Earthquakes in California in 1894, by Charles D. Perrine. 1895. 8°. 25 pp. Price 5 cents.
130. Bibliography and Index of North American Geology, Paleontology, Petrology, and Mineralogy for 1892 and 1893, by Fred Boughton Weeks. 1896. 8°. 210 pp. Price 20 cents.
131. Report of Progress of the Division of Hydrography for the Calendar Years 1893 and 1894, by Frederick Haynes Newell, Topographer in Charge. 1895. 8°. 126 pp. Price 15 cents.
132. The Disseminated Lead Ores of Southeastern Missouri, by Arthur Winslow. 1896. 8°. 31 pp. Price 5 cents.
133. Contributions to the Cretaceous Paleontology of the Pacific Coast: The Fauna of the Knoxville Beds, by T. W. Stanton. 1895. 8°. 132 pp. 20 pl. Price 15 cents.
134. The Cambrian Rocks of Pennsylvania, by Charles Doolittle Walcott. 1896. 8°. 43 pp. 15 pl. Price 15 cents.
135. Bibliography and Index of North American Geology, Paleontology, Petrology, and Mineralogy for the Year 1894, by F. B. Weeks. 1896. 8°. 141 pp. Price 15 cents.
136. Volcanic Rocks of South Mountain, Pennsylvania, by Florence Baseom. 1896. 8°. 124 pp. 28 pl. Price 15 cents.
137. The Geology of the Fort Riley Military Reservation and Vicinity, Kansas, by Robert Hay. 1896. 8°. 35 pp. 8 pl. Price 5 cents.
138. Artesian-Well Prospects in the Atlantic Coastal Plain Region, by N. H. Darton. 1896. 8°. 228 pp. 19 pl. Price 20 cents.
139. Geology of the Castle Mountain Mining District, Montana, by W. H. Weed and L. V. Pirsson. 1896. 8°. 164 pp. 17 pl. Price 15 cents.
140. Report of Progress of the Division of Hydrography for the Calendar Year 1895, by Frederick Haynes Newell, Hydrographer in Charge. 1896. 8°. 356 pp. Price 25 cents.
141. The Eocene Deposits of the Middle Atlantic Slope in Delaware, Maryland, and Virginia, by William Bullock Clark. 1896. 8°. 167 pp. 40 pl. Price 15 cents.
142. A Brief Contribution to the Geology and Paleontology of Northwestern Louisiana, by T. Wayland Vaughan. 1896. 8°. 65 pp. 4 pl. Price 10 cents.
143. A Bibliography of Clays and the Ceramic Arts, by John C. Branner. 1896. 8°. 114 pp. Price 15 cents.
144. The Moraines of the Missouri Coteau and their Attendant Deposits, by James Edward Todd. 1896. 8°. 71 pp. 21 pl. Price 10 cents.
145. The Potomac Formation in Virginia, by W. M. Fontaine. 1896. 8°. 149 pp. 2 pl. Price 15 cents.
146. Bibliography and Index of North American Geology, Paleontology, Petrology, and Mineralogy for the Year 1895, by F. B. Weeks. 1896. 8°. 130 pp. Price 15 cents.
147. Earthquakes in California in 1895, by Charles D. Perrine, Assistant Astronomer in Charge of Earthquake Observations at the Lick Observatory. 1896. 8°. 23 pp. Price 5 cents.
148. Analyses of Rocks, with a Chapter on Analytical Methods, Laboratory of the United States Geological Survey, 1880 to 1896, by F. W. Clarke and W. F. Hillebrand. 1897. 8°. 306 pp. Price 20 cents.
149. Bibliography and Index of North American Geology, Paleontology, Petrology, and Mineralogy for the Year 1896, by Fred Boughton Weeks. 1897. 8°. 152 pp. Price 15 cents.
150. The Educational Series of Rock Specimens Collected and Distributed by the United States Geological Survey, by Joseph Silas Diller. 1898. 8°. 400 pp. 47 pl. Price 25 cents.
151. The Lower Cretaceous Gryphaeas of the Texas Region, by R. T. Hill and T. Wayland Vaughan. 1898. 8°. 139 pp. 35 pl. Price 15 cents.
152. A Catalogue of the Cretaceous and Tertiary Plants of North America, by F. H. Knowlton. 1898. 8°. 247 pp. Price 20 cents.
153. A Bibliographic Index of North American Carboniferous Invertebrates, by Stuart Weller. 1898. 8°. 653 pp. Price 35 cents.
154. A Gazetteer of Kansas, by Henry Gannett. 1898. 8°. 246 pp. 6 pl. Price 20 cents.

155. Earthquakes in California in 1896 and 1897, by Charles D. Perrine, Assistant Astronomer in Charge of Earthquake Observations at the Lick Observatory. 1898. 8°. 47 pp. Price 5 cents.
156. Bibliography and Index of North American Geology, Paleontology, Petrology, and Mineralogy for the Year 1897, by Fred Boughton Weeks. 1898. 8°. 130 pp. Price 15 cents.
157. The Gneisses, Gabbro-Schists, and Associated Rocks of Southwestern Minnesota, by Christopher Webber Hall. 1899. 8°. 160 pp. 27 pl. Price 45 cents.
158. The Moraines of Southeastern South Dakota and their Attendant Deposits, by James Edward Todd. 1899. 8°. 171 pp. 27 pl. Price 25 cents.
159. The Geology of Eastern Berkshire County, Massachusetts, by B. K. Emerson. 1899. 8°. 139 pp. 9 pl. Price 20 cents.
160. A Dictionary of Altitudes in the United States (Third Edition), compiled by Henry Gannett. 1899. 8°. 775 pp. Price 40 cents.
161. Earthquakes in California in 1898, by Charles D. Perrine, Assistant Astronomer in Charge of Earthquake Observations at the Lick Observatory. 1899. 8°. 31 pp. 1 pl. Price 5 cents.
162. Bibliography and Index of North American Geology, Paleontology, Petrology, and Mineralogy for the Year 1898, by Fred Boughton Weeks. 1899. 8°. 163 pp. Price 15 cents.
163. Flora of the Montana Formation, by Frank Hall Knowlton. 1900. 8°. 118 pp. Price 15 cents.
164. Reconnaissance in the Rio Grande Coal Fields of Texas, by Thomas Wayland Vaughan, including a Report on Igneous Rocks from the San Carlos Coal Field, by E. C. E. Lord. 1900. 8°. 100 pp. 11 pl. and maps. Price 20 cents.
165. Contributions to the Geology of Maine, by Henry S. Williams and Herbert E. Gregory. 1900. 8°. 212 pp. 14 pl. Price 25 cents.
166. A Gazetteer of Utah, by Henry Gannett. 1900. 8°. 43 pp. 1 map. Price 15 cents.
167. Contributions to Chemistry and Mineralogy from the Laboratory of the United States Geological Survey, Frank W. Clarke, Chief Chemist. 1900. 8°. 166 pp. Price 15 cents.
168. Analyses of Rocks, Laboratory of the United States Geological Survey, 1880 to 1899, tabulated by F. W. Clarke, Chief Chemist. 1900. 8°. 308 pp. Price 20 cents.
169. Altitudes in Alaska, by Henry Gannett. 1900. 8°. 13 pp. Price 5 cents.

In preparation:

170. Idaho-Montana Boundary Line, by Richard Urquhart Goode.
- Bibliography and Catalogue of the Fossil Vertebrata of North America, by Oliver Perry Hay.

WATER-SUPPLY AND IRRIGATION PAPERS.

By act of Congress approved June 11, 1896, the following provision was made:

"*Provided*, That hereafter the reports of the Geological Survey in relation to the gauging of streams and to the methods of utilizing the water resources may be printed in octavo form, not to exceed one hundred pages in length and five thousand copies in number; one thousand copies of which shall be for the official use of the Geological Survey, one thousand five hundred copies shall be delivered to the Senate, and two thousand five hundred copies shall be delivered to the House of Representatives, for distribution."

Under this law the following papers have been published:

1. Pumping Water for Irrigation, by Herbert M. Wilson. 1896. 8°. 57 pp. 9 pl.
2. Irrigation near Phoenix, Arizona, by Arthur P. Davis. 1897. 8°. 97 pp. 31 pl.
3. Sewage Irrigation, by George W. Rafter. 1897. 8°. 100 pp. 4 pl.
4. A Reconnaissance in Southeastern Washington, by Israel Cook Russell. 1897. 8°. 96 pp. 7 pl.
5. Irrigation Practice on the Great Plains, by Elias Branson Cowgill. 1897. 8°. 39 pp. 12 pl.
6. Underground Waters of Southwestern Kansas, by Erasmus Haworth. 1897. 8°. 65 pp. 12 pl.
7. Seepage Waters of Northern Utah, by Samuel Fortier. 1897. 8°. 50 pp. 3 pl.
8. Windmills for Irrigation, by E. C. Murphy. 1897. 8°. 49 pp. 8 pl.
9. Irrigation near Greeley, Colorado, by David Boyd. 1897. 8°. 90 pp. 21 pl.
10. Irrigation in Mesilla Valley, New Mexico, by F. C. Barker. 1898. 8°. 51 pp. 11 pl.
11. River Heights for 1896, by Arthur P. Davis. 1897. 8°. 100 pp.
12. Underground waters of Southeastern Nebraska, by N. H. Darton. 1898. 8°. 56 pp. 21 pl.
13. Irrigation Systems in Texas, by William Ferguson Hutson. 1898. 8°. 67 pp. 10 pl.
14. New Tests of Pumps and Water-Lifts used in Irrigation, by O. P. Hood. 1898. 8°. 91 pp. 1 pl.
15. Operations at River Stations, 1897, Part I. 1898. 8°. 100 pp.
16. Operations at River Stations, 1897, Part II. 1898. 8°. 101-200 pp.
17. Irrigation near Bakersfield, California, by C. E. Grunsky. 1898. 8°. 96 pp. 16 pl.
18. Irrigation near Fresno, California, by C. E. Grunsky. 1898. 8°. 94 pp. 14 pl.
19. Irrigation near Merced, California, by C. E. Grunsky. 1899. 8°. 59 pp. 11 pl.
20. Experiments with Windmills, by T. O. Perry. 1899. 8°. 97 pp. 12 pl.
21. Wells of Northern Indiana, by Frank Leverett. 1899. 8°. 82 pp. 2 pl.
22. Sewage Irrigation, Part II, by George W. Rafter. 1899. 8°. 100 pp. 7 pl.
23. Water-right Problems of the Bighorn Mountains, by Elwood Mead. 1899. 8°. 62 pp. 7 pl.
24. Water Resources of the State of New York, Part I, by G. W. Rafter. 1899. 8°. 99 pp. 13 pl.

25. Water Resources of the State of New York, Part II, by G. W. Rafter. 1899. 8°. 101-200 pp. 12 pl.
26. Wells of Southern Indiana (Continuation of No. 21), by Frank Leverett. 1899. 8°. 64 pp.
27. Operations at River Stations for 1898, Part I. 1899. 8°. 100 pp.
28. Operations at River Stations for 1898, Part II. 1899. 8°. 101-200 pp.
29. Wells and Windmills in Nebraska, by Erwin H. Barbour. 1899. 8°. 85 pp. 27 pl.
30. Water Resources of the Lower Peninsula of Michigan, by Alfred C. Lane. 1899. 8°. 97 pp. 7 pl.
31. Lower Michigan Mineral Waters, by Alfred C. Lane. 1899. 8°. 97 pp. 4 pl.
32. Water Resources of Puerto Rico, by Herbert M. Wilson. 1899. 8°. 48 pp. 17 pl.
33. Storage of Water on Gila River, Arizona, by Joseph B. Lippincott. 1900. 8°. 98 pp. 33 pl.
34. Geology and water resources of SE. South Dakota, by J. E. Todd. 1900. 8°. 34 pp. 19 pls.

In preparation:

35. Operations at River Stations, 1899, Part I.
36. Operations at River Stations, 1899, Part II.
37. Operations at River Stations, 1899, Part III.

TOPOGRAPHIC MAP OF THE UNITED STATES.

When, in 1882, the Geological Survey was directed by law to make a geologic map of the United States, there was in existence no suitable topographic map to serve as a base for the geologic map. The preparation of such a topographic map was therefore immediately begun. About one-fifth of the area of the country, excluding Alaska, has now been thus mapped. The map is published in atlas sheets, each sheet representing a small quadrangular district, as explained under the next heading. The separate sheets are sold at 5 cents each when fewer than 100 copies are purchased, but when they are ordered in lots of 100 or more copies, whether of the same sheet or of different sheets, the price is 2 cents each. The mapped areas are widely scattered, nearly every State being represented. About 900 sheets have been engraved and printed; they are tabulated by States in the Survey's "List of Publications," a pamphlet which may be had on application.

The map sheets represent a great variety of topographic features, and with the aid of descriptive text they can be used to illustrate topographic forms. This has led to the projection of an educational series of topographic folios, for use wherever geography is taught in high schools, academies, and colleges. Of this series the first two folios have been issued, viz:

1. Physiographic types, by Henry Gannett, 1898, folio, consisting of the following sheets and 4 pages of descriptive text: Fargo (N. Dak.-Minn.), a region in youth; Charleston (W. Va.), a region in maturity; Caldwell (Kans.), a region in old age; Palmyra (Va.), a rejuvenated region; Mount Shasta (Cal.), a young volcanic mountain; Eagle (Wis.), moraines; Sun Prairie (Wis.), drumlins; Donaldsonville (La.), river flood plains; Boothbay (Me.), a fiord coast; Atlantic City (N. J.), a barrier-beach coast.

2. Physiographic types, by Henry Gannett, 1900, folio, consisting of the following sheets and 11 pages of descriptive text: Norfolk (Va.-N. C.), a coast swamp; Marshall (Mo.), a graded river; Lexington (Nebr.), an overloaded stream; Harrisburg (Pa.), Appalachian ridges; Poteau Mountain (Ark.-Ind. T.), Ozark ridges; Marshall (Ark.), Ozark Plateau; West Denver (Colo.), hogbacks; Mount Taylor (N. Mex.), volcanic peaks, plateaus, and necks; Cucamonga (Cal.), alluvial cones; Crater Lake special (Oreg.), a crater.

GEOLOGIC ATLAS OF THE UNITED STATES.

The Geologic Atlas of the United States is the final form of publication of the topographic and geologic maps. The atlas is issued in parts, or folios, progressively as the surveys are extended, and is designed ultimately to cover the entire country.

Under the plan adopted the entire area of the country is divided into small rectangular districts (designated *quadrangles*), bounded by certain meridians and parallels. The unit of survey is also the unit of publication, and the maps and descriptions of each rectangular district are issued as a folio of the Geologic Atlas.

Each folio contains topographic, geologic, economic, and structural maps, together with textual descriptions and explanations, and is designated by the name of a principal town or of a prominent natural feature within the district.

Two forms of issue have been adopted, a "library edition" and a "field edition." In both the sheets are bound between heavy paper covers, but the library copies are permanently bound, while the sheets and covers of the field copies are only temporarily wired together.

Under the law a copy of each folio is sent to certain public libraries and educational institutions. The remainder are sold at 25 cents each, except such as contain an unusual amount of matter, which are priced accordingly. Prepayment is obligatory. The folios ready for distribution are here listed.

No.	Name of sheet.	State.	Limiting meridians.	Limiting parallels.	Area, in square miles.	Price, in cents.		
1	Livingston	Montana..	110°-111°	45°-46°	3,354	25		
2	Ringgold	Georgia... Tennessee }	85°-85° 30'	34° 30'-35°	980	25		
3	Placerville.....	California	120° 30'-121°	38° 30'-39°	932	25		
4	Kingston <i>a</i>	Tennessee	84° 30'-85°	35° 30'-36°	969	25		
5	Sacramento.....	California	121°-121° 30'	38° 30'-39°	932	25		
6	Chattanooga.....	Tennessee	85°-85° 30'	35°-35° 30'	975	25		
7	Pikes Peak <i>a</i>	Colorado..	105°-105° 30'	38° 30'-39°	932	25		
8	Sewanee	Tennessee	85° 30'-86°	35°-35° 30'	975	25		
9	Anthracite-Crest- ed Butte.	Colorado.. Virginia.. }	106° 45'-107° 15'	38° 45'-39°	465	50		
10	Harpers Ferry ..	West Va.. }	77° 30'-78°	39°-39° 30'	925	25		
11	Jackson	Maryland. California }	120° 30'-121°	38°-38° 30'	938	25		
12	Estillville	Virginia.. Kentucky }	82° 30'-83°	36° 30'-37°	957	25		
13	Fredericksburg..	Tennessee Maryland. Virginia.. }	77°-77° 30'	38°-38° 30'	938	25		
14	Staunton	Virginia.. West Va.. }	79°-79° 30'	38°-38° 30'	938	25		
15	Lassen Peak.....	California.	121°-122°	40°-41°	3,634	25		
16	Knoxville.....	Tennessee N.Carolina }	83° 30'-84°	35° 30'-36°	925	25		
17	Marysville.....	California	121° 30'-122°	39°-39° 30'	925	25		
18	Smartsville.....	California	121°-121° 30'	39°-39° 30'	925	25		
19	Stevenson	Alabama.. Georgia... Tennessee }	85° 30'-86°	34° 30'-35°	980	25		
20	Cleveland.....	Tennessee	84° 30'-85°	35°-35° 30'	975	25		
21	Pikeville.....	Tennessee	85°-85° 30'	35° 30'-36°	969	25		
22	McMinnville.....	Tennessee	85° 30'-86°	35° 30'-36°	969	25		
23	Nomini	Maryland. Virginia.. }	76° 30'-77°	38°-38° 30'	938	25		
24	Three Forks.....	Montana..	111°-112°	45°-46°	3,354	50		
25	Loudon	Tennessee	84°-84° 30'	35° 30'-36°	969	25		
26	Pocahontas	Virginia.. West Va.. }	81°-81° 30'	37°-37° 30'	951	25		
27	Morristown.....	Tennessee	83°-83° 30'		963	25		
28	Piedmont.....	Maryland. West Va.. }	79°-79° 30'	39°-39° 30'	925	25		
29	Nevada City: Nevada City. Grass Valley. Banner Hill..	California.	{ 121° 00' 25"-121° 03' 45" 121° 01' 35"-121° 05' 04" 120° 57' 05"-121° 00' 25"	{ 39° 13' 50"-39° 17' 16" 39° 10' 22"-39° 13' 50" 39° 13' 50"-39° 17' 16"	{ 11.65 12.09 11.65	50		
30	Yellowstone Na- tional Park: Gallatin		Wyoming.	110°-111°	44°-45°		3,412	75
	Canyon							
	Shoshone							
	Lake							
31	Pyramid Peak.....	California.	120°-120° 30'	38° 30'-39°	932	25		
32	Franklin	Virginia.. West Va.. }	79°-79° 30'	38° 30'-39°	932	25		
33	Briceville.....	Tennessee	84°-84° 30'	36°-36° 30'	963	25		
34	Buckhannon.....	West Va..	80°-80° 30'	38° 30'-39°	932	25		
35	Gadsden	Alabama..	86°-86° 30'	34°-34° 30'	986	25		
36	Pueblo	Colorado..	104° 30'-105°	38°-38° 30'	938	50		
37	Downieville	California.	120° 30'-121°	39° 30'-40°	919	25		
38	Butte Special	Montana..	112° 29' 30"-112° 36' 42"	45° 59' 28"-46° 02' 54"	22.80	50		
39	Truckee	California.	120°-120° 30'	39°-39° 30'	925	25		
40	Wartburg	Tennessee	84° 30'-85°	36°-36° 30'	963	25		
41	Sonora	California.	120°-120° 30'	37° 30'-38°	944	25		
42	Nueces	Texas	100°-100° 30'	29° 30'-30°	1,035	25		
43	Bidwell Bar	California.	121°-121° 30'	39° 30'-40°	918	25		
44	Tazewell	Virginia.. West Va.. }	81° 30'-82°	37°-37° 30'	950	25		
45	Boise	Idaho	116°-116° 30'	43° 30'-44°	864	25		
46	Richmond	Kentucky	84°-84° 30'	37° 30'-38°	944	25		
47	London	Kentucky	84°-84° 30'	37°-37° 30'	950	25		
48	Tennile District Special.	Colorado..	106° 8'-106° 16'	39° 22' 30"-39° 30' 30"	55	25		
49	Roseburg	Oregon ...	123°-123° 30'	43°-43° 30'	871	25		
50	Holyoke	Mass	72° 30'-73°	42°-42° 30'	885	50		
51	Big Trees.....	Conn						
52	Absaroka.....	California	120°-120° 30'	38°-38° 30'	938	25		
	Crandall	Wyoming.	109° 30'-110°	44°-44° 30'	1,706	25		
	Ishawooa							
53	Standingstone	Tennessee	85°-85° 30'	36°-36° 30'	963	25		

a Out of stock.

No.	Name of sheet.	State.	Limiting meridians.	Limiting parallels.	Area, in square miles.	Price, in cents.
54	Tacoma.....	Washing- ton.	122°-122° 30'	47°-47° 30'	812	25
55	Fort Benton.....	Montana..	110°-111°	47°-48°	3, 273	25
56	Little Belt Mts....	Montana..	110°-111°	46°-47°	3, 295	25
57	Telluride.....	Colorado..	107° 45'-108°	37° 45'-38°	236	25
58	Elmoro.....	Colorado..	104°-104° 30'	37°-37° 30'	950	25

STATISTICAL PAPERS.

Mineral Resources of the United States, 1882, by Albert Williams, jr. 1883. 8°. xvii, 813 pp. Price 50 cents.

Mineral Resources of the United States, 1883 and 1884, by Albert Williams, jr. 1885. 8°. xiv, 1016 pp. Price 60 cents.

Mineral Resources of the United States, 1885. Division of Mining Statistics and Technology. 1886. 8°. vii, 576 pp. Price 40 cents.

Mineral Resources of the United States, 1886, by David T. Day. 1887. 8°. viii, 813 pp. Price 50 cents.

Mineral Resources of the United States, 1887, by David T. Day. 1888. 8°. vii, 832 pp. Price 50 cents.

Mineral Resources of the United States, 1888, by David T. Day. 1890. 8°. vii, 652 pp. Price 50 cents.

Mineral Resources of the United States, 1889 and 1890, by David T. Day. 1892. 8°. viii, 671 pp. Price 50 cents.

Mineral Resources of the United States, 1891, by David T. Day. 1893. 8°. vii, 630 pp. Price 50 cents.

Mineral Resources of the United States, 1892, by David T. Day. 1893. 8°. vii, 850 pp. Price 50 cents.

Mineral Resources of the United States, 1893, by David T. Day. 1894. 8°. viii, 810 pp. Price 50 cents.

On March 2, 1895, the following provision was included in an act of Congress:

"*Provided*, That hereafter the report of the mineral resources of the United States shall be issued, as a part of the report of the Director of the Geological Survey."

In compliance with this legislation the following reports have been published:

Mineral Resources of the United States, 1894, David T. Day, Chief of Division. 1895. 8°. xv, 646 pp., 23 pl.; xix, 735 pp., 6 pl. Being Parts III and IV of the Sixteenth Annual Report.

Mineral Resources of the United States, 1895, David T. Day, Chief of Division. 1896. 8°. xxiii, 542 pp., 8 pl. and maps; iii, 543-1058 pp., 9-13 pl. Being Part III (in 2 vols.) of the Seventeenth Annual Report.

Mineral Resources of the United States, 1896, David T. Day, Chief of Division. 1897. 8°. xii, 642 pp., 1 pl.; 643-1400 pp. Being Part V (in 2 vols.) of the Eighteenth Annual Report.

Mineral Resources of the United States, 1897, David T. Day, Chief of Division. 1898. 8°. viii, 651 pp., 11 pl.; viii, 706 pp. Being Part VI (in 2 vols.) of the Nineteenth Annual Report.

Mineral Resources of the United States, 1898, David T. Day, Chief of Division. 1899. 8°. viii, 616 pp.; ix, 804 pp., 1 pl. Being Part VI (in 2 vols.) of the Twentieth Annual Report.

The money received from the sale of the Survey publications is deposited in the Treasury, and the Secretary of the Treasury declines to receive bank checks, drafts, or postage stamps; all remittances, therefore, must be by MONEY ORDER, made payable to the Director of the United States Geological Survey, or in CURRENCY—the exact amount. Correspondence relating to the publications of the Survey should be addressed to—

THE DIRECTOR,

UNITED STATES GEOLOGICAL SURVEY,

WASHINGTON, D. C.

WASHINGTON, D. C., May, 1900.



